



- ▶ Age : 15 years old
- ▶ Reason of consultation
/ Chief complaint
/ main request : TMJ disorder
- ▶ Clinical Exam: TMD




How to prevent/correct TMJ disorder after orthodontic treatment?

** Modjaw en la práctica ortodóncica diaria: a proposito de un caso tratado con alineadores FAS. Martin D, Canabaz A, Aznar J, Fa*



Functional DC (CR/ first contact)



Mandibular position /
Selection and transfer



TM disorder

Why MODJAW? Select mandibular position, reeducate and target new therapeutic endpoint

CLINICAL CASE

SITUATION

CHALLENGE

IMPLICATION

SOLUTION



Patient dissatisfaction




Chair time



Profitability



Appointment #1




**Optimal / Complete patient data acquisition
(TMJ disorder diagnostic – optimized
therapeutic position)**



**Ability to calculate an appropriately lower
jaw position to fit the splint and stabilize the
patient in the right position**



Appointment #2



Functional evaluation after Splint therapy

CLINICAL CASE – Splint therapy

SITUATION


CHALLENGE

IMPLICATION

SOLUTION



Appointment #2



**Optimal/ Complete patient data acquisition
(Optimized therapeutic position after splint
therapy – Functional envelope)**

Lab work



**Control, reduce the room for assumptions
(Align the teeth starting from the optimized
therapeutic position)**




Appointment #3





Appointment #3



**High level of predictable functional
and aesthetic outcomes
(No more TMJ disorder – stable mandibular
position – correct functional envelope)**

**Optimal/ Complete patient data acquisition
(Optimized therapeutic position after splint
therapy – Functional envelope)**



Patient satisfaction

Excellent ortho result aesthetic and functional with no pain



Predictable clinical outcomes

Stable mandibular position – correct functional envelope



Clinical profitability

No post treatment complication – stable treatment – less chair time