PHYS1401 Midterm#2 (105/100 pts)

NOTE(IMPORTANT!!): Please detach, sign, and submit the equation sheet to receive credit.

Print Name:_____

Student ID:_____ **Course Section:**_____

Part I: Multiple-choice Questions (Only the answer shall be credited. Each question has only one correct answer.)


1. (5pts) Consider an object doing a uniform circular motion. Which of the following statements is correct?
   A. The acceleration of the object is zero since the speed is constant.
   B. The acceleration changes the magnitude of the velocity.
   C. The acceleration of the object is non-zero since the velocity direction changes.
   D. None of the above is correct.

2. (5pts) A 1500 kg fridge is on a horizontal kitchen floor with a coefficient of static friction $\mu_s = 0.44$ and a coefficient of kinetic friction $\mu_k = 0.35$. A horizontal force of 7500 N is applied to the fridge. What is the magnitude of the frictional force?
   A. 6500 N.
   B. 7500 N.
   C. 5100 N.
   D. 15000 N.


3. (5pts) A 74 kg person is standing on a scale in an elevator that is moving downward at a constant velocity of 1.3 m/s. What is the person's apparent weight?
   A. 0 N
   B. 820 N.
   C. 630 N.
   D. 730 N.

4. Given a swimmer's velocity relative to the river's current as $\vec{V}_{sr}$, the velocity of the river's current relative to the bank as $\vec{V}_{rb}$, and the swimmer's velocity relative to the bank as $\vec{V}_{sb}$, identify the correct vector tip-to-tail triangle representing the relationships between $\vec{V}_{sr}$, $\vec{V}_{rb}$ and $\vec{V}_{sb}$.


A




B



C



D



5. (5pts) A ball (mass 0.40 kg) is initially moving to the left at 20 m/s. After hitting the wall, the ball moves to the right at 20 m/s. Which of the following statements is correct?
   A. The momentum of the ball does not change during the collision with the wall.
   B. The impulse of the net force on the ball during the collision is 32 kg·m/s to the left.
   C. The impulse of the net force on the ball during the collision is 32 kg·m/s to the right
   D. The impulse of the net force on the ball during the collision is 0 kg·m/s.

6. (5pts) The electric force is given by the equation $\vec{F}_{e,12} = k\frac{q_1 q_2}{r_{12}^2}\hat{r}_{12}$ and the universal gravitational force is given by the equation $\vec{F}_{g,12} = -G\frac{m_1 m_2}{r_{12}^2}\hat{r}_{12}$ According to your understanding of these two equations, which of the following statements is correct?
   A. The direction of $\vec{F}_{g,12}$ is opposite to the direction of $\hat{r}_{12}$.
   B. The direction of $\vec{F}_{g,12}$ is always the same as the direction of $\hat{r}_{12}$
   C. The direction of $\vec{F}_{e,12}$ is opposite to the direction of $\hat{r}_{12}$.
   D. The direction of $\vec{F}_{e,12}$ is always the same as the direction of $\hat{r}_{12}$

7. (5pts) A passenger is standing on a scale in an elevator. The scale sits on the floor of the elevator. The elevator now begins to descend. Starting from rest, it takes the elevator 6 seconds to get up to its downward speed of 12 m/s. Which of the following statements is correct?




A. There are two forces exerted on the passenger: the gravitational force and the normal force from the scale.

B. The net force acting on the passenger is zero.

C. There are three forces exerted on the passenger: the gravitational force, the normal force from the scale and the normal force from the elevator.


D. The acceleration of the passenger is 9.8 m/s^2.

8. (5pts) Two boxes are stacked as shown below. According to Newton's third law, which of the following is a correct pair of action-reaction (Newton's Third Law) forces?



A. The normal force exerted by object #2 on object #1 and the gravitational force on object #1.
B. The gravitational force on object #1 and the downward force that object #1 exerts on object #2.
C. The normal force exerted by object #2 on object #1 and the gravitational force on object #2.
D. The normal force exerted by object #2 on object #1 and the downward force that object #1 exerts on object #2.

9. (5pts) What is the average force of the impulse in the following *F*-*t* plot?



A. 12N
B. 20N
C. 16N
D. 13N

10. (5pts) A passenger on a Ferris wheel of radius R = 20 m has a mass of 70 kg. At the top of the ride, the passenger has an apparent weight of 672 N. What is the passenger's apparent weight at the bottom of the ride?




A. 700 N
B. 704 N
C. 696 N
D. 692 N

Part II: Problem-solving Question (Please present the complete problem-solving procedure. Answers with no supportive calculation shall not be credited)

12. (25pts) A small box with mass 320 g slides down a ramp with initial velocity $0m/s$ as shown below. The ramp makes an angle $75°$ with the vertical direction and the coefficient of kinetic friction between the box and the ramp is 0.14. If the length down the ramp is a distance 2.1 m, what is the magnitude of the velocity when the box reaches the end of the ramp?


    a. (5pts) Please draw the free-body diagram for the box with **a proper coordinate system**.



    b. (20pts) This problem can be solved using kinematics, the work-energy theorem, or the principle of energy conservation. Choose any two of these three methods to solve the problem and verify that both approaches yield the same result.

13. (15pts) Train cars are coupled by colliding with each other. Suppose two loaded train cars are moving toward one another on a horizontal track: the first car has a mass of 211,000 kg and is traveling at 0.291 m/s to the right, while the second car has a mass of 63,500 kg and is moving at 0.122 m/s to the left. What are the magnitude and direction of their final velocity (in meters per second) after the collision? Using the five-step problem-solving approach, solve the problem.

14. (15pts) A ball of mass m is thrown vertically upward and reaches a maximum height h. Take the zero of gravitational potential energy to be at ground level. Assume air resistance is negligible. Using the given variables (g, m, h), write expressions for the following questions. Note: Not all variables may be needed.



a. (5pts) Derive an expression for the ball's initial speed.

b. (5pts) Derive an expression for the work done by the gravitational force as the ball moves from the ground to its maximum height.

c. (5pts) Derive an expression for the change in gravitational potential energy as the ball drops from the maximum height to a height of $h/2$.

Equation Sheet

Gravitational constant: $g = 9.8$ m/s^2
Universal gravitational constant: $G = 6.674 \times 10^{-11}$ N·m^2/kg^2

Coulomb's constant: $k = 8.99 \times 10^9 N \cdot m^2/C^2$

Newton's second law: $\sum \vec{F} = m\vec{a}$
Newton's third law: $\vec{F}_{AB} = -\vec{F}_{BA}$

The acceleration of uniform circular motion: $a = v^2/R$
The period of uniform circular motion: $T = \frac{2\pi r}{v}$
Frequency: $f = 1/T$

Weight: $F_g = mg$
Kinetic friction: $f = \mu_k N$
Static friction: $f \leq \mu_s N$
Universal gravitational force: $\vec{F}_{g,12} = -G\frac{m_1 m_2}{r_{12}^2}\hat{r}_{12}$
Electric force: $\vec{F}_{e,12} = k\frac{q_1 q_2}{r_{12}^2}\hat{r}_{12}$

Relative Motion: $\vec{V}_{A,B} = \vec{V}_{A,C} + \vec{V}_{C,B}$ and $\vec{V}_{A,B} = -\vec{V}_{B,A}$

Momentum: $\vec{p} = m\vec{v}$
Newton's second law in terms of momentum: $\sum \vec{F} = \frac{d\vec{p}}{dt}$
Average Force: $F_{avg} = J/\Delta t$
Impulse-momentum theorem: $\vec{J} = \Delta\vec{p}$
Momentum conservation: $\sum \vec{p}_i = \sum \vec{p}_f$

Work: $W = \vec{F} \cdot \vec{l}$ (constant force)
Kinetic energy: $K = \frac{1}{2}mv^2$
Work-energy theorem: $W_{tot} = \Delta K$

Gravitational potential energy: $\Delta U_{grav} = mg\Delta y$
$U_{grav} = -G\frac{m_1 m_2}{r_{12}}$

Mechanical energy conservation: $\Delta K + \Delta U = 0$
Energy conservation: $\Delta K + \Delta U + \Delta U_{loss} = 0$

Dot product: $\vec{A} \cdot \vec{B} = |\vec{A}||\vec{B}| \cos\phi$ or $\vec{A} \cdot \vec{B} = A_x B_x + A_y B_y + A_z B_z$