

## EC2104 Tutorial 5 solution

ling

## Section 1

### Question 1

# Question 1

Given  $\pi = p(Q) \cdot Q$ , find relationship between  $MR$  and  $\epsilon_D$

- What do I need? (1)  $MR$ , (2)  $\epsilon_D$ , (3) relationship between them

# Question 1

Given  $\pi = p(Q) \cdot Q$ , find relationship between  $MR$  and  $\epsilon_D$

- What do I need? (1)  $MR$ , (2)  $\epsilon_D$ , (3) relationship between them
- $MR: \frac{d\pi}{dQ} = P(Q) + Q \frac{dP}{dQ}$

# Question 1

Given  $\pi = p(Q) \cdot Q$ , find relationship between  $MR$  and  $\epsilon_D$

- What do I need? (1)  $MR$ , (2)  $\epsilon_D$ , (3) relationship between them
- $MR: \frac{d\pi}{dQ} = P(Q) + Q \frac{dP}{dQ}$
- $\epsilon_D = \frac{dQ}{dP} \left( \frac{P}{Q} \right)$

# Question 1

Given  $\pi = p(Q) \cdot Q$ , find relationship between  $MR$  and  $\epsilon_D$

- What do I need? (1)  $MR$ , (2)  $\epsilon_D$ , (3) relationship between them
- $MR: \frac{d\pi}{dQ} = P(Q) + Q \frac{dP}{dQ}$
- $\epsilon_D = \frac{dQ}{dP} \left( \frac{P}{Q} \right)$
- Relationship:  $Q \frac{dP}{dQ} = \frac{P}{\epsilon_D}$

# Question 1

Given  $\pi = p(Q) \cdot Q$ , find relationship between  $MR$  and  $\epsilon_D$

- What do I need? (1)  $MR$ , (2)  $\epsilon_D$ , (3) relationship between them
- $MR$ :  $\frac{d\pi}{dQ} = P(Q) + Q \frac{dP}{dQ}$
- $\epsilon_D = \frac{dQ}{dP} \left( \frac{P}{Q} \right)$
- Relationship:  $Q \frac{dP}{dQ} = \frac{P}{\epsilon_D}$
- Result:  $MR = P + Q \frac{dP}{dQ} = P + \frac{P}{\epsilon_D}$  (ans: a)

## Section 2

### Question 2



## Question 2

Given equation system, find  $\frac{du}{dv}$ ,  $\frac{du}{dy}$ ,  $\frac{dv}{dx}$ ,  $\frac{dv}{dy}$

$$\begin{aligned}\ln(x + u) + uv - y^2 e^y + y &= 0 \\ u^2 - x^v &= v\end{aligned}$$

- What do I need? (1) Solve differentials (2) Substitute values at the point (3) Solve system of linear equations (4) Find coefficients

## Question 2

Given equation system, find  $\frac{du}{dv}$ ,  $\frac{du}{dy}$ ,  $\frac{dv}{dx}$ ,  $\frac{dv}{dy}$

$$\begin{aligned}\ln(x + u) + uv - y^2 e^y + y &= 0 \\ u^2 - x^v &= v\end{aligned}$$

- What do I need? (1) Solve differentials (2) Substitute values at the point (3) Solve system of linear equations (4) Find coefficients
- Differentials:

$$\begin{aligned}\frac{dx + du}{x + u} + u dv + v du - 2ye^y dy + y^2 e^y dv + dy &= 0 \\ 2u du - vx^{v-1} dx - x^v \ln(x) dv &= dv\end{aligned}$$

## Question 2 Cont

- What do I need? (1) Solve differentials (2) Substitute values at the point (3) Solve system of linear equations (4) Find coefficients

## Question 2 Cont

- What do I need? (1) Solve differentials (2) Substitute values at the point (3) Solve system of linear equations (4) Find coefficients
- Substitute  $(x, y, u, v) = (2, 1, -1, 0)$

$$dx + du - dv - 2dy - dv + dy = 0 \Rightarrow du - 2dv = dy - dx$$

$$-2du - \ln(2)dv = dv \Rightarrow dv = -\frac{2}{1 + \ln(2)}du$$

$$\Rightarrow du = -\frac{1 + \ln(2)}{2}dv$$

## Question 2 Cont

- What do I need? (1) Solve differentials (2) Substitute values at the point (3) Solve system of linear equations (4) Find coefficients
- Substitute  $(x, y, u, v) = (2, 1, -1, 0)$

$$dx + du - dv - 2dy - dv + dy = 0 \Rightarrow du - 2dv = dy - dx$$

$$-2du - \ln(2)dv = dv \Rightarrow dv = -\frac{2}{1 + \ln(2)}du$$

$$\Rightarrow du = -\frac{1 + \ln(2)}{2}dv$$

- Solve:  $du - 2dv = dy - dx$

$$du + \frac{4}{1 + \ln(2)}du = dy - dx \Rightarrow du = \frac{1 + \ln(2)}{5 + \ln(2)}(dy - dx)$$

$$-\frac{1 + \ln(2)}{2}dv + 2dv = dy - dx \Rightarrow dv = -\frac{2}{5 + \ln(2)}(dy - dx)$$

## Question 2 Cont

- What do I need? (1) Solve differentials (2) Substitute values at the point (3) Solve system of linear equations (4) Find coefficients
- Substitute  $(x, y, u, v) = (2, 1, -1, 0)$

$$dx + du - dv - 2dy - dv + dy = 0 \Rightarrow du - 2dv = dy - dx$$

$$-2du - \ln(2)dv = dv \Rightarrow dv = -\frac{2}{1 + \ln(2)}du$$

$$\Rightarrow du = -\frac{1 + \ln(2)}{2}dv$$

- Solve:  $du - 2dv = dy - dx$

$$du + \frac{4}{1 + \ln(2)}du = dy - dx \Rightarrow du = \frac{1 + \ln(2)}{5 + \ln(2)}(dy - dx)$$

$$-\frac{1 + \ln(2)}{2}dv + 2dv = dy - dx \Rightarrow dv = -\frac{2}{5 + \ln(2)}(dy - dx)$$

- Find coefficients:

$$\frac{du}{dy} = \frac{1 + \ln(2)}{5 + \ln(2)}, \frac{du}{dx} = -\frac{1 + \ln(2)}{5 + \ln(2)}, \frac{dv}{dy} = -\frac{2}{5 + \ln(2)}, \frac{dv}{dx} = \frac{2}{5 + \ln(2)} \text{ (ans: a)}$$

## Section 3

Question 3, 4, 5

## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?



## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

- ①  $f$  is decreasing on  $[a, b]$

## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

- ①  $f$  is decreasing on  $[a, b]$ 
  - False,  $f'(x) > 0$ .  $f$  is increasing

## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

- 1  $f$  is decreasing on  $[a, b]$ 
  - False,  $f'(x) > 0$ .  $f$  is increasing
- 2  $f$  has no local extrema on  $(a, b)$  using  $f'(x) = 0$

## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

- 1  $f$  is decreasing on  $[a, b]$ 
  - False,  $f'(x) > 0$ .  $f$  is increasing
- 2  $f$  has no local extrema on  $(a, b)$  using  $f'(x) = 0$ 
  - True,  $f'(x) > 0$ . No result using FOC

## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

- 1  $f$  is decreasing on  $[a, b]$ 
  - False,  $f'(x) > 0$ .  $f$  is increasing
- 2  $f$  has no local extrema on  $(a, b)$  using  $f'(x) = 0$ 
  - True,  $f'(x) > 0$ . No result using FOC
- 3  $f$  is a constant function on  $(a, b)$

## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

- 1  $f$  is decreasing on  $[a, b]$ 
  - False,  $f'(x) > 0$ .  $f$  is increasing
- 2  $f$  has no local extrema on  $(a, b)$  using  $f'(x) = 0$ 
  - True,  $f'(x) > 0$ . No result using FOC
- 3  $f$  is a constant function on  $(a, b)$ 
  - False,  $f'(x) > 0$ . Constant function has  $f'(x) = 0$

## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

- 1  $f$  is decreasing on  $[a, b]$ 
  - False,  $f'(x) > 0$ .  $f$  is increasing
- 2  $f$  has no local extrema on  $(a, b)$  using  $f'(x) = 0$ 
  - True,  $f'(x) > 0$ . No result using FOC
- 3  $f$  is a constant function on  $(a, b)$ 
  - False,  $f'(x) > 0$ . Constant function has  $f'(x) = 0$
- 4  $f$  is convex on  $(a, b)$



## Question 3, 4, 5

(True/False) If  $g(x) = x^5$ ,  $\lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} = 80$

- Applying L'H rule:  $\Rightarrow \lim_{x \rightarrow 2} g'(x) = 5(2)^4 = 80$  (True)

(True/False) If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$

- Applying chain rule:  $\frac{d}{dx} f(\sqrt{x}) = f'(\sqrt{x}) \frac{d\sqrt{x}}{dx} = \frac{f'(\sqrt{x})}{2\sqrt{x}}$  (False)

Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) > 0$  on  $(a, b)$ . Which is necessarily true?

- 1  $f$  is decreasing on  $[a, b]$ 
  - False,  $f'(x) > 0$ .  $f$  is increasing
- 2  $f$  has no local extrema on  $(a, b)$  using  $f'(x) = 0$ 
  - True,  $f'(x) > 0$ . No result using FOC
- 3  $f$  is a constant function on  $(a, b)$ 
  - False,  $f'(x) > 0$ . Constant function has  $f'(x) = 0$
- 4  $f$  is convex on  $(a, b)$ 
  - False, not necessarily true. Require knowledge on  $f''(x)$

## Section 4

### Question 6 (a)

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x)dx$ ,  $F(x)$  is called Cumulative distribution function

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$ 
    - ① Substitution method:  $u = \exp(-\lambda x) + a$

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$ 
    - 1 Substitution method:  $u = \exp(-\lambda x) + a$
    - 2 Observation:  $\frac{d}{dx} \frac{1}{\exp(-\lambda x) + a} = \frac{\lambda \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$ 
    - 1 Substitution method:  $u = \exp(-\lambda x) + a$
    - 2 Observation:  $\frac{d}{dx} \frac{1}{\exp(-\lambda x) + a} = \frac{\lambda \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$
- Determine  $\lim_{x \rightarrow +\infty} F(x)$



## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$ 
    - 1 Substitution method:  $u = \exp(-\lambda x) + a$
    - 2 Observation:  $\frac{d}{dx} \frac{1}{\exp(-\lambda x) + a} = \frac{\lambda \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$
- Determine  $\lim_{x \rightarrow +\infty} F(x)$ 
  - Note that for all CDF,  $\lim_{x \rightarrow +\infty} F(x) = 1$

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$ 
    - 1 Substitution method:  $u = \exp(-\lambda x) + a$
    - 2 Observation:  $\frac{d}{dx} \frac{1}{\exp(-\lambda x) + a} = \frac{\lambda \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$
- Determine  $\lim_{x \rightarrow +\infty} F(x)$ 
  - Note that for all CDF,  $\lim_{x \rightarrow +\infty} F(x) = 1$
  - Formally:  $\lim_{x \rightarrow +\infty} \frac{a}{\exp(-\lambda x) + a} = \frac{a}{a} = 1 \because \lim_{x \rightarrow +\infty} \exp(-\lambda x) = 0$

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$ 
    - 1 Substitution method:  $u = \exp(-\lambda x) + a$
    - 2 Observation:  $\frac{d}{dx} \frac{1}{\exp(-\lambda x) + a} = \frac{\lambda \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$
- Determine  $\lim_{x \rightarrow +\infty} F(x)$ 
  - Note that for all CDF,  $\lim_{x \rightarrow +\infty} F(x) = 1$
  - Formally:  $\lim_{x \rightarrow +\infty} \frac{a}{\exp(-\lambda x) + a} = \frac{a}{a} = 1 \because \lim_{x \rightarrow +\infty} \exp(-\lambda x) = 0$
- Determine  $\lim_{x \rightarrow -\infty} F(x)$

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$ 
    - 1 Substitution method:  $u = \exp(-\lambda x) + a$
    - 2 Observation:  $\frac{d}{dx} \frac{1}{\exp(-\lambda x) + a} = \frac{\lambda \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$
- Determine  $\lim_{x \rightarrow +\infty} F(x)$ 
  - Note that for all CDF,  $\lim_{x \rightarrow +\infty} F(x) = 1$
  - Formally:  $\lim_{x \rightarrow +\infty} \frac{a}{\exp(-\lambda x) + a} = \frac{a}{a} = 1 \because \lim_{x \rightarrow +\infty} \exp(-\lambda x) = 0$
- Determine  $\lim_{x \rightarrow -\infty} F(x)$ 
  - Note that for all CDF,  $\lim_{x \rightarrow -\infty} F(x) = 0$

## Question 6 (a)

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ , given  $C = 0$ 
  - Note  $F(x) = \int f(x) dx$ ,  $F(x)$  is called Cumulative distribution function
  - Solving  $\int \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} dx$ 
    - 1 Substitution method:  $u = \exp(-\lambda x) + a$
    - 2 Observation:  $\frac{d}{dx} \frac{1}{\exp(-\lambda x) + a} = \frac{\lambda \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$
- Determine  $\lim_{x \rightarrow +\infty} F(x)$ 
  - Note that for all CDF,  $\lim_{x \rightarrow +\infty} F(x) = 1$
  - Formally:  $\lim_{x \rightarrow +\infty} \frac{a}{\exp(-\lambda x) + a} = \frac{a}{a} = 1 \because \lim_{x \rightarrow +\infty} \exp(-\lambda x) = 0$
- Determine  $\lim_{x \rightarrow -\infty} F(x)$ 
  - Note that for all CDF,  $\lim_{x \rightarrow -\infty} F(x) = 0$
  - Formally:  $\lim_{x \rightarrow -\infty} \frac{a}{\exp(-\lambda x) + a} = 0 \because \lim_{x \rightarrow -\infty} \exp(-\lambda x) = \infty$

## Question 6b

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $\int_{-\infty}^x f(t) dt = F(x)$

## Question 6b

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $\int_{-\infty}^x f(t) dt = F(x)$ 
  - Note: integrating pdf = cdf

## Question 6b

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $\int_{-\infty}^x f(t)dt = F(x)$ 
  - Note: integrating pdf = cdf
  - Formally:  $\int_{-\infty}^x f(t)dt = \lim_{s \rightarrow -\infty} F(x) - F(s) = F(x)$



## Question 6b

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $\int_{-\infty}^x f(t) dt = F(x)$ 
  - Note: integrating pdf = cdf
  - Formally:  $\int_{-\infty}^x f(t) dt = \lim_{s \rightarrow -\infty} F(x) - F(s) = F(x)$
- Show that  $F(x)$  is strictly increasing

## Question 6b

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $\int_{-\infty}^x f(t) dt = F(x)$ 
  - Note: integrating pdf = cdf
  - Formally:  $\int_{-\infty}^x f(t) dt = \lim_{s \rightarrow -\infty} F(x) - F(s) = F(x)$
- Show that  $F(x)$  is strictly increasing
  - Note: in general cdf is an increasing function, this question asks a narrower definition of strictly increasing function

## Question 6b

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $\int_{-\infty}^x f(t) dt = F(x)$ 
  - Note: integrating pdf = cdf
  - Formally:  $\int_{-\infty}^x f(t) dt = \lim_{s \rightarrow -\infty} F(x) - F(s) = F(x)$
- Show that  $F(x)$  is strictly increasing
  - Note: in general cdf is an increasing function, this question asks a narrower definition of strictly increasing function
  - Formally:  $f(x) > 0 \because \exp(\cdot) > 0$

## Question 6b

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Show that  $\int_{-\infty}^x f(t) dt = F(x)$ 
  - Note: integrating pdf = cdf
  - Formally:  $\int_{-\infty}^x f(t) dt = \lim_{s \rightarrow -\infty} F(x) - F(s) = F(x)$
- Show that  $F(x)$  is strictly increasing
  - Note: in general cdf is an increasing function, this question asks a narrower definition of strictly increasing function
  - Formally:  $f(x) > 0 \because \exp(\cdot) > 0$
  - Can you think of why this cdf is strictly increasing? Look at the support of the pdf

## Question 6c

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Compute  $F''(x)$

## Question 6c

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Compute  $F''(x)$

- $F''(x) = f'(x) = \frac{d}{dx} \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} =$   
$$\frac{-\lambda^2 a \exp(-\lambda x) (\exp(-\lambda x) + a)^2 - (\lambda a \exp(-\lambda x)) 2(\exp(-\lambda x) + a) \exp(-\lambda x) (-\lambda)}{(\exp(-\lambda x) + a)^4} =$$
  
$$\frac{a \lambda^2 \exp(-\lambda x) (\exp(-\lambda x) - a)}{(\exp(-\lambda x) + a)^3}$$

## Question 6c

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Compute  $F''(x)$

- $$F''(x) = f'(x) = \frac{d}{dx} \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} =$$
$$\frac{-\lambda^2 a \exp(-\lambda x) (\exp(-\lambda x) + a)^2 - (\lambda a \exp(-\lambda x)) 2(\exp(-\lambda x) + a) \exp(-\lambda x) (-\lambda)}{(\exp(-\lambda x) + a)^4} =$$
$$\frac{a \lambda^2 \exp(-\lambda x) (\exp(-\lambda x) - a)}{(\exp(-\lambda x) + a)^3}$$

- Find the inflection point at  $(x_0, F(x_0))$

## Question 6c

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Compute  $F''(x)$

- $$F''(x) = f'(x) = \frac{d}{dx} \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} =$$
$$\frac{-\lambda^2 a \exp(-\lambda x)(\exp(-\lambda x) + a)^2 - (\lambda a \exp(-\lambda x))2(\exp(-\lambda x) + a)\exp(-\lambda x)(-\lambda)}{(\exp(-\lambda x) + a)^4} =$$
$$\frac{a\lambda^2 \exp(-\lambda x)(\exp(-\lambda x) - a)}{(\exp(-\lambda x) + a)^3}$$

- Find the inflection point at  $(x_0, F(x_0))$

- At point of inflection:  $F(x_0) = \frac{1}{2}$ . Data is equal likely to be less than and more than  $x_0$



## Question 6c

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Compute  $F''(x)$

- $$F''(x) = f'(x) = \frac{d}{dx} \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} =$$
$$\frac{-\lambda^2 a \exp(-\lambda x)(\exp(-\lambda x) + a)^2 - (\lambda a \exp(-\lambda x))2(\exp(-\lambda x) + a)\exp(-\lambda x)(-\lambda)}{(\exp(-\lambda x) + a)^4} =$$
$$\frac{a\lambda^2 \exp(-\lambda x)(\exp(-\lambda x) - a)}{(\exp(-\lambda x) + a)^3}$$

- Find the inflection point at  $(x_0, F(x_0))$

- At point of inflection:  $F(x_0) = \frac{1}{2}$ . Data is equal likely to be less than and more than  $x_0$
  - Formally: Inflection at  $F''(x) = 0 \Rightarrow \exp(-\lambda x_0) - a = 0$  (other terms are positive)

## Question 6c

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Compute  $F''(x)$

- $$F''(x) = f'(x) = \frac{d}{dx} \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} =$$
$$\frac{-\lambda^2 a \exp(-\lambda x)(\exp(-\lambda x) + a)^2 - (\lambda a \exp(-\lambda x))2(\exp(-\lambda x) + a)\exp(-\lambda x)(-\lambda)}{(\exp(-\lambda x) + a)^4} =$$
$$\frac{a\lambda^2 \exp(-\lambda x)(\exp(-\lambda x) - a)}{(\exp(-\lambda x) + a)^3}$$

- Find the inflection point at  $(x_0, F(x_0))$

- At point of inflection:  $F(x_0) = \frac{1}{2}$ . Data is equal likely to be less than and more than  $x_0$
  - Formally: Inflection at  $F''(x) = 0 \Rightarrow \exp(-\lambda x_0) - a = 0$  (other terms are positive)
    - $\exp(-\lambda x_0) = a \Rightarrow x_0 = -\frac{\ln(a)}{\lambda}$

## Question 6c

Given pdf:  $f(x) = \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2}$ ,  $F(x) = \frac{a}{\exp(-\lambda x) + a}$ ,  $x \in \mathbf{R}$ ,  $a, \lambda > 0$

- Compute  $F''(x)$

$$\begin{aligned} \bullet \quad F''(x) &= f'(x) = \frac{d}{dx} \frac{\lambda a \exp(-\lambda x)}{(\exp(-\lambda x) + a)^2} = \\ &= \frac{-\lambda^2 a \exp(-\lambda x)(\exp(-\lambda x) + a)^2 - (\lambda a \exp(-\lambda x))2(\exp(-\lambda x) + a)\exp(-\lambda x)(-\lambda)}{(\exp(-\lambda x) + a)^4} = \\ &= \frac{a\lambda^2 \exp(-\lambda x)(\exp(-\lambda x) - a)}{(\exp(-\lambda x) + a)^3} \end{aligned}$$

- Find the inflection point at  $(x_0, F(x_0))$

- At point of inflection:  $F(x_0) = \frac{1}{2}$ . Data is equal likely to be less than and more than  $x_0$
- Formally: Inflection at  $F''(x) = 0 \Rightarrow \exp(-\lambda x_0) - a = 0$  (other terms are positive)
  - $\exp(-\lambda x_0) = a \Rightarrow x_0 = -\frac{\ln(a)}{\lambda}$
- $(x_0, F(x_0)) = \left(-\frac{\ln(a)}{\lambda}, \frac{1}{2}\right)$

## Section 5

### Question 7

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function
  - ① Show by definition

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

- ① Show by definition

- ① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

- ① Show by definition

- ① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

- ②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$



## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

- ① Show by definition

- ① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

- ②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

① Show by definition

- ① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$
- ②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$
- ③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$
- ④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0, L > 0$

- Show  $H$  is homothetic function

① Show by definition

① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

② Using Thm. Rewrite  $H(Q) = h(Q(K, L))$  where  $h(x) = x^2$

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0, L > 0$

- Show  $H$  is homothetic function

- ① Show by definition

- ① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

- ②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

- ② Using Thm. Rewrite  $H(Q) = h(Q(K, L))$  where  $h(x) = x^2$

- ① Show  $Q$  is homogenous:  $Q(tK, tL) = t^{\alpha+\beta} AK^\alpha L^\beta = t^{\alpha+\beta} Q(K, L)$

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0, L > 0$

- Show  $H$  is homothetic function

- ① Show by definition

- ① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

- ②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

- ② Using Thm. Rewrite  $H(Q) = h(Q(K, L))$  where  $h(x) = x^2$

- ① Show  $Q$  is homogenous:  $Q(tK, tL) = t^{\alpha+\beta} AK^\alpha L^\beta = t^{\alpha+\beta} Q(K, L)$

- ② Show  $h$  is increasing function:  $\frac{dh}{dQ} = 2Q > 0 \because K > 0, L > 0 \Rightarrow Q > 0$

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

- ① Show by definition

- ① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

- ②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

- ② Using Thm. Rewrite  $H(Q) = h(Q(K, L))$  where  $h(x) = x^2$

- ① Show  $Q$  is homogenous:  $Q(tK, tL) = t^{\alpha+\beta} AK^\alpha L^\beta = t^{\alpha+\beta} Q(K, L)$

- ② Show  $h$  is increasing function:  $\frac{dh}{dQ} = 2Q > 0 \because K > 0, L > 0 \Rightarrow Q > 0$

- Is  $H$  homogenous function?

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

- ① Show by definition

- ① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

- ②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

- ④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

- ② Using Thm. Rewrite  $H(Q) = h(Q(K, L))$  where  $h(x) = x^2$

- ① Show  $Q$  is homogenous:  $Q(tK, tL) = t^{\alpha+\beta} AK^\alpha L^\beta = t^{\alpha+\beta} Q(K, L)$

- ② Show  $h$  is increasing function:  $\frac{dh}{dQ} = 2Q > 0 \because K > 0, L > 0 \Rightarrow Q > 0$

- Is  $H$  homogenous function?

- Yes, since  $H(tK, tL) = t^{2(\alpha+\beta)} AK^\alpha L^\beta = t^{2(\alpha+\beta)} H(K, L)$

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0, L > 0$

- Show  $H$  is homothetic function

① Show by definition

① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

② Using Thm. Rewrite  $H(Q) = h(Q(K, L))$  where  $h(x) = x^2$

① Show  $Q$  is homogenous:  $Q(tK, tL) = t^{\alpha+\beta} AK^\alpha L^\beta = t^{\alpha+\beta} Q(K, L)$

② Show  $h$  is increasing function:  $\frac{dh}{dQ} = 2Q > 0 \because K > 0, L > 0 \Rightarrow Q > 0$

- Is  $H$  homogenous function?

- Yes, since  $H(tK, tL) = t^{2(\alpha+\beta)} AK^\alpha L^\beta = t^{2(\alpha+\beta)} H(K, L)$

- In general, does homotheticity imply homogeneity?



## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

① Show by definition

① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

② Using Thm. Rewrite  $H(Q) = h(Q(K, L))$  where  $h(x) = x^2$

① Show  $Q$  is homogenous:  $Q(tK, tL) = t^{\alpha+\beta} AK^\alpha L^\beta = t^{\alpha+\beta} Q(K, L)$

② Show  $h$  is increasing function:  $\frac{dh}{dQ} = 2Q > 0 \because K > 0, L > 0 \Rightarrow Q > 0$

- Is  $H$  homogenous function?

- Yes, since  $H(tK, tL) = t^{2(\alpha+\beta)} AK^\alpha L^\beta = t^{2(\alpha+\beta)} H(K, L)$

- In general, does homotheticity imply homogeneity?

- No, there are homothetic function that are not homogenous

## Question 7

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Show  $H$  is homothetic function

① Show by definition

① If  $H(k_1, l_1) = H(k_2, l_2)$ , WTS:  $H(tk_1, tl_1) = H(tk_2, tl_2)$

②  $H(tk_1, tl_1) = t^{2(\alpha+\beta)} H(k_1, l_1) = t^{2(\alpha+\beta)} H(k_2, l_2)$

③  $H(tk_2, tl_2) = t^{2(\alpha+\beta)} H(k_2, l_2)$

④ Since  $H(tk_1, tl_1) = H(tk_2, tl_2) \Rightarrow$  homothetic

② Using Thm. Rewrite  $H(Q) = h(Q(K, L))$  where  $h(x) = x^2$

① Show  $Q$  is homogenous:  $Q(tK, tL) = t^{\alpha+\beta} AK^\alpha L^\beta = t^{\alpha+\beta} Q(K, L)$

② Show  $h$  is increasing function:  $\frac{dh}{dQ} = 2Q > 0 \because K > 0, L > 0 \Rightarrow Q > 0$

- Is  $H$  homogenous function?

- Yes, since  $H(tK, tL) = t^{2(\alpha+\beta)} AK^\alpha L^\beta = t^{2(\alpha+\beta)} H(K, L)$

- In general, does homotheticity imply homogeneity?

- No, there are homothetic function that are not homogenous

- E.g.  $f(x, y) = xy + \exp(xy)$

## Question 7 c

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Find marginal rate of technical substitution for the isoquants of  $H$

## Question 7 c

Given  $H(Q) = Q^2$ ,  $Q(K, L) = AK^\alpha L^\beta$ ,  $K > 0$ ,  $L > 0$

- Find marginal rate of technical substitution for the isoquants of  $H$ 
  - MRTS:

$$\frac{H'_K}{H'_L} = \frac{2(AK^\alpha L^\beta)\alpha K^{\alpha-1}AL^\beta}{2(AK^\alpha L^\beta)\beta L^{\beta-1}AK^\alpha} = \frac{\alpha L}{\beta K}$$