EC2104 Lecture 4 - Integration

ling

Section 1

Integration

Integration

Indefinite integrals

If $F'(x) = f(x)$, then $\int f(x)dx = F(x) + C$. Given
$A = F(0) + C \Rightarrow C = A - F(0)$

Definite integrals

When $F'(x) = f(x)$ for all $x \in (a, b)$, the definite integral of f over $[a, b]$ is
$\int_a^b f(x)dx = F(b) - F(a)$

Integration as Area Under Curve

Let $A(t)$ be a function measures the area under the graph of f over the interval $[a, t] \Rightarrow A(t) = \int_a^b f(t)dt$

Subsection 1

Integration as Area Under Curve

Integration as Area Under Curve

Consider: Find area under the curve for $f(x) = cos(x)$ for $x \in [0, \pi]$

Note:

- $\int cos(x)dx = sin(x) + C$
- When finding area under a curve, one need to know which area are under the $x - axis$
- For area under the $x - axis$, take $-\int f(x)dx$ instead

Wrong method:

$$\int_0^{\pi} cos(x)dx = sin(\pi) - sin(0) = 0$$

Correct method:

$$\int_0^{\pi/2} cos(x)dx - \int_{\pi/2}^{\pi} cos(x)dx = sin(\pi/2) - sin(0) - sin(\pi) + sin(\pi/2) = 2$$

Section 2

Techniques in integration

Techniques in integration

1. Anti-differentiation
$$\int f(x)dx = F(x) + C$$
2. Integration by Substitution
$$\int f(g(x))\frac{dg(x)}{dx}dx = \int f(u)du = F(g(x)) + C$$
   - Differentials: $du = \frac{du}{dx}dx = g'(x)dx$, where $u = g(x)$
3. Integration by Parts
$$\int f(x)\frac{dg(x)}{dx} = f(x)g(x) - \int g(x)\frac{df(x)}{dx}dx$$
4. Leibniz Integral Rule (next slide)

Subsection 1

Leibniz Integral Rule

Leibniz Integral Rule

$$\frac{d}{dt}\int_{a(t)}^{b(t)} f(x,t)dx = \int_{a(t)}^{b(t)} \frac{\partial}{\partial t} f(x,t)dx + f(b(t),t)\frac{db(t)}{dt} - f(a(t),t)\frac{da(t)}{dt}$$

- If $a(t), b(t)$ are constants

$$\frac{d}{dt}\int_{a}^{b} f(x,t)dx = \int_{a}^{b} \frac{\partial}{\partial t} f(x,t)dx$$

- Variable limit form

$$\frac{d}{dt}\int_{a(t)}^{b(t)} f(x)dx = f(b(t))\frac{db(t)}{dt} - f(a(t))\frac{da(t)}{dt}$$

Subsection 2

Integrating infinite bounds

Integrating infinite bounds

To solve

$$\int_0^\infty f(x)dx$$

1. Express ∞ as s
   - i.e. $\int_0^s f(x)dx$
2. Solve the integration in terms of the symbol s
   - i.e. $\int_0^s f(x)dx = F(s) - F(0)$
3. Take limit to infinity
   - i.e. $\lim_{s\to\infty} F(s) - F(0)$

Note:

- $\int_{-\infty}^{\infty} f(x)dx = \int_{-\infty}^{0} f(x)dx + \int_0^\infty f(x)dx$