A Physics-Based Monte Carlo Framework for Space Debris Cascade Risk and Insurance Pricing

Lingxi Lu
Independent Researcher

October 2026 | Version 3.0 | Working Paper

Abstract

The commercial space insurance market, valued at approximately $1.2 billion in 2025, prices satellite risk using historical launch failure rates. No existing actuarial model accounts for cascade risk — the chain-reaction scenario in which one collision generates debris that triggers further collisions, creating correlated losses across an insurance portfolio. I present a coupled physics-actuarial model that simulates debris cascade dynamics in Low Earth Orbit (LEO) using Monte Carlo methods driven by first-principles collision physics, and translates the resulting loss distributions into risk-adjusted insurance premiums via Value at Risk (VaR) and Conditional VaR (CVaR). My baseline simulation at 800 km altitude demonstrates that debris populations grow from 36,500 to ~1.51 million objects over 50 years under business-as-usual conditions, with cascade-adjusted premiums **16.5× higher** than historical-loss pricing ($2,058M vs. $125M per year for a 500-satellite portfolio). The framework is calibrated against three documented collision events (Iridium 33 × Cosmos 2251, Fengyun-1C, Kosmos 1408) and decomposes premiums into expected loss plus risk, capital, and expense loadings following the Solvency II framework.

Keywords: *space debris, Kessler Syndrome, satellite insurance, Monte Carlo simulation, cascade risk, Solvency II, orbital sustainability, space traffic management, insurance pricing*

1. Introduction

1.1 The Unpriced Risk to Orbital Sustainability

The space debris environment in Low Earth Orbit (LEO) has changed. It is no longer a background condition. It is an active operating cost. Tracking networks currently count approximately 36,500 objects larger than 10 cm in Earth orbit. On top of that, there are roughly 1 million objects between 1-10 cm and 130 million smaller fragments, according to the ESA Space Environment Report (2025). The World Economic Forum put a price tag on this: $25.8-$42.3 billion in debris-related costs for LEO assets between 2025 and 2035, assuming nothing changes. That kind of trajectory does not square with the long-term sustainability goals in the UN Space 2030 Agenda or the IADC Space Debris Mitigation Guidelines.

And yet the $1.2 billion space insurance market still prices risk the old way, using historical loss data. Mostly launch and in-orbit failure experience plus single-asset collision probabilities. *Nobody is modeling the chance that one debris event sets off a cascade*, generating correlated claims across a whole

portfolio at once. That gap creates a moral hazard. Operators flying in congested shells pay premiums that have nothing to do with the systemic risk they impose on everyone else in orbit.

1.2 Prior Work and Gap

Freeman et al. (2021) came closest. They built an actuarial modifier for On-Orbit Servicing (OOS) satellite insurance, folding debris mitigation into premiums, but only for direct collision risk. Cascade dynamics, the chain-reaction scenario Kessler (1978) laid out, were not part of their model. Liou and Johnson (2008) ran numerical simulations that showed LEO satellite populations are unstable. They did not take the step of linking those results to insurance pricing. To my knowledge, nobody has coupled first-principles collision physics with actuarial risk metrics to produce regulatory-compliant insurance premiums.

1.3 Contribution

I present the first framework that: (1) couples collision physics with actuarial science so debris growth is driven by actual collision rates, with no separately tuned cascade-rate parameter; (2) implements Poisson-distributed collision events with event-anchored log-normal fragment yields calibrated to observed fragment counts; (3) distinguishes debris–satellite and debris–debris collision channels, with asymmetric yields that prevent unphysical runaway growth from naive N^2 scaling; and (4) calibrates against three documented collision events (Iridium–Cosmos 2009, Fengyun-1C 2007, Kosmos 1408 2021).

2. Methodology

2.1 Physical Layer: Collision-Driven Debris Dynamics

I model a representative LEO shell at altitude h = 800 km with thickness dh = 200 km (700-900 km). The shell volume works out to $V = 1.29 \times 10^{11}$ km^3. I picked this altitude because it has the most defunct satellites and is where most mega-constellations operate. Drag at 800 km is weak enough that debris can stick around for decades, sometimes centuries.

2.1.1 Collision Rate

For collision rates between debris and satellites, I use the Kessler-Flournoy formula:

$$R_{ds} = N_d \cdot N_s \cdot \sigma \cdot v_{rel} \cdot T / V \quad (1)$$

Debris-debris collisions drive the cascade. The rate R_{dd} is proportional to $N_d(N_d-1)$, which means it scales with N_d^2. Double the debris, you quadruple the collision rate. This quadratic feedback is the mathematical root of the cascade. At current population levels (N_d = 36,500, N_s = 8,000), I compute about 7 debris-satellite and 16 debris-debris collisions per year.

2.1.2 Fragment Generation

The *asymmetric fragment yield* between collision types is physically motivated: a fragment striking an intact satellite releases far more energy (and creates more debris) than two small fragments colliding:

Collision Type	Distribution	Median	Physical Rationale
Debris-Satellite (DS)	LogNormal(5.5, 0.8)	~245	Large satellite breakup
Debris-Debris (DD)	LogNormal(1.1, 0.5)	~3	Small fragment impact

Without this distinction, the N^2 scaling of DD rates combined with high yields (~245) would produce unphysical runaway growth. The asymmetry reflects the energy differential: a fragment striking an intact satellite releases orders of magnitude more energy than two small fragments colliding.

2.1.3 Stochastic Time-Stepping

I model collisions as a Poisson process. I broke time into 0.1-year steps, about 36.5 days each. Within each step, I draw DS and DD collision counts, generate fragments from log-normal distributions, apply orbital decay ($\lambda = 0.005$/year with multiplicative Gaussian noise, standard deviation 10%, clipped below at 0.1), and update the population. *There is no separately tuned cascade-rate parameter*; the collision terms are fixed by orbital mechanics. That is not the same as "no free parameters": initial debris count, yield parameters, the 0.5% benchmark, satellite value, and portfolio size are modelling choices, and their influence is quantified in the limitations. Each Monte Carlo path uses a distinct random seed for reproducibility.

2.2 Economic Layer: From Debris to Insurance Premiums

Getting from debris population to financial loss takes four steps. First, I compute per-satellite annual hit probability $p_{hit} = N_d \cdot \sigma \cdot v \cdot T / V$. Second, I calculate per-policy expected loss = $p_{hit} \cdot V_{sat}$ ($50M average). Third, I aggregate the portfolio: expected loss is $p_{hit} \cdot V_{sat} \cdot 500$. An inter-policy loss correlation ($\rho = 0.3$, driven by the shared debris environment) widens the tail but does not change the portfolio mean. Fourth, I pull out VaR and CVaR risk metrics from 1,000 Monte Carlo paths.

2.2.1 Solvency-II-style Premium Decomposition

The total premium is decomposed into four components: P = Expected Loss + Risk Loading + Capital Charge + Expense Loading. The risk loading captures the tail risk beyond the expected loss; the capital charge covers extreme scenarios; and the expense loading accounts for operational costs. The specific multipliers ($\lambda_R = 1.5$, $r = 0.04$, $\lambda_E = 0.15$) follow standard industry practice.

2.3 Consistency Check Against Observed Events

Event	Date	Fragments (>10cm)	Mass
Fengyun-1C (ASAT test)	2007-01-11	~3,500	880 kg
Iridium 33 x Cosmos 2251	2009-02-10	~2,300	1,510 kg
Kosmos 1408 (ASAT test)	2021-11-15	~1,500	2,200 kg

These three events collectively created about 7,300 catalogued fragments. The counts are catalogue totals rather than per-collision yields, so they are used as an order-of-magnitude consistency check, not to fit the operative log-normal parameters. Both ASAT tests (Fengyun-1C, Kosmos 1408) and the accidental collision (Iridium-Cosmos) are represented, ensuring the model captures both intentional and unintentional debris generation mechanisms.

2.4 Validation Against Current Observations

Metric	Model Value	Observed/Market Data	Source
Tracked debris (>10cm), 2025	36,500	36,500	[9]
Global insurance market, 2025	-	$1.2B	[8]
In-orbit failure benchmark	0.5%	Assumption, not observed	Model assumption
Fengyun-1C fragments	~3,500	~3,500	[4]

Iridium-Cosmos fragments	~2,300	~2,300	[4]
Kosmos 1408 fragments	~1,500	~1,500	[4]
Cascade-adjusted premium	$2,058M/year	Not yet priced	Model prediction

The model's initial conditions align with current observational data from the ESA Space Environment Report 2025 [9]. The event counts are order-of-magnitude consistency checks against IADC records and NASA fragmentation data [4], not a formal calibration. The cascade-adjusted premium of $2,058M/year represents what insurance pricing should reflect if cascade risk were properly accounted for, compared with a $125M/year benchmark built from a 0.5% in-orbit failure assumption, not an observed market premium.

3. Results

3.1 Debris Projection

Parameter	Value
Monte Carlo paths	1,000
Time horizon	50 years (2025-2075)
Time step	0.1 year (36.5 days)
Initial debris (>10 cm)	36,500
Active satellites in shell	8,000
Insured portfolio	500 satellites at $50M each
Final debris (mean)	1,509,425
Final debris (median)	1,429,504
Final debris (95th pctile)	2,236,440

Debris grows 41x over 50 years under business-as-usual conditions. The 95th percentile path reaches 2.24 million objects, representing a 61x increase. Growth accelerates over time because debris-debris collisions create new debris that causes further collisions - a positive feedback loop.

3.2 Actuarial Risk Metrics

Metric	Definition	Value
Expected Loss (EL)	Mean worst-year portfolio loss	$921.4M
VaR (95%)	95th pctile worst-year loss	$1,365M
CVaR (95%) / ES	Avg loss beyond VaR(95%)	$1,565M
VaR (99%)	99th pctile worst-year loss	$1,721M
CVaR (99%) / ES	Avg loss beyond VaR(99%)	$1,854M

3.3 Premium Decomposition

Component	Formula	Amount
Expected Loss (EL)	Mean worst-year loss	$921.4M
Risk Loading	1.5 x (CVaR95 - EL)	$966.0M
Capital Charge	4% x (VaR99 - EL)	$32.0M
Expense Loading	15% x EL	$138.2M
TOTAL PREMIUM		$2,058M / year
Per-policy premium	Total / 500 policies	$4,115,098 / year
Historical benchmark	0.5% x $50M x 500	$125.0M / year
CASCADE FACTOR	Cascade / Benchmark	16.5x

Time alignment. *The premium is computed on the worst-year sample, and in 100% of the 1,000 paths the worst year is the final step, so the headline is the fair annual premium at the end of the horizon, not today's. The same model gives expected losses of $22.3M at t = 0 (0.18x the static benchmark) and $111.0M at t = 25 (0.89x). The 16.5x therefore measures how far a static benchmark falls behind a growing cascade by year 50; it is not a claim that current premiums are 16.5x too low today.*

3.4 Key Findings

1. Debris grows 41x over 50 years (36,500 to 1.51M). The growth accelerates over time because debris-debris collisions create new debris that causes further collisions - a positive feedback loop. The qualitative conclusion that LEO is unstable at current population levels is consistent with Liou and Johnson (2008), but no quantitative 41x alignment with their scenario is claimed.

2. The premium gap grows to 16.5x over the horizon. At year 50 the cascade-adjusted premium ($2,058M) exceeds the 0.5% in-orbit failure benchmark ($125M) by $1,933M per year. That gap combines physics with Solvency-II-style loading; on expected loss alone it is 7.4x. The benchmark is all-peril while the model prices debris collisions only, so the ratio is conservative on coverage. The same model gives $22.3M of expected loss at year 0, which is 0.18x the benchmark, so the gap measures how a static premium falls behind a growing cascade rather than how mispriced today's market is. For a 500-satellite portfolio concentrated in one shell, the year-50 gap exceeds the estimated $1.2B global market, which is a solvency signal rather than a market-wide premium forecast. It is a lower bound on the fair premium because the model uses expected claim counts.

3. Tail risk is substantial. CVaR(95%) ($1,565M) exceeds expected loss ($921M) by 70%, indicating that extreme cascade scenarios produce losses well above the mean.

4. Risk loading dominates the premium. The CVaR-based risk loading ($966M) exceeds the expected loss ($921M). This reflects the right-skewed loss distribution and the choice of worst-year loss.

4. Discussion

4.1 Systemic Risk to the Insurance Market

The $1,933M annual gap between the benchmark and cascade-adjusted pricing is a *solvency concern for concentrated books*. When debris density is high, all satellites face elevated risk at the same time. Claims would be correlated across the entire portfolio; correlation widens the tail rather than raising expected

loss. A cascade event at static benchmark levels could exceed insurer reserves on a concentrated book and propagate losses to reinsurers. The historical record gives three reference points: Fengyun-1C in 2007, Iridium-Cosmos in 2009, Kosmos 1408 in 2021 - three events, each creating thousands of trackable fragments, all within a 14-year span.

4.2 Implications for International Space Governance

Right now, the UN Space 2030 Agenda and IADC guidelines set norms for sustainable space operations. But there is no enforcement behind them. This framework proposes an *economic instrument* to go along with regulatory approaches: risk-adjusted insurance pricing that internalizes the externality of debris generation.

Under cascade-aware pricing, operators who place satellites in less congested shells or invest in end-of-life deorbiting would get lower premiums. This gives operators a market-based incentive for sustainability without requiring direct regulation. Operators who contribute to orbital congestion would face premiums that reflect the systemic risk they impose on all space users.

4.3 Policy Recommendations

1. Insurance regulators (e.g., EIOPA, NAIC) should require space insurers to disclose whether cascade risk is incorporated into pricing models.

2. The IADC and UNCOPUOS should commission independent studies on the macroeconomic implications of underpriced cascade risk.

3. Space agencies should explore premium differentiation as a condition for launch licensing -- satellites in congested shells pay higher insurance, which in funds debris mitigation.

4. Reinsurance markets should develop cascade-risk-specific instruments (e.g., catastrophe bonds tied to debris density thresholds) to transfer systemic risk off insurer balance sheets.

4.4 Limitations

Several limitations warrant discussion. First, the single-shell approximation (800 km only) does not capture the full 3D orbital environment, and the framework covers only >10 cm catalogue objects; the roughly one million 1-10 cm objects, which are the actual largest contributors to current satellite loss risk, are not modeled. Second, the debris-debris collision rate is an upper bound: both channels share a 10 m2 satellite cross-section, whereas the true DD cross-section for two >10 cm objects is ~0.01 m2, 2-3 orders of magnitude smaller. The 16.5x cascade factor is materially dependent on the DD term. Third, the orbital decay parameter is an effective removal rate lumping atmospheric drag, collisional grinding, and catalogue dropout, not pure drag lifetime. Fourth, the 0.5% benchmark is an in-orbit failure assumption, not an observed market premium: 0.3% gives 27.5x and 1% gives 8.2x. Fifth, the fragment yield is an event-anchored log-normal prior calibrated to observed fragment counts from three historical events, not a physics-based calibration using collision mass and energy.

4.5 Future Work

Future research directions include: (1) multi-shell model with cross-altitude migration; (2) NORAD TLE data integration for empirical debris tracking; (3) dynamic constellation deployment modeling; (4) ADR scenario analysis and premium reduction quantification; (5) reinsurance layer pricing (excess-of-loss, stop-loss, catastrophe bonds).

5. Conclusion

This framework shows that incorporating physics-based cascade dynamics into insurance pricing produces a *16.5x gap* relative to a 0.5% in-orbit failure benchmark (7.4x on expected loss alone). The main innovation is replacing ad-hoc cascade parameters with first-principles collision physics. Debris growth is driven by actual collision rates, with event-anchored fragment yields and asymmetric treatment of debris-satellite versus debris-debris collisions.

The $1,933M annual pricing gap is a systemic risk to the global space insurance market, but it is also an opportunity. Risk-adjusted pricing can work as an economic instrument for orbital sustainability, going along with the regulatory framework of the UN Space 2030 Agenda. By aligning private insurance costs with systemic debris risk, this framework gives us a market-based pathway toward long-term sustainability of outer space activities. The magnitude is conditional on the single-shell approximation, the DD cross-section upper bound, and the choice of benchmark; these caveats are discussed in the limitations.

References

[1] Kessler, D.J. and Cour-Palais, B.G. (1978). "Collision Frequency of Artificial Satellites: The Creation of a Debris Belt." *J. Geophys. Res.*, 83(A6), 2637-2646.

[2] Kessler, D.J. (1981). "Derivation of the collision probability between orbiting objects." *Icarus*, 48(1), 39-48.

[3] Liou, J.-C. and Johnson, N.L. (2008). "Instability of the present LEO satellite populations." *Advances in Space Research*, 41(7), 1046-1053.

[4] Johnson, N.L. et al. (2008). "History of On-Orbit Satellite Fragmentations, 14th Edition." NASA/TM-2008-214779, NASA Orbital Debris Program Office.

[5] Freeman, R. et al. (2021). "An actuarial modifier for underwriting OOS satellite insurance." *J. Space Operations*.

[6] European Commission (2009). "Solvency II Directive." 2009/138/EC.

[7] IADC (2007). "Space Debris Mitigation Guidelines." Inter-Agency Space Debris Coordination Committee.

[8] UNCOPUOS (2019). "Space 2030 Agenda." United Nations Committee on the Peaceful Uses of Outer Space.

[9] World Economic Forum / Centre for Space Futures (2026). "Clear Orbit, Secure Future: A Call to Action on Space Debris."

[10] ESA Space Environment Report (2025). European Space Agency.

Data and Code Availability: The simulation code is released under the Apache License 2.0 (see LICENSE); the manuscript text is distributed under CC BY 4.0. Source code (Python package *cascade_risk/*), configuration files, and simulation output data are available from the author upon request.

Correspondence: ling.xi.lu@gmail.com