Conditional Probability: Intermediate: Takeaways



by Dataquest Labs, Inc. - All rights reserved © 2020

Concepts

- Given events A and B:
 - P(A) means finding the probability of A
 - P(A|B) means finding the conditional probability of A (given that B occurs)
 - $P(A \cap B)$ means finding the probability that both A and B occur
 - $P(A \cup B)$ means finding the probability that A occurs or B occurs (this doesn't exclude the situation where both A and B occur)
- For any events A and B, it's true that:

- The order of conditioning is important, so P(A|B) is different P(B|A).
- If event A occurs and the probability of B remains unchanged (and vice versa), then events A and B are said to be **statistically independent** (although the term "independent" is used more often). Mathematically, statistical independence between A and B implies that:

• If events events A and B are **statistically dependent**, it means that the occurrence of event A changes the probability of event B and vice versa. In mathematical terms, this means that:

• If three events A, B, C are **mutually independent**, then two conditions must hold: they should be pairwise independent, but also independent together. If any of these two conditions doesn't hold, then the events are not mutually independent.

•										
	Tho	mult	inlica	tion	rulo	for	donon	dont	events:	
	THE	mull	IDIICo	ILIUII	Tuic	TOT	ucbell	uciii	CVCIILS.	

• The mutliplication rule for independent events:

Resources

- An intuitive approach to understanding independent events
- An easy intro to some basic conditional probability concepts
- A brief reminder on set complements



Takeaways by Dataquest Labs, Inc. - All rights reserved © 2020