RESEARCH STATEMENT

Jianfeng Lin

My research lies in gauge theory, Floer homotopy theory, and low-dimensional topology, with an emphasis on the smooth topology of four-manifolds. Dimension four exhibits distinctive phenomena, including infinitely many smooth structures on a single closed manifold and infinitely generated mapping class groups of simply connected closed manifolds. My work reveals further manifestations of this exceptional topology through two themes: developing Floer and Seiberg–Witten invariants and their homotopy refinements, and using configuration spaces and geometric techniques to study diffeomorphisms and embedded submanifolds.

Floer homotopy theory and gauge-theoretic applications. Floer homology and its homotopy refinements connect three-dimensional geometry with four-dimensional topology. With Khandhawit and Sasahira, we constructed the unfolded Seiberg–Witten–Floer spectra, relative Bauer–Furuta invariants for four-manifolds with boundary, and a gluing theorem [1, 2]. These constructions retain equivariant information lost upon passage to homology, enabling applications of equivariant stable homotopy theory to four-manifold topology.

A major application of Floer homotopy theory concerns the 11/8-conjecture: every closed smooth spin four-manifold X should satisfy $b_2(X) \geq \frac{11}{8}|\sigma(X)|$. Furuta proved $b_2(X) \geq \frac{10}{8}|\sigma(X)| + 2$ for nonzero signature by studying $\mathrm{Pin}(2)$-equivariant maps between representation spheres. With Hopkins, Shi, and Xu, we completely resolved the resulting Furuta–Jones problem using equivariant Mahowald invariants [6]. For $|\sigma(X)| \geq 32$, our theorem gives $b_2(X) \geq \frac{10}{8}|\sigma(X)| + c(|\sigma(X)|)$, where the explicit correction term is 4, 6, or 8. This determines the optimal restrictions on intersection forms obtainable through this equivariant sphere-map approach.

Another application concerns stabilization. Wall's work and subsequent developments show that many exotic phenomena on simply connected four-manifolds disappear after sufficiently many stabilizations (i.e., connected sums with $S^2 \times S^2$). Whether one stabilization is enough has been an open question for a long time. I proved that the exotic Dehn twist on $K3\#K3$, studied by Kronheimer and Mrowka, survives one stabilization: it remains topologically isotopic to the identity but not smoothly so [7]. This was the first such example on a simply connected four-manifold and its shows that one stabilization is not always enough. The key is the $\mathrm{Pin}(2)$-equivariant families Bauer–Furuta invariant, whose additional equivariant information detects nontriviality even when the corresponding S^1-equivariant invariant vanishes.

I also developed families Seiberg–Witten theory to study diffeomorphism groups and moduli spaces. My gluing formula proves McDuff's conjecture that $\pi_1(\mathrm{Symp}(X)) \to \pi_1(\mathrm{Diff}(X))$ is not surjective for any closed symplectic four-manifold containing a smoothly embedded sphere of self-intersection -1 or -2 [8]. With Konno, we constructed gauge-theoretic characteristic classes for four-manifold bundles and proved that homological stability fails for the moduli spaces of every simply connected closed smooth four-manifold [9], contrasting with the high-dimensional theorem of Galatius and Randal-Williams.

With Konno, Mukherjee, and Muñoz-Echániz, we connected families gauge theory with singularity theory by proving infinite-order results for boundary Dehn twists on symplectic fillings of links of minimally elliptic singularities [10]. Applications include Milnor monodromy and compactly supported exotic diffeomorphisms of exotic $\mathbb{R}^4$'s. More recently, families Bauer–Furuta invariants yielded constraints on Lefschetz fibrations with four-dimensional fibers. We constructed closed simply connected symplectic four-manifolds with smoothly nontrivial Torelli symplectomorphisms, showing that their symplectic Torelli mapping class groups are not generated by squared Dehn–Seidel twists [11]. Du and Li independently obtained counterexamples to Donaldson's generation question using a different construction [12].

In three papers with Ruberman and Saveliev, we studied the Seiberg–Witten invariant λ_{SW} of homology $S^1 \times S^3$'s. We proved a splitting formula relating it to a Lefschetz number on reduced monopole Floer homology and the Frøyshov invariant [3]. We then computed monopole Lefschetz numbers for branched-covering involutions and finite-order covering transformations [4, 5]. These formulas make λ_{SW} computable for classes of mapping tori, relate it to the Furuta–Ohta invariant, and yield applications to

positive scalar curvature, homology cobordism, and knot concordance.

With Majewski and Rzemieniecki, we applied families gauge theory and parametrized equivariant homotopy theory to exceptional holonomy. We proved that every Cayley fibration of a compact torsion-free Spin(7)-manifold with full holonomy must have singular fibers [13], establishing a new application of four-dimensional gauge theory to higher-dimensional geometry.

Configuration spaces, diffeomorphism groups, and submanifolds. My second direction uses configuration-space invariants to study four-dimensional isotopy and smooth structures. Watanabe disproved the four-dimensional Smale conjecture using Kontsevich's configuration-space integrals. Answering a question of Freedman, Xie and I proved that these integrals depend only on a formal smooth structure: a vector bundle structure on the topological tangent microbundle [14]. The central construction is a Fulton–MacPherson compactification that requires no genuine smooth structure. Consequently, $\mathrm{Top}(4)$, the homeomorphism group of $\mathbb{R}^4$, has nonzero rational homotopy groups in infinitely many degrees, and $\mathrm{Diff}(X) \to \mathrm{Homeo}(X)$ is not rationally 2-connected for any compact oriented smooth four-manifold X. This contrasts with the homotopy equivalence of these groups in dimensions at most three.

With Wu, Xie, and Zhang, we extended the Dax invariant from disks to closed surfaces and obtained many applications. For every closed oriented surface Σ of positive genus, we classified embeddings $\Sigma \hookrightarrow \Sigma \times S^2$ geometrically dual to $\{\mathrm{pt}\} \times S^2$, answering Gabai's question, and proved that the mapping class group of $\Sigma \times S^2$ is infinitely generated [15]. Combining this construction with J-holomorphic curve techniques, we answered questions of McDuff–Salamon and Cieliebak–Eliashberg–Mishachev on symplectic 4-manifolds. On every irrational ruled surface, we found infinitely many connected components of symplectic forms in each fixed symplectic cohomology class. We also constructed infinitely many cohomologous, formally homotopic symplectic forms that are pairwise nonhomotopic through symplectic forms. These give the first such failures of connectedness and of the corresponding h-principle on closed four-manifolds.

In subsequent work with Xie and Zhang, we developed methods based on Budney–Gabai's configuration-space invariant w_3. For every compact connected oriented three-manifold Y with infinite fundamental group, we proved that the mapping class group of $Y \times [0,1]$ relative to the boundary is infinitely generated [16]. We also showed that every smooth two-component split sphere link in S^4 admits infinitely many smooth splitting three-spheres that are pairwise topologically non-isotopic [17], extending Tatsuoka's theorem from the unlink to arbitrarily knotted components. These results expose failures of isotopy uniqueness through configuration-space methods complementary to gauge theory.

References

[1] T. Khandhawit, J. Lin, and H. Sasahira. The unfolded Seiberg–Witten–Floer spectra I: Definition, invariance and examples. *Geometry & Topology* **22**(4) (2018), 2027–2114.

[2] T. Khandhawit, J. Lin, and H. Sasahira. The unfolded Seiberg–Witten–Floer spectra II: Relative invariant and the gluing theorem. *Journal of Differential Geometry* **124**(2) (2023), 231–316.

[3] J. Lin, D. Ruberman, and N. Saveliev. A splitting theorem for the Seiberg–Witten invariant of a homology $S^1 \times S^3$. *Geometry & Topology* **22**(5) (2018), 2865–2942.

[4] J. Lin, D. Ruberman, and N. Saveliev. On the Frøyshov invariant and monopole Lefschetz number. *Journal of Differential Geometry* **123**(3) (2023), 523–593.

[5] J. Lin, D. Ruberman, and N. Saveliev. On the monopole Lefschetz number of finite order diffeomorphisms. *Geometry & Topology* **25** (2021), 3591–3628.

[6] M. J. Hopkins, J. Lin, X. D. Shi, and Z. Xu. Intersection forms of spin 4-manifolds and the Pin(2)-equivariant Mahowald invariant. *Communications of the American Mathematical Society* **2** (2022), 22–132.

[7] J. Lin. Isotopy of the Dehn twist on $K3\#K3$ after a single stabilization. *Geometry & Topology* **27**(5) (2023), 1987–2012.

[8] J. Lin. The family Seiberg–Witten invariant and nonsymplectic loops of diffeomorphisms. Accepted in *Geometry & Topology*. arXiv:2208.12082.

[9] H. Konno and J. Lin. Homological instability for moduli spaces of smooth 4-manifolds. *Geometry & Topology* **30** (2026), 2993–3068.

[10] H. Konno, J. Lin, A. Mukherjee, and J. Muñoz-Echániz. On four-dimensional Dehn twists and Milnor fibrations. Accepted in *Duke Mathematical Journal*. arXiv:2409.11961.

[11] H. Konno, J. Lin, A. Mukherjee, and J. Muñoz-Echániz. Constraints on Lefschetz fibrations with four-dimensional fibers from Seiberg–Witten theory. Preprint (2025), arXiv:2511.00601.

[12] Y. Du and T. Li. Elliptic twists and Dehn twists on the one point Kähler blowup of the K3 surface. In preparation (2025).

[13] J. Lin, V. F. Majewski, and J. Rzemieniecki. Non-existence of smooth full-holonomy Cayley fibrations. Preprint (2026), arXiv:2603.26920.

[14] J. Lin and Y. Xie. Configuration space integrals and formal smooth structures. Preprint (2023), arXiv:2310.14156.

[15] J. Lin, W. Wu, Y. Xie, and B. Zhang. Dax invariants, light bulbs, and isotopies of symplectic structures. Preprint (2025), arXiv:2501.16083.

[16] J. Lin, Y. Xie, and B. Zhang. Pseudo-isotopies of 3-manifolds with infinite fundamental groups. Preprint (2026), arXiv:2602.09454.

[17] J. Lin, Y. Xie, and B. Zhang. Splitting spheres for S^2-links in S^4. Preprint (2026), arXiv:2608.02785.