

# Mapping Hydrodynamics for the Facilitated Exclusion and Zero-range Processes

**Linjie Zhao**

joint work with **Clément Erignoux** and **Marielle Simon**

Inria Lille-Nord Europe, France

March 2022

## Notation and Results

## Formal Proof

## Mapping

# Introduction

- ▶ Hydrodynamic limit: understanding macroscopic behaviors, usually described by PDEs via a long-time and large-space scaling limit, of microscopic systems.
- ▶ Symmetric exclusion: heat equation. Asymmetric exclusion: conservation law.
- ▶ Goal: deriving Stefan problems from interacting particle systems.

# Introduction


- ▶ Hydrodynamic limit: understanding macroscopic behaviors, usually described by PDEs via a long-time and large-space scaling limit, of microscopic systems.
- ▶ Symmetric exclusion: heat equation. Asymmetric exclusion: conservation law.
- ▶ Goal: deriving Stefan problems from interacting particle systems.

# Introduction

- ▶ Hydrodynamic limit: understanding macroscopic behaviors, usually described by PDEs via a long-time and large-space scaling limit, of microscopic systems.
- ▶ Symmetric exclusion: heat equation. Asymmetric exclusion: conservation law.
- ▶ Goal: deriving Stefan problems from interacting particle systems.

# Facilitated Exclusion Process


The **state space** is  $\Sigma_N := \{0, 1\}^{\mathbb{L}_N}$ , where  $\mathbb{L}_N = \mathbb{Z}$  or  $\mathbb{T}_N$ . For a **configuration**  $\eta = (\eta_x)_{x \in \mathbb{L}_N} \in \Sigma_N$ ,  $\eta_x = 1$  if and only if  $x$  is occupied by a particle.



- ▶ **Exclusion** rule: there is at most one particle at each site.
- ▶ **Facilitated** rule: a particle has to be pushed forward in order to jump.
- ▶ **Symmetric**:  $p = p'$ ,  $\mathbb{L}_N = \mathbb{T}_N$ . **Asymmetric**:  $p > p'$ ,  $\mathbb{L}_N = \mathbb{Z}$ .

# Facilitated Exclusion Process


The **state space** is  $\Sigma_N := \{0, 1\}^{\mathbb{L}_N}$ , where  $\mathbb{L}_N = \mathbb{Z}$  or  $\mathbb{T}_N$ . For a **configuration**  $\eta = (\eta_x)_{x \in \mathbb{L}_N} \in \Sigma_N$ ,  $\eta_x = 1$  if and only if  $x$  is occupied by a particle.



- **Exclusion** rule: there is at most one particle at each site.
- **Facilitated** rule: a particle has to be pushed forward in order to jump.
- **Symmetric**:  $p = p'$ ,  $\mathbb{L}_N = \mathbb{T}_N$ . **Asymmetric**:  $p > p'$ ,  $\mathbb{L}_N = \mathbb{Z}$ .

# Facilitated Exclusion Process

The **state space** is  $\Sigma_N := \{0, 1\}^{\mathbb{L}_N}$ , where  $\mathbb{L}_N = \mathbb{Z}$  or  $\mathbb{T}_N$ . For a **configuration**  $\eta = (\eta_x)_{x \in \mathbb{L}_N} \in \Sigma_N$ ,  $\eta_x = 1$  if and only if  $x$  is occupied by a particle.




- ▶ **Exclusion** rule: there is at most one particle at each site.
- ▶ **Facilitated** rule: a particle has to be pushed forward in order to jump.
- ▶ **Symmetric**:  $p = p'$ ,  $\mathbb{L}_N = \mathbb{T}_N$ . **Asymmetric**:  $p > p'$ ,  $\mathbb{L}_N = \mathbb{Z}$ .



# Facilitated Exclusion Process

The **state space** is  $\Sigma_N := \{0, 1\}^{\mathbb{L}_N}$ , where  $\mathbb{L}_N = \mathbb{Z}$  or  $\mathbb{T}_N$ . For a **configuration**  $\eta = (\eta_x)_{x \in \mathbb{L}_N} \in \Sigma_N$ ,  $\eta_x = 1$  if and only if  $x$  is occupied by a particle.



- ▶ **Exclusion** rule: there is at most one particle at each site.
- ▶ **Facilitated** rule: a particle has to be pushed forward in order to jump.
- ▶ **Symmetric**:  $p = p'$ ,  $\mathbb{L}_N = \mathbb{T}_N$ . **Asymmetric**:  $p > p'$ ,  $\mathbb{L}_N = \mathbb{Z}$ .

# Facilitated Exclusion Process

- ▶ For  $x, y \in \mathbb{L}_N$ ,  $\eta^{x,y}$  denotes the configuration obtained from  $\eta$  by swapping the values at sites  $x$  and  $y$ .
- ▶ The **jump rate** is given by

$$c_{x,x+1}(\eta) = p\eta_{x-1}\eta_x(1 - \eta_{x+1}) + p'(1 - \eta_x)\eta_{x+1}\eta_{x+2}.$$




- ▶ For any local function  $f: \Sigma_N \rightarrow \mathbb{R}$ , the **infinitesimal generator** is

$$\mathcal{L}_N^{\text{FEX}} f(\eta) := \sum_{x \in \mathbb{L}_N} c_{x,x+1}(\eta) (f(\eta^{x,x+1}) - f(\eta)).$$

# Facilitated Exclusion Process

- ▶ For  $x, y \in \mathbb{L}_N$ ,  $\eta^{x,y}$  denotes the configuration obtained from  $\eta$  by swapping the values at sites  $x$  and  $y$ .
- ▶ The **jump rate** is given by

$$c_{x,x+1}(\eta) = p\eta_{x-1}\eta_x(1 - \eta_{x+1}) + p'(1 - \eta_x)\eta_{x+1}\eta_{x+2}.$$




- ▶ For any local function  $f: \Sigma_N \rightarrow \mathbb{R}$ , the **infinitesimal generator** is

$$\mathcal{L}_N^{\text{FEX}} f(\eta) := \sum_{x \in \mathbb{L}_N} c_{x,x+1}(\eta) (f(\eta^{x,x+1}) - f(\eta)).$$

# Facilitated Exclusion Process

- ▶ For  $x, y \in \mathbb{L}_N$ ,  $\eta^{x,y}$  denotes the configuration obtained from  $\eta$  by swapping the values at sites  $x$  and  $y$ .
- ▶ The **jump rate** is given by

$$c_{x,x+1}(\eta) = p\eta_{x-1}\eta_x(1 - \eta_{x+1}) + p'(1 - \eta_x)\eta_{x+1}\eta_{x+2}.$$




- ▶ For any local function  $f: \Sigma_N \rightarrow \mathbb{R}$ , the **infinitesimal generator** is

$$\mathcal{L}_N^{\text{FEX}} f(\eta) := \sum_{x \in \mathbb{L}_N} c_{x,x+1}(\eta) (f(\eta^{x,x+1}) - f(\eta)).$$

# Phase Transitions

The process displays a **phase transition** at the critical particle density  $\rho_c = 1/2$ .

- ▶ If initially  $\rho > 1/2$ , then the system evolves until there are no longer two neighboring empty sites.




- ▶ If initially  $\rho < 1/2$ , then the system evolves until all particles are isolated, and thus can no longer move.



# Phase Transitions

The process displays a **phase transition** at the critical particle density  $\rho_c = 1/2$ .

- ▶ If initially  $\rho > 1/2$ , then the system evolves until there are no longer two neighboring empty sites.




- ▶ If initially  $\rho < 1/2$ , then the system evolves until all particles are isolated, and thus can no longer move.



# Phase Transitions

The process displays a **phase transition** at the critical particle density  $\rho_c = 1/2$ .

- ▶ If initially  $\rho > 1/2$ , then the system evolves until there are no longer two neighboring empty sites.



- ▶ If initially  $\rho < 1/2$ , then the system evolves until all particles are isolated, and thus can no longer move.



# Invariant Measures $\pi_\rho$ on $\{0, 1\}^{\mathbb{Z}}$

- Fix  $\rho > 1/2$ . For some  $\sigma = (\sigma_x, 1 \leq x \leq \ell)$  where  $\sigma_x \in \{0, 1\}$  and  $(\sigma_x, \sigma_{x+1}) \neq (0, 0)$ , let  $k = \sum_{x=1}^{\ell} \sigma_x$ .

$$\begin{aligned}\pi_\rho(\eta : \eta_x = \sigma_x, \forall 1 \leq x \leq \ell) \\ = (1 - \rho) \left( \frac{1 - \rho}{\rho} \right)^{\ell-1-k} \left( \frac{2\rho - 1}{\rho} \right)^{2k - \ell + 1 - \sigma(1) - \sigma(\ell)}\end{aligned}$$

- Put a hole in some position with probability  $1 - \rho$ , then put a random **geometric number** of parameter  $\frac{1-\rho}{\rho}$  particles to its right, then a hole, starts again and so on.

$$\begin{aligned}\pi_\rho(11) &= \pi_\rho(1) - \pi_\rho(01) = \rho - (1 - \rho) \times 1 = 2\rho - 1. \\ \pi_\rho(111) &= \pi_\rho(11) - \pi_\rho(011) \\ &= (2\rho - 1) - (1 - \rho) \times 1 \times \frac{2\rho - 1}{\rho} = \frac{(2\rho - 1)^2}{\rho}.\end{aligned}$$



# Invariant Measures $\pi_\rho$ on $\{0, 1\}^{\mathbb{Z}}$

- Fix  $\rho > 1/2$ . For some  $\sigma = (\sigma_x, 1 \leq x \leq \ell)$  where  $\sigma_x \in \{0, 1\}$  and  $(\sigma_x, \sigma_{x+1}) \neq (0, 0)$ , let  $k = \sum_{x=1}^{\ell} \sigma_x$ .

$$\begin{aligned}\pi_\rho(\eta : \eta_x = \sigma_x, \forall 1 \leq x \leq \ell) \\ = (1 - \rho) \left( \frac{1 - \rho}{\rho} \right)^{\ell-1-k} \left( \frac{2\rho - 1}{\rho} \right)^{2k - \ell + 1 - \sigma(1) - \sigma(\ell)}\end{aligned}$$

- Put a hole in some position with probability  $1 - \rho$ , then put a random **geometric number** of parameter  $\frac{1-\rho}{\rho}$  particles to its right, then a hole, starts again and so on.

$$\begin{aligned}\pi_\rho(11) &= \pi_\rho(1) - \pi_\rho(01) = \rho - (1 - \rho) \times 1 = 2\rho - 1. \\ \pi_\rho(111) &= \pi_\rho(11) - \pi_\rho(011) \\ &= (2\rho - 1) - (1 - \rho) \times 1 \times \frac{2\rho - 1}{\rho} = \frac{(2\rho - 1)^2}{\rho}.\end{aligned}$$

# Invariant Measures $\pi_\rho$ on $\{0, 1\}^{\mathbb{Z}}$

- Fix  $\rho > 1/2$ . For some  $\sigma = (\sigma_x, 1 \leq x \leq \ell)$  where  $\sigma_x \in \{0, 1\}$  and  $(\sigma_x, \sigma_{x+1}) \neq (0, 0)$ , let  $k = \sum_{x=1}^{\ell} \sigma_x$ .

$$\begin{aligned}\pi_\rho(\eta : \eta_x = \sigma_x, \forall 1 \leq x \leq \ell) \\ = (1 - \rho) \left( \frac{1 - \rho}{\rho} \right)^{\ell-1-k} \left( \frac{2\rho - 1}{\rho} \right)^{2k - \ell + 1 - \sigma(1) - \sigma(\ell)}\end{aligned}$$

- Put a hole in some position with probability  $1 - \rho$ , then put a random **geometric number** of parameter  $\frac{1-\rho}{\rho}$  particles to its right, then a hole, starts again and so on.

$$\begin{aligned}\pi_\rho(11) &= \pi_\rho(1) - \pi_\rho(01) = \rho - (1 - \rho) \times 1 = 2\rho - 1. \\ \pi_\rho(111) &= \pi_\rho(11) - \pi_\rho(011) \\ &= (2\rho - 1) - (1 - \rho) \times 1 \times \frac{2\rho - 1}{\rho} = \frac{(2\rho - 1)^2}{\rho}.\end{aligned}$$

# Invariant Measures $\pi_\rho$ on $\mathbb{Z}$

- ▶ For  $\rho \leq 1/2$ ,  $\pi_\rho$  is concentrated on frozen configurations.

- ▶  $\rho = 1/2$ ,

...0101010101...

- ▶  $\rho = 1/3$ ,

...001001001...

# Invariant Measures $\pi_\rho$ on $\mathbb{Z}$

- ▶ For  $\rho \leq 1/2$ ,  $\pi_\rho$  is concentrated on frozen configurations.
- ▶  $\rho = 1/2$ ,

...0101010101...

- ▶  $\rho = 1/3$ ,

...001001001...

# Invariant Measures $\pi_\rho$ on $\mathbb{Z}$

► For  $\rho \leq 1/2$ ,  $\pi_\rho$  is concentrated on frozen configurations.

►  $\rho = 1/2$ ,

...0101010101...

►  $\rho = 1/3$ ,

...001001001...

# Hydrodynamic Limit

- If at the initial time, there exists  $\rho^{\text{ini}} : \mathbb{T}$  (or  $\mathbb{R}$ )  $\rightarrow \mathbb{R}$

$$\eta_x(0) \approx \rho^{\text{ini}}\left(\frac{x}{N}\right),$$

then, at any time  $t > 0$ ,

$$\eta_x(t\theta(N)) \approx \rho\left(t, \frac{x}{N}\right)$$

for some  $\rho(t, u)$  with initial condition  $\rho^{\text{ini}}$ .



$$\theta(N) = \begin{cases} N^2 \text{ (diffusive scaling),} & \text{in the symmetric case;} \\ N \text{ (hyperbolic scaling),} & \text{in the asymmetric case.} \end{cases}$$

# Hydrodynamic Limit

- If at the initial time, there exists  $\rho^{\text{ini}} : \mathbb{T}$  (or  $\mathbb{R}$ )  $\rightarrow \mathbb{R}$

$$\eta_x(0) \approx \rho^{\text{ini}}\left(\frac{x}{N}\right),$$

then, at any time  $t > 0$ ,

$$\eta_x(t\theta(N)) \approx \rho\left(t, \frac{x}{N}\right)$$

for some  $\rho(t, u)$  with initial condition  $\rho^{\text{ini}}$ .



$$\theta(N) = \begin{cases} N^2 \text{ (diffusive scaling),} & \text{in the symmetric case;} \\ N \text{ (hyperbolic scaling),} & \text{in the asymmetric case.} \end{cases}$$

$$p = p' = 1$$

- Space/Time scaling:

	Macroscopic	Microscopic
Space	$u \approx \frac{x}{N}$	$x$
Time	$t$	$tN^2$

- Free boundary problem:

$$\begin{cases} \partial_t \rho = \partial_u^2 \left( \frac{2\rho-1}{\rho} \mathbf{1}_{\{\rho > 1/2\}} \right) & \text{in } \{t > 0\} \times \mathbb{T}, \\ \rho(0, \cdot) = \rho^{\text{ini}} & \text{in } \mathbb{T}. \end{cases}$$

- The weak solution is given in terms of integration by parts formula.



$$p = p' = 1$$

- Space/Time scaling:

	Macroscopic	Microscopic
Space	$u \approx \frac{x}{N}$	$x$
Time	$t$	$tN^2$

- Free boundary problem:

$$\begin{cases} \partial_t \rho = \partial_u^2 \left( \frac{2\rho-1}{\rho} \mathbf{1}_{\{\rho > 1/2\}} \right) & \text{in } \{t > 0\} \times \mathbb{T}, \\ \rho(0, \cdot) = \rho^{\text{ini}} & \text{in } \mathbb{T}. \end{cases}$$

- The weak solution is given in terms of integration by parts formula.

$$p = p' = 1$$

- Space/Time scaling:

	Macroscopic	Microscopic
Space	$u \approx \frac{x}{N}$	$x$
Time	$t$	$tN^2$

- Free boundary problem:

$$\begin{cases} \partial_t \rho = \partial_u^2 \left( \frac{2\rho-1}{\rho} \mathbf{1}_{\{\rho > 1/2\}} \right) & \text{in } \{t > 0\} \times \mathbb{T}, \\ \rho(0, \cdot) = \rho^{\text{ini}} & \text{in } \mathbb{T}. \end{cases}$$

- The weak solution is given in terms of integration by parts formula.

$$p = p' = 1$$

- Define the **empirical measure**  $\pi_t^N(du)$ , which is a random measure on  $\mathbb{T}$ , as

$$\pi_t^N(du) = \frac{1}{N} \sum_{x \in \mathbb{T}_N} \eta_x(tN^2) \delta_{x/N}(du).$$

- For a sequence of measures  $\{\mu_N\}_{N \geq 1}$ ,  $\mu$  on  $\mathbb{T}$  with total mass bounded by one,

$$\lim_{N \rightarrow \infty} \mu_N = \mu$$

if for any continuous function  $f: \mathbb{T} \rightarrow \mathbb{R}$ ,

$$\lim_{N \rightarrow \infty} \int f d\mu_N = \int f d\mu.$$

$$p = p' = 1$$

- Define the **empirical measure**  $\pi_t^N(du)$ , which is a random measure on  $\mathbb{T}$ , as

$$\pi_t^N(du) = \frac{1}{N} \sum_{x \in \mathbb{T}_N} \eta_x(tN^2) \delta_{x/N}(du).$$

- For a sequence of measures  $\{\mu_N\}_{N \geq 1}$ ,  $\mu$  on  $\mathbb{T}$  with total mass bounded by one,

$$\lim_{N \rightarrow \infty} \mu_N = \mu$$

if for any continuous function  $f: \mathbb{T} \rightarrow \mathbb{R}$ ,

$$\lim_{N \rightarrow \infty} \int f d\mu_N = \int f d\mu.$$

$$p = p' = 1$$

Theorem [Blondel *et al.* '20, '21, Erignoux, Simon and Z. '22]

Under mild conditions on the initial density profile  $\rho^{\text{ini}}$ , if

$$\lim_{N \rightarrow \infty} \pi_0^N(du) = \rho^{\text{ini}}(u)du \quad \text{in probability,}$$

then, for any time  $t > 0$ ,

$$\lim_{N \rightarrow \infty} \pi_t^N(du) = \rho(t, u)du \quad \text{in probability,}$$

where  $\rho(t, u)$  is the unique weak solution to

$$\partial_t \rho = \partial_u^2 \left( \frac{2\rho - 1}{\rho} \mathbf{1}_{\{\rho > 1/2\}} \right)$$

with initial condition  $\rho^{\text{ini}}$ .

$$p = 1 - p' \in (1/2, 1]$$

- Space/Time scaling:

	Macroscopic	Microscopic
Space	$u \approx \frac{x}{N}$	$x$
Time	$t$	$tN$

- Hyperbolic free boundary problem:

$$\begin{cases} \partial_t \rho + (2\rho - 1) \partial_u \left( \frac{(1-\rho)(2\rho-1)}{\rho} \mathbf{1}_{\{\rho > 1/2\}} \right) = 0 & \text{in } \{t > 0\} \times \mathbb{R}, \\ \rho(0, \cdot) = \rho^{\text{ini}} & \text{in } \mathbb{R}. \end{cases}$$

- The weak solutions are **not** unique! Physical relevant solution is the **entropy solution**.

$$p = 1 - p' \in (1/2, 1]$$

- Space/Time scaling:

	Macroscopic	Microscopic
Space	$u \approx \frac{x}{N}$	$x$
Time	$t$	$tN$

- Hyperbolic free boundary problem:

$$\begin{cases} \partial_t \rho + (2\rho - 1) \partial_u \left( \frac{(1-\rho)(2\rho-1)}{\rho} \mathbf{1}_{\{\rho > 1/2\}} \right) = 0 & \text{in } \{t > 0\} \times \mathbb{R}, \\ \rho(0, \cdot) = \rho^{\text{ini}} & \text{in } \mathbb{R}. \end{cases}$$

- The weak solutions are **not** unique! Physical relevant solution is the **entropy solution**.

$$p = 1 - p' \in (1/2, 1]$$

- Space/Time scaling:

	Macroscopic	Microscopic
Space	$u \approx \frac{x}{N}$	$x$
Time	$t$	$tN$

- Hyperbolic free boundary problem:

$$\begin{cases} \partial_t \rho + (2\rho - 1) \partial_u \left( \frac{(1-\rho)(2\rho-1)}{\rho} \mathbf{1}_{\{\rho > 1/2\}} \right) = 0 & \text{in } \{t > 0\} \times \mathbb{R}, \\ \rho(0, \cdot) = \rho^{\text{ini}} & \text{in } \mathbb{R}. \end{cases}$$

- The weak solutions are **not** unique! Physical relevant solution is the **entropy solution**.



# Entropy solution

Let  $\mathfrak{H}(\rho) = \frac{(1-\rho)(2\rho-1)}{\rho} \mathbf{1}_{\{\rho > 1/2\}}$ . An entropy solution satisfies

- (*entropy inequality*) for any **non-negative** test function  $\varphi \in C^{1,1}(\mathbb{R}_+ \times \mathbb{R})$  with compact support in  $(0, \infty) \times \mathbb{R}$ , for any  $0 \leq c \leq 1$ ,

$$\int_0^\infty \langle |\rho_t - c|, \partial_t \varphi_t \rangle + (2\rho - 1) \langle \mathfrak{q}(\rho_t; c, \mathfrak{H}), \partial_u \varphi_t \rangle dt \geq 0,$$

where  $\mathfrak{q}(\rho; c, \mathfrak{H}) = \text{sign}(\rho - c)(\mathfrak{H}(\rho) - \mathfrak{H}(c))$  ;

- (*initial condition*) for any  $A > 0$ ,

$$\lim_{t \rightarrow 0} \int_{-A}^A |\rho_t(u) - \rho^{\text{ini}}(u)| du = 0.$$

# Entropy solution

Let  $\mathfrak{H}(\rho) = \frac{(1-\rho)(2\rho-1)}{\rho} \mathbf{1}_{\{\rho > 1/2\}}$ . An entropy solution satisfies

- (*entropy inequality*) for any *non-negative* test function  $\varphi \in C^{1,1}(\mathbb{R}_+ \times \mathbb{R})$  with compact support in  $(0, \infty) \times \mathbb{R}$ , for any  $0 \leq c \leq 1$ ,

$$\int_0^\infty \langle |\rho_t - c|, \partial_t \varphi_t \rangle + (2\rho - 1) \langle \mathfrak{q}(\rho_t; c, \mathfrak{H}), \partial_u \varphi_t \rangle dt \geq 0,$$

where  $\mathfrak{q}(\rho; c, \mathfrak{H}) = \text{sign}(\rho - c)(\mathfrak{H}(\rho) - \mathfrak{H}(c))$  ;

- (*initial condition*) for any  $A > 0$ ,

$$\lim_{t \rightarrow 0} \int_{-A}^A |\rho_t(u) - \rho^{\text{ini}}(u)| du = 0.$$

$$p = 1 - p' \in (1/2, 1)$$

Theorem [Erignoux, Simon and Z. '22]

Under mild conditions on the initial density profile  $\rho^{\text{ini}}$ , if

$$\lim_{N \rightarrow \infty} \pi_0^N(du) = \rho^{\text{ini}}(u)du \quad \text{in probability,}$$

then, for any time  $t > 0$ ,

$$\lim_{N \rightarrow \infty} \pi_t^N(du) = \rho(t, u)du \quad \text{in probability,}$$

where  $\rho(t, u)$  is the unique entropy solution to

$$\partial_t \rho + (2p - 1) \partial_u \left( \frac{(1 - \rho)(2\rho - 1)}{\rho} \mathbf{1}_{\{\rho > 1/2\}} \right) = 0$$

with initial condition  $\rho^{\text{ini}}$ .

Notation and Results

Formal Proof

Mapping

$$p = p' = 1$$

► First recall

$$\rho(t, \frac{x}{N}) \approx \mathbb{E}[\eta_x(tN^2)], \quad x \in \mathbb{T}_N,$$

or equivalently,

$$\rho(t, u) \approx \mathbb{E}[\eta_{uN}(tN^2)], \quad u \in \mathbb{T}.$$

► Conservation law

$$\frac{d}{dt} \mathbb{E}[\eta_{uN}(tN^2)] = N^2 \left( \mathbb{E}[j_{uN-1, uN}(tN^2)] - \mathbb{E}[j_{uN, uN+1}(tN^2)] \right),$$

where  $j_{x, x+1}(t)$  is the instantaneous current from  $x$  to  $x+1$ ,

$$j_{x, x+1}(t) = \eta_{x-1}(t)\eta_x(t)(1 - \eta_{x+1}(t)) - \eta_{x+2}(t)\eta_{x+1}(t)(1 - \eta_x(t)).$$

$$p = p' = 1$$

► First recall

$$\rho(t, \frac{x}{N}) \approx \mathbb{E}[\eta_x(tN^2)], \quad x \in \mathbb{T}_N,$$

or equivalently,

$$\rho(t, u) \approx \mathbb{E}[\eta_{uN}(tN^2)], \quad u \in \mathbb{T}.$$

► Conservation law

$$\frac{d}{dt} \mathbb{E}[\eta_{uN}(tN^2)] = N^2 \left( \mathbb{E}[j_{uN-1, uN}(tN^2)] - \mathbb{E}[j_{uN, uN+1}(tN^2)] \right),$$

where  $j_{x, x+1}(t)$  is the instantaneous current from  $x$  to  $x+1$ ,

$$j_{x, x+1}(t) = \eta_{x-1}(t)\eta_x(t)(1 - \eta_{x+1}(t)) - \eta_{x+2}(t)\eta_{x+1}(t)(1 - \eta_x(t)).$$

$$p = p' = 1$$

► Gradient condition

$$\begin{aligned} j_{x,x+1}(t) &= \eta_{x-1}(t)\eta_x(t)(1 - \eta_{x+1}(t)) - \eta_{x+2}(t)\eta_{x+1}(t)(1 - \eta_x(t)) \\ &= h_x(t) - h_{x+1}(t), \end{aligned}$$

where

$$h_x(t) = \eta_{x-1}(t)\eta_x(t) + \eta_x(t)\eta_{x+1}(t) - \eta_{x-1}(t)\eta_x(t)\eta_{x+1}(t).$$

► Therefore,

$$\partial_t \rho(t, u) \approx N^2 \left( \mathbb{E}[h_{uN-1}(tN^2)] + \mathbb{E}[h_{uN+1}(tN^2)] - 2\mathbb{E}[h_{uN}(tN^2)] \right).$$

$$p = p' = 1$$

► Gradient condition

$$\begin{aligned} j_{x,x+1}(t) &= \eta_{x-1}(t)\eta_x(t)(1 - \eta_{x+1}(t)) - \eta_{x+2}(t)\eta_{x+1}(t)(1 - \eta_x(t)) \\ &= h_x(t) - h_{x+1}(t), \end{aligned}$$

where

$$h_x(t) = \eta_{x-1}(t)\eta_x(t) + \eta_x(t)\eta_{x+1}(t) - \eta_{x-1}(t)\eta_x(t)\eta_{x+1}(t).$$

► Therefore,

$$\partial_t \rho(t, u) \approx N^2 \left( \mathbb{E}[h_{uN-1}(tN^2)] + \mathbb{E}[h_{uN+1}(tN^2)] - 2\mathbb{E}[h_{uN}(tN^2)] \right).$$



$$p = p' = 1$$

- **Local ergodicity:** the distribution of the process at time  $tN^2$  around site  $uN$  is approximately  $\pi_{\rho(t,u)}$ .
- For  $\rho(t, u) > 1/2$ ,

$$\begin{aligned}\mathbb{E}[h_{uN}(tN^2)] &\approx \pi_{\rho(t,u)}(h) = 2\pi_{\rho(t,u)}(11) - \pi_{\rho(t,u)}(111) \\ &= \frac{2\rho(t, u) - 1}{\rho(t, u)}.\end{aligned}$$

The above term is zero if  $\rho(t, u) \leq 1/2$ .

- Let  $\mathcal{H}(\rho) = \frac{(2\rho-1)}{\rho} \mathbf{1}_{\rho>1/2}$ . Then,

$$\begin{aligned}\partial_t \rho(t, u) &\approx N^2 \left( \mathcal{H}(\rho(t, u - \tfrac{1}{N})) + \mathcal{H}(\rho(t, u + \tfrac{1}{N})) - 2\mathcal{H}(\rho(t, u)) \right) \\ &\approx \partial_u^2 \mathcal{H}(\rho(t, u)).\end{aligned}$$

$$p = p' = 1$$

- ▶ **Local ergodicity:** the distribution of the process at time  $tN^2$  around site  $uN$  is approximately  $\pi_{\rho(t,u)}$ .
- ▶ For  $\rho(t, u) > 1/2$ ,

$$\begin{aligned}\mathbb{E}[h_{uN}(tN^2)] &\approx \pi_{\rho(t,u)}(h) = 2\pi_{\rho(t,u)}(11) - \pi_{\rho(t,u)}(111) \\ &= \frac{2\rho(t, u) - 1}{\rho(t, u)}.\end{aligned}$$

The above term is zero if  $\rho(t, u) \leq 1/2$ .

- ▶ Let  $\mathcal{H}(\rho) = \frac{(2\rho-1)}{\rho} \mathbf{1}_{\rho > 1/2}$ . Then,

$$\begin{aligned}\partial_t \rho(t, u) &\approx N^2 \left( \mathcal{H}(\rho(t, u - \tfrac{1}{N})) + \mathcal{H}(\rho(t, u + \tfrac{1}{N})) - 2\mathcal{H}(\rho(t, u)) \right) \\ &\approx \partial_u^2 \mathcal{H}(\rho(t, u)).\end{aligned}$$

$$p = p' = 1$$

- ▶ **Local ergodicity:** the distribution of the process at time  $tN^2$  around site  $uN$  is approximately  $\pi_{\rho(t,u)}$ .
- ▶ For  $\rho(t, u) > 1/2$ ,

$$\begin{aligned}\mathbb{E}[h_{uN}(tN^2)] &\approx \pi_{\rho(t,u)}(h) = 2\pi_{\rho(t,u)}(11) - \pi_{\rho(t,u)}(111) \\ &= \frac{2\rho(t, u) - 1}{\rho(t, u)}.\end{aligned}$$

The above term is zero if  $\rho(t, u) \leq 1/2$ .

- ▶ Let  $\mathcal{H}(\rho) = \frac{(2\rho-1)}{\rho} \mathbf{1}_{\rho>1/2}$ . Then,

$$\begin{aligned}\partial_t \rho(t, u) &\approx N^2 \left( \mathcal{H}(\rho(t, u - \tfrac{1}{N})) + \mathcal{H}(\rho(t, u + \tfrac{1}{N})) - 2\mathcal{H}(\rho(t, u)) \right) \\ &\approx \partial_u^2 \mathcal{H}(\rho(t, u)).\end{aligned}$$

$$p = 1 - p' \in (1/2, 1]$$

- ▶ The above arguments only show that  $\rho(t, u)$  is a weak solution.
- ▶ We need to prove a microscopic version of the entropy inequality.
- ▶ For attractive systems, *cf.* [Rezakhanlou'91].

$$p = 1 - p' \in (1/2, 1]$$

- ▶ The above arguments only show that  $\rho(t, u)$  is a weak solution.
- ▶ We need to prove a microscopic version of the entropy inequality.
- ▶ For attractive systems, *cf.* [Rezakhanlou'91].

$$p = 1 - p' \in (1/2, 1]$$

- ▶ The above arguments only show that  $\rho(t, u)$  is a weak solution.
- ▶ We need to prove a microscopic version of the entropy inequality.
- ▶ For attractive systems, *cf.* [Rezakhanlou'91].

Notation and Results


Formal Proof

Mapping

# Facilitated Zero Range Process


- In the exclusion process, label the empty sites from the left to the right in an increasing order. Let  $X_y(t)$  be the position of the  $y$ -th empty site at time  $t$ .

EP:



- Let  $\omega_y(t) \in \{0, 1, 2, \dots\}$  be the number of particles between the  $y$ -th and  $(y+1)$ -th empty sites at time  $t$ .


ZRP:






# Facilitated Zero Range Process

- In the exclusion process, label the empty sites from the left to the right in an increasing order. Let  $X_y(t)$  be the position of the  $y$ -th empty site at time  $t$ .




- Let  $\omega_y(t) \in \{0, 1, 2, \dots\}$  be the number of particles between the  $y$ -th and  $(y+1)$ -th empty sites at time  $t$ .



# Facilitated Zero Range Process


- ▶ If there are at least two particles at a site, then one of them jumps to the right (resp. to the left) at rate  $p$  (resp.  $p'$ ).



- ▶ The critical particle density is  $\alpha_c = 1$ .
- ▶ The process has a family of **product** invariant measures when the particle density  $\alpha > 1$ .
- ▶ The process is **attractive**.

# Facilitated Zero Range Process


- ▶ If there are at least two particles at a site, then one of them jumps to the right (resp. to the left) at rate  $p$  (resp.  $p'$ ).



- ▶ The critical particle density is  $\alpha_c = 1$ .
- ▶ The process has a family of **product** invariant measures when the particle density  $\alpha > 1$ .
- ▶ The process is **attractive**.

# Facilitated Zero Range Process


- ▶ If there are at least two particles at a site, then one of them jumps to the right (resp. to the left) at rate  $p$  (resp.  $p'$ ).



- ▶ The critical particle density is  $\alpha_c = 1$ .
- ▶ The process has a family of **product** invariant measures when the particle density  $\alpha > 1$ .
- ▶ The process is **attractive**.

# Facilitated Zero Range Process


- ▶ If there are at least two particles at a site, then one of them jumps to the right (resp. to the left) at rate  $p$  (resp.  $p'$ ).



- ▶ The critical particle density is  $\alpha_c = 1$ .
- ▶ The process has a family of **product** invariant measures when the particle density  $\alpha > 1$ .
- ▶ The process is **attractive**.


# Attractiveness

Attractiveness: if  $\omega_y(0) \leq \tilde{\omega}_y(0)$  for all  $y$ , then there exists a coupling such that  $\omega_y(t) \leq \tilde{\omega}_y(t)$  for all  $y$  and  $t > 0$ .




# Attractiveness

Attractiveness: if  $\omega_y(0) \leq \tilde{\omega}_y(0)$  for all  $y$ , then there exists a coupling such that  $\omega_y(t) \leq \tilde{\omega}_y(t)$  for all  $y$  and  $t > 0$ .




# Attractiveness

Attractiveness: if  $\omega_y(0) \leq \tilde{\omega}_y(0)$  for all  $y$ , then there exists a coupling such that  $\omega_y(t) \leq \tilde{\omega}_y(t)$  for all  $y$  and  $t > 0$ .





# Proof Outline



# Macroscopic Mapping

- ▶ Exclusion  $\eta_x$ ,  $\rho(t, u)$ ; Zero range  $\omega_y$ ,  $\alpha(t, v)$ .
- ▶ Consider the process on  $\mathbb{Z}$ . For any  $\varphi \in C_c(\mathbb{R})$ ,

$$\frac{1}{N} \sum_{x \in \mathbb{Z}} [1 - \eta_x(tN)] \varphi(x/N) = \frac{1}{N} \sum_{y \in \mathbb{Z}} \varphi(X_y(tN)/N).$$

- ▶ Note that

$$X_y = \sum_{y'=1}^y [X_{y'} - X_{y'-1}] + X_0 = \sum_{y'=1}^y [1 + \omega_{y'-1}] + X_0.$$

- ▶ Let  $\sigma_t = \lim_{N \rightarrow \infty} X_0(tN)/N$ ,

$$\int (1 - \rho(t, u)) \varphi(u) du = \int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv$$

# Macroscopic Mapping

- ▶ Exclusion  $\eta_x$ ,  $\rho(t, u)$ ; Zero range  $\omega_y$ ,  $\alpha(t, v)$ .
- ▶ Consider the process on  $\mathbb{Z}$ . For any  $\varphi \in C_c(\mathbb{R})$ ,

$$\frac{1}{N} \sum_{x \in \mathbb{Z}} [1 - \eta_x(tN)] \varphi(x/N) = \frac{1}{N} \sum_{y \in \mathbb{Z}} \varphi(X_y(tN)/N).$$

- ▶ Note that

$$X_y = \sum_{y'=1}^y [X_{y'} - X_{y'-1}] + X_0 = \sum_{y'=1}^y [1 + \omega_{y'-1}] + X_0.$$

- ▶ Let  $\sigma_t = \lim_{N \rightarrow \infty} X_0(tN)/N$ ,

$$\int (1 - \rho(t, u)) \varphi(u) du = \int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv$$

# Macroscopic Mapping

- ▶ Exclusion  $\eta_x$ ,  $\rho(t, u)$ ; Zero range  $\omega_y$ ,  $\alpha(t, v)$ .
- ▶ Consider the process on  $\mathbb{Z}$ . For any  $\varphi \in C_c(\mathbb{R})$ ,

$$\frac{1}{N} \sum_{x \in \mathbb{Z}} [1 - \eta_x(tN)] \varphi(x/N) = \frac{1}{N} \sum_{y \in \mathbb{Z}} \varphi(X_y(tN)/N).$$

- ▶ Note that

$$X_y = \sum_{y'=1}^y [X_{y'} - X_{y'-1}] + X_0 = \sum_{y'=1}^y [1 + \omega_{y'-1}] + X_0.$$

- ▶ Let  $\sigma_t = \lim_{N \rightarrow \infty} X_0(tN)/N$ ,

$$\int (1 - \rho(t, u)) \varphi(u) du = \int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv$$

# Macroscopic Mapping

- ▶ Exclusion  $\eta_x, \rho(t, u)$ ; Zero range  $\omega_y, \alpha(t, v)$ .
- ▶ Consider the process on  $\mathbb{Z}$ . For any  $\varphi \in C_c(\mathbb{R})$ ,

$$\frac{1}{N} \sum_{x \in \mathbb{Z}} [1 - \eta_x(tN)] \varphi(x/N) = \frac{1}{N} \sum_{y \in \mathbb{Z}} \varphi(X_y(tN)/N).$$

- ▶ Note that

$$X_y = \sum_{y'=1}^y [X_{y'} - X_{y'-1}] + X_0 = \sum_{y'=1}^y [1 + \omega_{y'-1}] + X_0.$$

- ▶ Let  $\sigma_t = \lim_{N \rightarrow \infty} X_0(tN)/N$ ,

$$\int (1 - \rho(t, u)) \varphi(u) du = \int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv$$

# Macroscopic Mapping

- ▶ We already have

$$\int (1 - \rho(t, u)) \varphi(u) du = \int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv$$

- ▶ Let

$$u(t, v) = \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t$$

It has an inverse  $v(t, u)$ .

- ▶ Then,

$$\int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv = \int \varphi(u) [1 + \alpha(t, v(t, u))]^{-1} du.$$

- ▶ We finally have

$$\rho(t, u) = \frac{\alpha(t, v(t, u))}{1 + \alpha(t, v(t, u))}.$$

# Macroscopic Mapping

- ▶ We already have

$$\int (1 - \rho(t, u)) \varphi(u) du = \int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv$$

- ▶ Let

$$u(t, v) = \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t$$

It has an inverse  $v(t, u)$ .

- ▶ Then,

$$\int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv = \int \varphi(u) [1 + \alpha(t, v(t, u))]^{-1} du.$$

- ▶ We finally have

$$\rho(t, u) = \frac{\alpha(t, v(t, u))}{1 + \alpha(t, v(t, u))}.$$

# Macroscopic Mapping

- ▶ We already have

$$\int (1 - \rho(t, u)) \varphi(u) du = \int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv$$

- ▶ Let

$$u(t, v) = \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t$$

It has an inverse  $v(t, u)$ .

- ▶ Then,

$$\int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv = \int \varphi(u) [1 + \alpha(t, v(t, u))]^{-1} du.$$

- ▶ We finally have

$$\rho(t, u) = \frac{\alpha(t, v(t, u))}{1 + \alpha(t, v(t, u))}.$$



# Macroscopic Mapping

- ▶ We already have

$$\int (1 - \rho(t, u)) \varphi(u) du = \int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv$$

- ▶ Let

$$u(t, v) = \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t$$

It has an inverse  $v(t, u)$ .

- ▶ Then,

$$\int \varphi \left( \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t \right) dv = \int \varphi(u) [1 + \alpha(t, v(t, u))]^{-1} du.$$

- ▶ We finally have

$$\rho(t, u) = \frac{\alpha(t, v(t, u))}{1 + \alpha(t, v(t, u))}.$$

# Macroscopic Mapping

- Relation between  $\alpha(t, v)$  and  $\rho(t, u)$ :

$$\rho(t, u) = \frac{\alpha(t, v(t, u))}{1 + \alpha(t, v(t, u))},$$

$$u(t, v) = \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t$$

.

- Prove

$\rho(t, u)$  is solution of FEP

$\Leftrightarrow \alpha(t, v)$  is solution of FZRP.

# Macroscopic Mapping

- Relation between  $\alpha(t, v)$  and  $\rho(t, u)$ :

$$\rho(t, u) = \frac{\alpha(t, v(t, u))}{1 + \alpha(t, v(t, u))},$$

$$u(t, v) = \int_0^v (1 + \alpha(t, v')) dv' + \sigma_t$$

.

- Prove

$\rho(t, u)$  is solution of FEP

$\Leftrightarrow \alpha(t, v)$  is solution of FZRP.

# Macroscopic Mapping

- ▶ Problem: the solutions to the Stefan problem and hyperbolic equation are not smooth.
- ▶ For the Stefan problem

$$\partial_t \rho = \partial_u^2 \mathcal{H}(\rho),$$

we approximate the function  $\mathcal{H}$  by some good  $\mathcal{H}^\varepsilon$ ,

$$\partial_t \rho^\varepsilon = \partial_u^2 \mathcal{H}^\varepsilon(\rho^\varepsilon).$$

- ▶ For the hyperbolic equation

$$\partial_t \rho + (2\rho - 1) \partial_u \mathfrak{H}(\rho) = 0,$$

we use viscous approximation method,

$$\partial_t \rho^\varepsilon + (2\rho - 1) \partial_u \mathfrak{H}(\rho^\varepsilon) = \varepsilon \partial_u^2 \rho^\varepsilon.$$

# Macroscopic Mapping

- ▶ Problem: the solutions to the Stefan problem and hyperbolic equation are not smooth.
- ▶ For the Stefan problem

$$\partial_t \rho = \partial_u^2 \mathcal{H}(\rho),$$

we approximate the function  $\mathcal{H}$  by some good  $\mathcal{H}^\varepsilon$ ,

$$\partial_t \rho^\varepsilon = \partial_u^2 \mathcal{H}^\varepsilon(\rho^\varepsilon).$$

- ▶ For the hyperbolic equation

$$\partial_t \rho + (2\rho - 1)\partial_u \mathfrak{H}(\rho) = 0,$$

we use viscous approximation method,

$$\partial_t \rho^\varepsilon + (2\rho - 1)\partial_u \mathfrak{H}(\rho^\varepsilon) = \varepsilon \partial_u^2 \rho^\varepsilon.$$

# Macroscopic Mapping

- ▶ Problem: the solutions to the Stefan problem and hyperbolic equation are not smooth.
- ▶ For the Stefan problem

$$\partial_t \rho = \partial_u^2 \mathcal{H}(\rho),$$

we approximate the function  $\mathcal{H}$  by some good  $\mathcal{H}^\varepsilon$ ,

$$\partial_t \rho^\varepsilon = \partial_u^2 \mathcal{H}^\varepsilon(\rho^\varepsilon).$$





- ▶ For the hyperbolic equation

$$\partial_t \rho + (2\rho - 1)\partial_u \mathfrak{H}(\rho) = 0,$$

we use viscous approximation method,

$$\partial_t \rho^\varepsilon + (2\rho - 1)\partial_u \mathfrak{H}(\rho^\varepsilon) = \varepsilon \partial_u^2 \rho^\varepsilon.$$

# References

-  Kipnis, C., & Landim, C. (1998). *Scaling limits of interacting particle systems* (Vol. 320). Springer Science & Business Media.
-  Blondel, O., Erignoux, C., Sasada, M., & Simon, M. (2020, February). Hydrodynamic limit for a facilitated exclusion process. *In Annales de l'Institut Henri Poincaré, Probabilités et Statistiques* (Vol. 56, No. 1, pp. 667-714).
-  Blondel, O., Erignoux, C., & Simon, M. (2021). Stefan problem for a nonergodic facilitated exclusion process. *Probability and Mathematical Physics*, 2(1), 127-178.
-  Erignoux, C., Simon, M., & Zhao, L. (2022). Mapping hydrodynamics for the facilitated exclusion and zero-range processes. *arXiv preprint arXiv:2202.04469*.

# Thanks!

*Email:* linjie.zhao@inria.fr