

Scaling Limits of the SSEP with a Slow Site

T. Franco¹, P. Gonçalves², R. Marinho² and L. Zhao³

1 Background

One of the biggest problems in *interacting particle systems* is the derivation of macroscopic equations as large-scale limits from these models. For example, the Navier-Stokes equation is recovered from the evolution of the momentum in the incompressible lattices gas model described in [2]. The attainment of these partial differential equations from particle systems is known as *hydrodynamic limit*. The analysis of the hydrodynamic behaviour of interacting particle systems gives validity to many partial differential equations in the literature [18].

Obtaining hydrodynamic equations becomes even more difficult when the desired ones have boundary conditions. It is well known that the hydrodynamics of symmetric simple exclusion processes (SSEP) is driven by the heat equation [18]. In order to obtain the heat equation with boundary conditions, the SSEP in different settings was considered. We mention here the SSEP with slow bonds [6, 8–10], the SSEP with slow boundary [1, 7, 13] and the SSEP with a slow site [11].

For the SSEP with a slow site, the asymmetry in the jump rates that a particle enters and leaves the origin makes the usual arguments much more difficult. In [11], they only proved the supercritical case. We solved the remaining cases recently.

2 Model and Main Results

The model is defined on the one dimensional discrete torus \mathbb{T}_n with n sites or on the integer lattice \mathbb{Z} . There is at most one particle per site. A particle at site x jumps to one of its neighbors $y \sim x$ at rate ξ_x^n provided that the target site is empty, where

$$\xi_x^n = \begin{cases} \alpha n^{-\beta}, & \text{if } x = 0, \\ 1, & \text{if } x \neq 0. \end{cases}$$

Here $\alpha > 0$ and $\beta \geq 0$.




Figure 1: At most one particle per site. A particle at the origin jumps at a lower rate $\alpha n^{-\beta}$.

In order to see the process on its diffusive time scaling, we will consider the process $\{\eta_t : t \geq 0\}$ with *time speeded up by n^2* . We recall from [11, Proposition 2.1] that for any $p \in [0, 1]$, the Bernoulli product measure ν_p defined by

$$\nu_p(\eta; \eta(x) = 1) =: m_p(x) = \begin{cases} \frac{p/(\alpha n^{-\beta})}{(1-p)+p/(\alpha n^{-\beta})}, & \text{if } x = 0, \\ p, & \text{if } x \neq 0, \end{cases} \quad (1)$$

is *reversible* for the process $\{\eta_t : t \geq 0\}$.

2.1 Hydrodynamics

The process is defined on \mathbb{T}_n . Fix a profile $\rho_0 : \mathbb{T} \rightarrow [0, 1]$, representing the initial density of particles. To avoid uninteresting technical complications, we assume that ρ_0 is continuous at all $x \in \mathbb{T} \setminus \{0\}$ and bounded from below by a positive constant,

$$\zeta := \inf_{x \in \mathbb{T}} \rho_0(x) > 0. \quad (2)$$

The *empirical measure* is defined as

$$\pi_t^n(du) = \frac{1}{n} \sum_x \eta_t(x) \delta_{x/n}(du). \quad (3)$$

Theorem 1 (Law of Large Numbers for the Empirical Measure)

Let the initial measure $\{\mu_n; n \in \mathbb{N}\}$ be a sequence of product measures on $\{0, 1\}^{\mathbb{T}_n}$ with marginals given by

$$\mu_n\{\eta : \eta(x) = 1\} = \rho_0(x/n). \quad (4)$$

Then, for every $t > 0$, $\pi_t^n(du)$ converges in the weak topology to $\rho(t, u)du$ in probability as $n \rightarrow \infty$, where $\rho(t, u)$ stands for the unique weak solution of

- the heat equation with *periodic boundary conditions* if $\beta \in [0, 1)$;
- the heat equation with *Robin boundary conditions* if $\beta = 1$;
- and the heat equation with *Neumann boundary conditions* if $\beta > 1$.

2.2 Equilibrium Fluctuations

The process is defined on \mathbb{Z} . The *fluctuation field*, which is the linear functional acting on test functions H , is defined as

$$\mathcal{Y}_t^n(H) = \frac{1}{\sqrt{n}} \sum_{x \in \mathbb{Z}} H\left(\frac{x}{n}\right) \bar{\eta}_t(x),$$

where $\bar{\eta}_t(x) = \eta_t(x) - m_p(x)$.

Domain of the Test Functions in Different Regimes

We will set \mathcal{S}_β

- for $\beta \in [0, 1)$, as the usual Schwartz space;
- for $\beta = 1$, as the space of functions $H : \mathbb{R} \rightarrow \mathbb{R}$ such that
 - (1) H is continuous and infinitely differentiable, except possibly at $x = 0$.
 - (2) $H(0) = \frac{1}{2}[H(0^+) + H(0^-)]$.
 - (3) For all integers $k, \ell \geq 0$,

$$\|H\|_{k,\ell} := \sup_{x \neq 0} \left| (1 + |x|^\ell) \frac{d^k H}{dx^k}(x) \right| < \infty.$$

- (4) For any integer $k \geq 0$,

$$\frac{d^{2k+1} H}{dx^{2k+1}}(0^+) = \frac{d^{2k+1} H}{dx^{2k+1}}(0^-) = \alpha \left(\frac{d^{2k} H}{dx^{2k}}(0^+) - \frac{d^{2k} H}{dx^{2k}}(0^-) \right).$$

- for $\beta > 1$, as the space of functions $H : \mathbb{R} \rightarrow \mathbb{R}$ that satisfy (1), (2') below, (3), (4) with $\alpha = 0$, (2') H is continuous from the right at zero.

Denote by $\mathcal{S}'_\beta(\mathbb{R})$ the dual space of $\mathcal{S}_\beta(\mathbb{R})$. We will define ∇_β and Δ_β

- for $\beta \in [0, 1)$, by the usual first and second space derivatives;
- for $\beta \geq 1$, by

$$\nabla_\beta H(u) = \begin{cases} \frac{dH}{du}(u), & \text{if } u \neq 0, \\ \lim_{u \rightarrow 0^+} \frac{dH}{du}(u), & \text{if } u = 0, \end{cases} \quad \Delta_\beta H(u) = \begin{cases} \frac{d^2 H}{du^2}(u), & \text{if } u \neq 0, \\ \lim_{u \rightarrow 0^+} \frac{d^2 H}{du^2}(u), & \text{if } u = 0. \end{cases}$$

Theorem 2 (Central Limit Theorem for the Density of Particles)

Fix a horizontal time $T > 0$. Consider the Markov process $\{\eta_t : t \geq 0\}$ starting from the invariant state ν_p . Then, the sequence of processes $\{\mathcal{Y}_t^n, 0 \leq t \leq T\}_{n \in \mathbb{N}}$ converges in distribution, as $n \rightarrow +\infty$, with respect to the Skorohod topology of $\mathcal{D}([0, T], \mathcal{S}'_\beta(\mathbb{R}))$ to \mathcal{Y}_t in $\mathcal{C}([0, T], \mathcal{S}'_\beta(\mathbb{R}))$, the generalized Ornstein-Uhlenbeck process which is the formal solution of the SPDE

$$d\mathcal{Y}_t = \Delta_\beta \mathcal{Y}_t dt + \sqrt{2\chi(p)} \nabla_\beta d\mathcal{W}_t \quad (5)$$

where \mathcal{W}_t is a Brownian motion on the space of tempered distributions and $\chi(p) = p(1-p)$ is the *compressibility* of the system.

Remark

- (i) As we said before, the case $\beta > 1$ was proved earlier in [11].
- (ii) The rigorous meaning of (5) is given in terms of martingale problems, see [9, 11] for similar definitions.

3 Strategy of the Proof

- To prove the hydrodynamic limit when $\beta \in [0, 1)$, we use the *entropy method* of Guo, Papanicolaou and Varadhan [15], after a smart trick partitioning the space of configurations.
- To prove the hydrodynamic limit when $\beta = 1$, we appeal to *Gordin's martingale approximation method* [14]. The method was introduced by Gordin in 1969 as an approach to obtain central limit theorems for Markov processes. It basically consists in approximating an additive functional $\int_0^t F(X_s) ds$ of a Markov process to a martingale. The idea is the following: given a Markov process $\{X_t; t \geq 0\}$ with generator \mathcal{L} and a function f in the domain of \mathcal{L} , the process

$$f(X_t) - f(X_0) - \int_0^t \mathcal{L} f(X_s) ds \quad (6)$$

is a mean-zero martingale with respect to the law of $\{X_t; t \geq 0\}$ (see [18, Appendix 1, Lemma 5.1]). Therefore, if we solve the Poisson equation

$$\mathcal{L} f = F \quad (7)$$

for an *appropriate* f , we can express the functional $\int_0^t F(X_s) ds$ as the sum of a martingale and a boundary term. This strategy will be used to prove the *replacement lemmas* for the case $\beta = 1$.

- To prove equilibrium fluctuations for both cases, we maintain the usual *entropy method*, which now works owing to the fact that since we are starting at the stationary measure, we do not need to pay an entropy price.

References

- [1] Baldasso, R., Menezes, O., Neumann, A., and Souza, R. R. *Exclusion process with slow boundary*. Journal of Statistical Physics 167.5 (2017): 1112-1142.
- [2] Beltrán, J., and C. Landim. *A lattice gas model for the incompressible Navier-Stokes equation*. Annales de l'IHP Probabilités et statistiques. Vol. 44. No. 5. 2008.
- [3] De Masi, A., et al. *Current reservoirs in the simple exclusion process*. Journal of Statistical Physics 144.6 (2011): 1151.
- [4] De Masi, A., et al. *Truncated correlations in the stirring process with births and deaths*. Electronic Journal of Probability 17 (2012).
- [5] De Masi, A., Presutti, E. and Tsagkarogiannis, D., Vares, M.E., *Non-equilibrium stationary state for the SEP with births and deaths*. Journal of Statistical Physics (2012).
- [6] Erhard, D., Franco, T., Gonçalves, P., Neumann, A., and Tavares, M. (2018). *Non-equilibrium fluctuations for the SSEP with a slow bond*. arXiv preprint arXiv:1809.04367.
- [7] Franco, T., Gonçalves, P., and Neumann, A. *Equilibrium Fluctuations for the Slow Boundary Exclusion Process*. Meeting on Particle Systems and PDE's. Springer, Cham, 2015.
- [8] Franco, T., Gonçalves, P., and Neumann, A. *Hydrodynamical behavior of symmetric exclusion with slow bonds*. Annales de l'IHP Probabilités et statistiques. Vol. 49. No. 2. 2013.
- [9] Franco, T., Gonçalves, P., and Neumann, A. *Phase transition in equilibrium fluctuations of symmetric slowed exclusion*. Stochastic Processes and their Applications 123.12 (2013): 4156-4185.
- [10] Franco, T., Gonçalves, P. and Neumann, A. *Phase transition of a heat equation with Robin's boundary conditions and exclusion process*. Transactions of the American Mathematical Society 367.9 (2015): 6131-6158.
- [11] Franco, T., Gonçalves, P. and Schütz, G. M. *Scaling limits for the exclusion process with a slow site*. Stochastic Processes and their applications 126.3 (2016): 800-831.
- [12] Franco, T., and Tavares, M. *Hydrodynamic Limit for the SSEP with a Slow Membrane*. arXiv preprint arXiv:1809.07911 (2018).
- [13] Gonçalves, P., Jara, M., Menezes, O., and Neumann, A. (2018). *Non-equilibrium and stationary fluctuations for the SSEP with slow boundary*. arXiv preprint arXiv:1810.05015.
- [14] Gordin, M. I. *The central limit theorem for stationary processes*. Doklady Akademii Nauk. Vol. 188. No. 4. Russian Academy of Sciences, 1969.
- [15] Guo, M. Z., G. C. Papanicolaou, and S. R. S. Varadhan. *Nonlinear diffusion limit for a system with nearest neighbor interactions*. Communications in Mathematical Physics 118.1 (1988): 31-59.
- [16] Jacod, J., and Shiryaev, A. *Limit theorems for stochastic processes*. Springer, 2003.
- [17] Holley, R. A., & Stroock, D. W. (1978). Generalized Ornstein-Uhlenbeck processes and infinite particle branching Brownian motions. Publications of the Research Institute for Mathematical Sciences, 14(3), 741-788.
- [18] Kipnis, C., and Landim, C. *Scaling Limits of Interacting Particle Systems*. Vol. 320. Springer Science and Business Media, 1998.
- [19] Kipnis, C., and Varadhan, S. *Central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusions*. Communications in Mathematical Physics 104.1 (1986): 1-19.
- [20] Liggett, T.M. *Interacting particle systems*. Vol. 276. Springer Science and Business Media, 2012.
- [21] Mitoma, I. (1983). Tightness of Probabilities On $C([0, 1]; \mathcal{S}')$ and $D([0, 1]; \mathcal{S}')$. The Annals of Probability, 11(4), 989-999.
- [22] M. Reed, B. Simon, Functional Analysis, Vol. 1, first ed., in: Methods of Modern Mathematical Physics, Academic Press, 1981.

¹Universidade Federal da Bahia, Brazil, ²Instituto Superior Técnico, Portugal, ³Peking University, China

Email: tertu@ufba.br, patricia.goncalves@math.tecnico.ulisboa.pt, rodrigo.marinho@tecnico.ulisboa.pt and zhaolinjie@pku.edu.cn