

Zero Range Process with Particle Destruction at the Origin

Linjie Zhao

joint work with **Clément Erignoux** and **Marielle Simon**

Inria Lille-Nord Europe, France

October 2021

Hydrodynamic Limit

ZRP

ZRP with Destruction

References

Hydrodynamic Limit

- Hydrodynamic limit: deriving PDEs from microscopic systems, such as interacting particle systems.

Hydrodynamic Limit

- Hydrodynamic limit: deriving PDEs from microscopic systems, such as interacting particle systems.
- The PDEs macroscopically describe the evolution of densities, energies and momentum of the particle system.

Hydrodynamic Limit

- Hydrodynamic limit: deriving PDEs from microscopic systems, such as interacting particle systems.
- The PDEs macroscopically describe the evolution of densities, energies and momentum of the particle system.
- See [Kipnis and Landim'98].

Microscopic/Macroscopic Scale

Let N be the scaling parameter.

	Microscopic	Macroscopic
Space	$x \in \mathbb{Z}^d$	$x/N \in \mathbb{R}^d$
Time	$t\Theta(N)$	t
Mass	1	$1/N^d$

Microscopic/Macroscopic Scale

Let N be the scaling parameter.

	Microscopic	Macroscopic
Space	$x \in \mathbb{Z}^d$	$x/N \in \mathbb{R}^d$
Time	$t\Theta(N)$	t
Mass	1	$1/N^d$

The particles travel a distance of order N during a time interval of order $\Theta(N)$.

Diffusive Scaling

- **Symmetric** case: under diffusive scaling, i.e., space divided by N , time speeded up by N^2 , the hydrodynamic equation is usually parabolic equation.

Diffusive Scaling

- **Symmetric** case: under diffusive scaling, i.e., space divided by N , time speeded up by N^2 , the hydrodynamic equation is usually parabolic equation.
- Recall CLT: $X_i, i \geq 1$, i.i.d. with zero mean and variance one, then

$$\frac{S_{N^2}}{N} \Rightarrow N(0, 1).$$

Diffusive Scaling

- **Symmetric** case: under diffusive scaling, i.e., space divided by N , time speeded up by N^2 , the hydrodynamic equation is usually parabolic equation.
- Recall CLT: X_i , $i \geq 1$, i.i.d. with zero mean and variance one, then

$$\frac{S_{N^2}}{N} \Rightarrow N(0, 1).$$

- Entropy method (GPV method), Yau's relative entropy method...

Hyperbolic Scaling

- **Asymmetric** case: under hyperbolic scaling, i.e., space divided by N , time speeded up by N , the hydrodynamic equation is usually hyperbolic equation.

Hyperbolic Scaling

- **Asymmetric** case: under hyperbolic scaling, i.e., space divided by N , time speeded up by N , the hydrodynamic equation is usually hyperbolic equation.
- Recall LLN: $X_i, i \geq 1$, i.i.d. with finite non-zero mean, then

$$\frac{S_N}{N} \rightarrow E[X_1] \quad \text{in probability.}$$

Hyperbolic Scaling

- **Asymmetric** case: under hyperbolic scaling, i.e., space divided by N , time speeded up by N , the hydrodynamic equation is usually hyperbolic equation.
- Recall LLN: $X_i, i \geq 1$, i.i.d. with finite non-zero mean, then

$$\frac{S_N}{N} \rightarrow E[X_1] \quad \text{in probability.}$$

- Attractiveness method, Fritz's compensated compactness method...

PDEs with Boundary Conditions


- Parabolic equation with boundary conditions: lots of papers.

PDEs with Boundary Conditions

- Parabolic equation with boundary conditions: lots of papers.
- Hyperbolic equation with boundary conditions: [Lamdim'96, Bahadoran'12, Xu'21].

Zero Range Process

Let $1/2 < p \leq 1$ and $g: \{0, 1, 2, \dots\} \rightarrow \mathbb{R}_+$ such that $g(0) = 0$.



Zero Range Process

- State space: $\{0, 1, 2, \dots\}^{\mathbb{Z}}$. For a configuration η , $\eta(x)$ is the number of particles at site x .

Zero Range Process

- State space: $\{0, 1, 2, \dots\}^{\mathbb{Z}}$. For a configuration η , $\eta(x)$ is the number of particles at site x .
- The infinitesimal generator is

$$L_{\text{ZRP}}f(\eta) = \sum_{x \in \mathbb{Z}} \left\{ pg(\eta(x))[f(\eta^{x, x+1}) - f(\eta)] \right. \\ \left. + (1-p)g(\eta(x+1))[f(\eta^{x+1, x}) - f(\eta)] \right\},$$

where

$$\eta^{x,y}(z) = \begin{cases} \eta(x) - 1, & z = x \\ \eta(y) + 1, & z = y \\ \eta(z), & \text{otherwise} \end{cases}$$

Zero Range Process

- The model is introduced by Spitzer in 1970.

Zero Range Process

- The model is introduced by Spitzer in 1970.
- Assume the function g is Lipschitz:

$$\sup_k |g(k+1) - g(k)| \leq \exists C_0.$$

This is needed to well define the process, cf. [Andjel'82].

Zero Range Process

- The model is introduced by Spitzer in 1970.
- Assume the function g is Lipschitz:

$$\sup_k |g(k+1) - g(k)| \leq \exists C_0.$$

This is needed to well define the process, cf. [Andjel'82].

- Attractiveness: $g(k+1) \geq g(k) \geq g(1) > 0$, $k \geq 1$.

Attractiveness

- We could couple processes together such that

$$\omega_0 \leq \eta_0 \quad \Rightarrow \quad \omega_t \leq \eta_t, \quad \forall t.$$

Attractiveness

- We could couple processes together such that

$$\omega_0 \leq \eta_0 \quad \Rightarrow \quad \omega_t \leq \eta_t, \quad \forall t.$$

- Current state:



Attractiveness

- We could couple processes together such that

$$\omega_0 \leq \eta_0 \quad \Rightarrow \quad \omega_t \leq \eta_t, \quad \forall t.$$



- Current state:





- Jump together to the right at rate $pg(1)$:


Attractiveness

- We could couple processes together such that

$$\omega_0 \leq \eta_0 \quad \Rightarrow \quad \omega_t \leq \eta_t, \quad \forall t.$$

- Current state: 

- Jump together to the right at rate $pg(1)$: 
- The red particle jumps alone to the right at rate

$$p(g(2) - g(1)) \geq 0:$$


Hydrodynamic Limit

- The process has a family of invariant measures $\{\nu_\rho\}_{\rho \geq 0}$, where ρ is the particle density. The one-site marginal of ν_ρ is

$$\nu_\rho^1(k) = \frac{1}{Z(\Phi(\rho))} \frac{\Phi(\rho)^k}{g(k)!}, \quad k \geq 0.$$

Hydrodynamic Limit

- The process has a family of invariant measures $\{\nu_\rho\}_{\rho \geq 0}$, where ρ is the particle density. The one-site marginal of ν_ρ is

$$\nu_\rho^1(k) = \frac{1}{Z(\Phi(\rho))} \frac{\Phi(\rho)^k}{g(k)!}, \quad k \geq 0.$$

- Let

$$\Phi(\rho) = \int g(\eta(0)) \nu_\rho(d\eta).$$

The function Φ is strictly increasing.

Hydrodynamic Limit

- The process has a family of invariant measures $\{\nu_\rho\}_{\rho \geq 0}$, where ρ is the particle density. The one-site marginal of ν_ρ is

$$\nu_\rho^1(k) = \frac{1}{Z(\Phi(\rho))} \frac{\Phi(\rho)^k}{g(k)!}, \quad k \geq 0.$$

- Let

$$\Phi(\rho) = \int g(\eta(0)) \nu_\rho(d\eta).$$

The function Φ is strictly increasing.

- Roughly speaking, if

$$\eta_0([uN]) \approx \exists \rho_0(u), \quad u \in \mathbb{R},$$

then

$$\eta_{tN}([uN]) \approx \exists \rho(t, u).$$

Hydrodynamic Limit

➤ Since $\eta_{tN}([uM]) \approx \rho(t, u)$, we have

$$\partial_t \rho(t, u) \approx NL_{\text{ZRP}} \eta_{tN}([uM]).$$

Hydrodynamic Limit

- Since $\eta_{tN}([uM]) \approx \rho(t, u)$, we have

$$\partial_t \rho(t, u) \approx NL_{\text{ZRP}} \eta_{tN}([uM]).$$

- By a simple calculation, $NL_{\text{ZRP}} \eta_{tN}([uM])$ is equal to

$$N \{ \rho g(\eta_{tN}([uM]) - 1) + (1 - \rho) g(\eta_{tN}([uM]) + 1) - g(\eta_{tN}([uM])) \}.$$

Hydrodynamic Limit

- Since $\eta_{tN}([uN]) \approx \rho(t, u)$, we have

$$\partial_t \rho(t, u) \approx NL_{\text{ZRP}} \eta_{tN}([uN]).$$

- By a simple calculation, $NL_{\text{ZRP}} \eta_{tN}([uN])$ is equal to

$$N \{ \rho g(\eta_{tN}([uN]) - 1) + (1 - \rho) g(\eta_{tN}([uN]) + 1) - g(\eta_{tN}([uN])) \}.$$

- Replacing g with its average Φ , the last formula is approximate to

$$\begin{aligned} N(\rho \Phi(\rho(t, u - 1/N)) + (1 - \rho) \Phi(\rho(t, u + 1/N)) - \Phi(\rho(t, u))) \\ \approx -(2\rho - 1) \partial_u \Phi(\rho(t, u)). \end{aligned}$$

Hydrodynamic Limit

- Since $\eta_{tN}([uM]) \approx \rho(t, u)$, we have

$$\partial_t \rho(t, u) \approx NL_{\text{ZRP}} \eta_{tN}([uM]).$$

- By a simple calculation, $NL_{\text{ZRP}} \eta_{tN}([uM])$ is equal to

$$N \{ \rho g(\eta_{tN}([uM]) - 1) + (1 - \rho) g(\eta_{tN}([uM]) + 1) - g(\eta_{tN}([uM])) \}.$$

- Replacing g with its average Φ , the last formula is approximate to

$$\begin{aligned} N(\rho \Phi(\rho(t, u - 1/N)) + (1 - \rho) \Phi(\rho(t, u + 1/N)) - \Phi(\rho(t, u))) \\ \approx -(2\rho - 1) \partial_u \Phi(\rho(t, u)). \end{aligned}$$

- Whence, the hydrodynamic equation should be

$$\partial_t \rho(t, u) + (2\rho - 1) \partial_u \Phi(\rho(t, u)) = 0.$$

Theorem [Rezakhanlou'91]

Fix an initial density profile $\rho_0 : \mathbb{R} \rightarrow \mathbb{R}_+$, which is bounded and locally integrable. Suppose initially, $\{\eta_0(x)\}_{x \in \mathbb{Z}}$ are independent and $\eta_0(x)$ has distribution $\nu_{\rho_0(x/N)}^1$. Then for every time $t > 0$, every $H \in C_c(\mathbb{R})$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{x \in \mathbb{Z}} \eta_{tN}(x) H\left(\frac{x}{N}\right) = \int_{\mathbb{R}} \rho(t, u) H(u) du \quad \text{in probability,}$$

where $\rho(t, u)$ is the unique entropy solution to

$$\begin{cases} \partial_t \rho(t, u) + (2\rho - 1) \partial_u \Phi(\rho(t, u)) = 0, \\ \rho(0, u) = \rho_0(u). \end{cases}$$

Entropy Solution

An entropy solution means

- (i) $\rho(t, u)$ is a weak solution;
- (ii) $\rho(t, u)$ satisfies the entropy inequality: for every $c \geq 0$,

$$\partial_t |\rho(t, u) - c| + (2p - 1) \partial_u |\Phi(\rho(t, u)) - \Phi(c)| \leq 0$$

in the sense of distributions;

- (iii) for every constant $k > 0$,

$$\lim_{t \rightarrow 0} \int_{|u| \leq k} |\rho(t, u) - \rho_0(u)| du = 0.$$

Example: Independent Random Walks

Let $g(k) = k$. Then $\Phi(\rho) = \rho$. The hydrodynamic equation is the simpler transport equation


$$\begin{cases} \partial_t \rho(t, u) + (2p - 1) \partial_u \rho(t, u) = 0, \\ \rho(0, u) = \rho_0(u). \end{cases}$$

The solution is explicitly given by

$$\rho(t, u) = \rho_0(u - (2p - 1)t).$$

ZRP with Destruction

Let $\alpha > 0, \beta \in \mathbb{R}$. N is the scaling parameter.



ZRP with Destruction

➤ The generator is $L_{ZRP} + L_N^{\alpha, \beta}$, where

$$L_N^{\alpha, \beta} f(\eta) = \alpha N^\beta g(\eta(0)) [f(\eta^{(0)}) - f(\eta)],$$

where

$$\eta^{(0)}(z) = \begin{cases} \eta(0) - 1, & z = 0 \\ \eta(z), & \text{otherwise} \end{cases}$$

ZRP with Destruction

- The generator is $L_{ZRP} + L_N^{\alpha, \beta}$, where

$$L_N^{\alpha, \beta} f(\eta) = \alpha N^\beta g(\eta(0)) [f(\eta^{(0)}) - f(\eta)],$$

where

$$\eta^{(0)}(z) = \begin{cases} \eta(0) - 1, & z = 0 \\ \eta(z), & \text{otherwise} \end{cases}$$

- We assume g is Lipschitz and non-decreasing so that the attractiveness property holds.

Hydrodynamic Limit

- The process still has product invariant measures. However, the invariant measures are **not** translation invariant.

Hydrodynamic Limit

- The process still has product invariant measures. However, the invariant measures are **not** translation invariant.
- Initially, $\{\eta_0(x)\}_{x \in \mathbb{Z}}$ are independent and $\eta_0(x)$ has distribution $\nu_{\rho_0(x/N)}^1$ for some bounded and locally integrable initial profile ρ_0 . In particular, every $H \in C_c(\mathbb{R})$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{x \in \mathbb{Z}} \eta_0(x) H\left(\frac{x}{N}\right) = \int_{\mathbb{R}} \rho_0(u) H(u) du \quad \text{in probability.}$$

Hydrodynamic Limit

Theorem for $\beta < 0$, [Erignoux, Simon and Z.'21]

If $\beta < 0$, then the hydrodynamic equation is

$$\begin{cases} \partial_t \rho(t, u) + (2p - 1) \partial_u \Phi(\rho(t, u)) = 0, \\ \rho(0, u) = \rho_0(u). \end{cases}$$

Theorem for $\beta > 0$, [Erignoux, Simon and Z.'21]

If $\beta > 0$, then the hydrodynamic equation is

$$\rho(t, u) = \rho_L(t, u)\mathbf{1}_{\{u < 0\}} + \rho_R(t, u)\mathbf{1}_{\{u \geq 0\}}$$

Theorem for $\beta > 0$, [Erignoux, Simon and Z.'21]

If $\beta > 0$, then the hydrodynamic equation is

$$\rho(t, u) = \rho_L(t, u)\mathbf{1}_{\{u < 0\}} + \rho_R(t, u)\mathbf{1}_{\{u \geq 0\}}$$

where ρ_L is the unique entropy solution to

$$\begin{cases} \partial_t \rho(t, u) + (2p - 1) \partial_u \Phi(\rho(t, u)) = 0, & u \in \mathbb{R} \\ \rho(0, u) = \rho_0(u), \end{cases}$$

Theorem for $\beta > 0$, [Erignoux, Simon and Z.'21]

If $\beta > 0$, then the hydrodynamic equation is

$$\rho(t, u) = \rho_L(t, u)\mathbf{1}_{\{u < 0\}} + \rho_R(t, u)\mathbf{1}_{\{u \geq 0\}}$$

where ρ_L is the unique entropy solution to

$$\begin{cases} \partial_t \rho(t, u) + (2p - 1) \partial_u \Phi(\rho(t, u)) = 0, & u \in \mathbb{R} \\ \rho(0, u) = \rho_0(u), \end{cases}$$

and ρ_R is the unique entropy solution to

$$\begin{cases} \partial_t \rho(t, u) + (2p - 1) \partial_u \Phi(\rho(t, u)) = 0, & u > 0, \\ \rho(t, 0) = 0, \\ \rho(0, u) = \rho_0(u). \end{cases}$$

Theorem for $\beta = 0$, [Erignoux, Simon and Z.'21]

If $\beta = 0$, then the hydrodynamic equation is

$$\rho(t, u) = \rho_L(t, u)\mathbf{1}_{\{u < 0\}} + \rho_R(t, u)\mathbf{1}_{\{u \geq 0\}}$$

where ρ_L is the unique entropy solution to

$$\begin{cases} \partial_t \rho(t, u) + (2p - 1) \partial_u \Phi(\rho(t, u)) = 0, & u \in \mathbb{R} \\ \rho(0, u) = \rho_0(u), \end{cases}$$

and ρ_R is the unique entropy solution to

$$\begin{cases} \partial_t \rho(t, u) + (2p - 1) \partial_u \Phi(\rho(t, u)) = 0, & u > 0, \\ \rho(t, 0) = \Phi^{-1} \left(\frac{2p-1}{\alpha+2p-1} \Phi(\rho_L(t, 0)) \right), \\ \rho(0, u) = \rho_0(u). \end{cases}$$

Entropy Solution

We have two equivalent definitions for

$$\begin{cases} \partial_t \rho(t, u) + (2p - 1) \partial_u \Phi(\rho(t, u)) = 0, & u > 0, \\ \rho(t, 0) = \varrho(t), \\ \rho(0, u) = \rho_0(u). \end{cases}$$

One is based on a weak formula, and the other is based on traces, cf. [Málek *et al.*'19]

Entropy Solution

Definition ($\beta = 0$) [Málek *et al.*'19]

A bounded function ρ is said to be the entropy solution of the above PDE if there exists $M > 0$ such that for every $0 \leq H \in C_c^{1,1}(\mathbb{R} \times \mathbb{R})$ and for every $c \geq 0$,

$$\begin{aligned} & \int_0^\infty \int_0^\infty \left\{ \partial_t H(t, u) (\rho(t, u) - c)^\pm \right. \\ & \quad \left. + (2p - 1) \partial_u H(t, u) (\Phi(\rho(t, u)) - \Phi(c))^\pm \right\} du dt \\ & + M \int_0^\infty H(t, 0) (\varrho(t) - c)^\pm dt + \int_0^\infty H(0, u) (\rho_0(u) - c)^\pm du \geq 0. \end{aligned}$$

Entropy Solution

Definition ($\beta > 0$) [Málek *et al.*'19]

A bounded function ρ is said to be the entropy solution of the above PDE if

- (i) the entropy inequality holds for every $0 \leq H \in C_c^{1,1}((0, \infty) \times (0, \infty))$;
- (ii) for every $T > 0$,

$$\lim_{u \rightarrow 0} \int_0^T |\Phi(\rho(t, u)) - \Phi(\varrho(t))| dt = 0;$$

- (iii) for every constant $k > 0$,


$$\lim_{t \rightarrow 0} \int_{|u| \leq k} |\rho(t, u) - \rho_0(u)| du = 0.$$

Why Using Two Definitions?

- The main idea is to couple the process with another process whose macroscopic profile is the constant c .


Why Using Two Definitions?

- The main idea is to couple the process with another process whose macroscopic profile is the constant c .
- If $\beta = 0$, the stationary profile looks like




Why Using Two Definitions?

- The main idea is to couple the process with another process whose macroscopic profile is the constant c .
- If $\beta = 0$, the stationary profile looks like




- If $\beta > 0$, the stationary profile looks like




Independent Random Walks with Destruction

Let $g(k) = k$. Let us consider the Riemann initial profile:




Independent Random Walks with Destruction

Let $g(k) = k$. Let us consider the Riemann initial profile:




If without destruction or $\beta < 0$, then at time $t > 0$




Independent Random Walks with Destruction

If $\beta > 0$, then at time $t > 0$




Independent Random Walks with Destruction

If $\beta > 0$, then at time $t > 0$



If $\beta = 0$, then at time $t > 0$



Independent Random Walks with Destruction

The solution $\rho(t, u)$ is explicitly given by

$$\rho(t, u) = (1 - \tilde{\alpha} \mathbf{1}\{0 \leq u < (2p - 1)t\}) \rho_0(u - (2p - 1)t),$$

where

$$\tilde{\alpha} = \begin{cases} 1 & \text{if } \beta > 0, \\ \alpha/(\alpha + 2p - 1) & \text{if } \beta = 0, \\ 0 & \text{if } \beta < 0. \end{cases}$$

Future Work

- Remove the assumption that g is increasing. Then the process is non-attractive.

Future Work

- Remove the assumption that g is increasing. Then the process is non-attractive.
- Use compensated compactness method.

Future Work

- Remove the assumption that g is increasing. Then the process is non-attractive.
- Use compensated compactness method.
- Joint work with Clément Erignoux and Lu Xu.



Andjel, E. D. (1982). Invariant measures for the zero range process. *The Annals of Probability*, 10(3), 525-547.



Bahadoran, C. (2012). Hydrodynamics and hydrostatics for a class of asymmetric particle systems with open boundaries. *Communications in Mathematical Physics*, 310(1), 1-24.



Spitzer, F. (1970). Interaction of Markov processes. *Advances in Mathematics*, 5(2), 246-290.



Kipnis, C., & Landim, C. (1998). *Scaling limits of interacting particle systems* (Vol. 320). Springer Science & Business Media.



Landim, C. (1996). Hydrodynamical limit for space inhomogeneous one-dimensional totally asymmetric zero-range processes. *The Annals of Probability*, 599-638.



Málek, J., Nečas, J., Rokyta, M., & Růžička, M. (2019). *Weak and measure-valued solutions to evolutionary PDEs*. Chapman and Hall/CRC.



Xu, L. (2021). Hydrodynamic limit for asymmetric simple exclusion with accelerated boundaries. *arXiv preprint arXiv:2108.09345*.

Thanks!

Email: linjie.zhao@inria.fr