

# STAT 583 Lecture 3

Linjun (Leon) Zhang

Department of Statistics  
Rutgers University

Feb. 4th

- Midterm is scheduled at 6:50-8:50pm on March 3rd.

# Last time

- Point Estimation
- Confidence intervals
- Confidence interval for a population mean ( $\mu$ )

# Recap

- When  $X_1, \dots, X_n$  is an *i.i.d.* sample from a population with known  $\sigma$ , and  $n$  is large,

$$\left[\bar{X} - 2\frac{\sigma}{\sqrt{n}}, \bar{X} + 2\frac{\sigma}{\sqrt{n}}\right]$$

is called an (approximate) 95% confidence interval (CI) for the population mean.

# Recap

- When  $X_1, \dots, X_n$  is an *i.i.d.* sample from a population with known  $\sigma$ , and  $n$  is large,

$$[\bar{X} - 2\frac{\sigma}{\sqrt{n}}, \bar{X} + 2\frac{\sigma}{\sqrt{n}}]$$

is called an (approximate) 95% confidence interval (CI) for the population mean.

- For large  $n$  and unknown  $\sigma$ , we can safely substitute  $s$  for  $\sigma$  to obtain a 95% confidence interval

$$[\bar{X} - 2\frac{s}{\sqrt{n}}, \bar{X} + 2\frac{s}{\sqrt{n}}],$$

where  $s = \sqrt{\frac{(X_1 - \hat{\mu})^2 + \dots + (X_n - \hat{\mu})^2}{n-1}}.$

# Recap

- When  $X_1, \dots, X_n$  is an *i.i.d.* sample from a population with known  $\sigma$ , and  $n$  is large,

$$[\bar{X} - 2\frac{\sigma}{\sqrt{n}}, \bar{X} + 2\frac{\sigma}{\sqrt{n}}]$$

is called an (approximate) 95% confidence interval (CI) for the population mean.

- For large  $n$  and unknown  $\sigma$ , we can safely substitute  $s$  for  $\sigma$  to obtain a 95% confidence interval

$$[\bar{X} - 2\frac{s}{\sqrt{n}}, \bar{X} + 2\frac{s}{\sqrt{n}}],$$

$$\text{where } s = \sqrt{\frac{(X_1 - \hat{\mu})^2 + \dots + (X_n - \hat{\mu})^2}{n-1}}.$$

- The 95% is a property of the procedure, not a specific interval. *In approximately 95% of all samples the confidence interval created according to this procedure will contain  $\mu$ .*

# Recap

A sample of 100 summer days measuring the number of ship passing near a power-plant location showed a mean of 7.2 ships per day with a sample variance of 8.8. Show a 95% CI for the true summer mean.

# Recap

A sample of 100 summer days measuring the number of ship passing near a power-plant location showed a mean of 7.2 ships per day with a sample variance of 8.8. Show a 95% CI for the true summer mean.

## Answer

- $[\bar{X} - 1.96 \frac{s}{\sqrt{n}}, \bar{X} + 1.96 \frac{s}{\sqrt{n}}]$



# Recap

A sample of 100 summer days measuring the number of ship passing near a power-plant location showed a mean of 7.2 ships per day with a sample variance of 8.8. Show a 95% CI for the true summer mean.

## Answer

- $[\bar{X} - 1.96 \frac{s}{\sqrt{n}}, \bar{X} + 1.96 \frac{s}{\sqrt{n}}]$
- $7.2 \pm 1.96 \cdot \sqrt{8.8}/\sqrt{100} = [6.62, 7.78].$

# Interpretation of the 95% Confidence Interval

- The 95% confidence interval for the summer mean is [6.62,7.78].
- What does it MEAN?
- Which of the following interpretations is correct?
  - A With 95% percent chance, the mean  $\mu$  lies inside the interval [6.62,7.78].
  - B Given the observed data, with 95% percent chance, the mean  $\mu$  lies inside the interval [6.62,7.78].
  - C 95% of the values of  $\mu$  lie inside the interval [6.62,7.78].
  - D The interval [6.62,7.78] captures the true value  $\mu$  95% of the time.
  - E None of the above.

# Interpretation of the 95% Confidence Interval

- The correct interpretation:

“In an infinitely long series of trials in which repeated samples of size  $n$  are drawn from the same distribution and 95% CI's for  $\mu$  are calculated using the same **procedure**, the proportion of intervals that actually include  $\mu$  will be 95%. ”

# Interpretation of the 95% Confidence Interval

- The correct interpretation:

“In an infinitely long series of trials in which repeated samples of size  $n$  are drawn from the same distribution and 95% CI's for  $\mu$  are calculated using the same procedure, the proportion of intervals that actually include  $\mu$  will be 95%. ”

- The 95% confidence level is about the procedure, not about any particular interval obtained by applying the method to the observed data

# Interpretation of the 95% Confidence Interval

- The correct interpretation:

“In an infinitely long series of trials in which repeated samples of size  $n$  are drawn from the same distribution and 95% CI's for  $\mu$  are calculated using the same procedure, the proportion of intervals that actually include  $\mu$  will be 95%. ”

- The 95% confidence level is about the procedure, not about any particular interval obtained by applying the method to the observed data
- For any particular CI obtained from the observed data, we do not know whether or not it contains  $\mu$

# Overview

- More on general CIs: sample size determination
- One sample inference
  - CI for a binomial parameter ( $p$ ) when  $n$  is large
  - CI for a population mean ( $\mu$ ) when  $\sigma$  is unknown and  $n$  is small
- Two-sample inference
  - CI for the difference between two binomial parameters  $p_1 - p_2$
  - CI for the difference between two means  $\mu_X - \mu_Y$ 
    - Large sample size
    - Small sample size
    - Paired samples

# Sample size determination

- How large a sample size do I need?

# Sample size determination

- How large a sample size do I need?
- Think about three things:



# Sample size determination

- How large a sample size do I need?
- Think about three things:
  - 1 What confidence level you want (say 95%).

# Sample size determination


- How large a sample size do I need?
- Think about three things:
  - 1 What confidence level you want (say 95%).
  - 2 What Margin of Error (MoE) you want. ( $2\sigma/\sqrt{n}$ )

# Sample size determination

- How large a sample size do I need?
- Think about three things:
  - 1 What confidence level you want (say 95%).
  - 2 What Margin of Error (MoE) you want. ( $2\sigma/\sqrt{n}$ )
  - 3 If  $\sigma$  is unknown, then you need an estimate for  $\sigma$ .

# Sample size determination

- How large a sample size do I need?
- Think about three things:
  - 1 What confidence level you want (say 95%).
  - 2 What Margin of Error (MoE) you want.  $(2\sigma/\sqrt{n})$
  - 3 If  $\sigma$  is unknown, then you need an estimate for  $\sigma$ .
- Recall that the MoE is the distance from the center to the edge of the interval.



# The sample size formula for a population mean, .

- As the  $MoE = 1.96\sigma/\sqrt{n}$  in the 95% confidence interval for  $\mu$ , it follows that:

$$n = \left(\frac{1.96\sigma}{MoE}\right)^2 \approx \left(\frac{2\sigma}{MoE}\right)^2 \approx \left(\frac{2s}{MoE}\right)^2.$$

- There are different formulas for different statistical questions (not just the mean), but sample size is always a legitimate question.

# Example

- An insurance company is being sued because it has paid bills late and then failed to pay interest on the late payments.
- Lawyers need to estimate the average amount of unpaid interest on the late bills.
- The population of bills is 300,000, far too many to review individually.
- How large a sample size do I need to make a 95% confidence interval for the mean amount of unpaid interest on bills paid late?
- We decide on a MoE of  $\pm \$2.00$  and from a previous study we estimate  $\sigma$  by  $s = \$25.25$ .

# Example

- An insurance company is being sued because it has paid bills late and then failed to pay interest on the late payments.
- Lawyers need to estimate the average amount of unpaid interest on the late bills.
- The population of bills is 300,000, far too many to review individually.
- How large a sample size do I need to make a 95% confidence interval for the mean amount of unpaid interest on bills paid late?
- We decide on a MoE of  $\pm \$2.00$  and from a previous study we estimate  $\sigma$  by  $s = \$25.25$ .
- Answer:

$$n = \left( \frac{2 \times 25.25}{2} \right)^2 = 638.$$

# Confidence Intervals for a binomial parameter

Suppose we have a random variable  $Y \sim \text{Bin}(n, p)$ , and we are interested in the CI for  $p$ . Let  $X_i$  be the binary outcome and  $p$  the success probability.

- Recall that  $Y = \sum_{i=1}^n X_i$ , where  $X_1, \dots, X_n$  are i.i.d. Bernoulli distribution with  $\mathbb{P}(X_i = 1) = p, \mathbb{E}(X_i) = p, \text{Var}(X_i) = p(1 - p)$ .



# Confidence Intervals for a binomial parameter

Suppose we have a random variable  $Y \sim \text{Bin}(n, p)$ , and we are interested in the CI for  $p$ . Let  $X_i$  be the binary outcome and  $p$  the success probability.

- Recall that  $Y = \sum_{i=1}^n X_i$ , where  $X_1, \dots, X_n$  are i.i.d. Bernoulli distribution with  $\mathbb{P}(X_i = 1) = p, \mathbb{E}(X_i) = p, \text{Var}(X_i) = p(1 - p)$ .
- We estimate  $p$  by  $\hat{p} = \frac{Y}{n}$ .

# Confidence Intervals for a binomial parameter

Suppose we have a random variable  $Y \sim \text{Bin}(n, p)$ , and we are interested in the CI for  $p$ . Let  $X_i$  be the binary outcome and  $p$  the success probability.

- Recall that  $Y = \sum_{i=1}^n X_i$ , where  $X_1, \dots, X_n$  are i.i.d. Bernoulli distribution with  $\mathbb{P}(X_i = 1) = p$ ,  $\mathbb{E}(X_i) = p$ ,  $\text{Var}(X_i) = p(1 - p)$ .
- We estimate  $p$  by  $\hat{p} = \frac{Y}{n}$ .
- When  $n > 30$  and both  $n\hat{p}$ ,  $n(1 - \hat{p})$  are larger than 10, the sampling distribution of  $\hat{p}$  is approximately:

$$\hat{p} \sim N\left(p, \frac{p(1 - p)}{n}\right).$$

# Confidence Intervals for a binomial parameter

- An approximate 95% CI for the binomial parameter is given by

$$[\hat{p} - 2sd(\hat{p}), \hat{p} + 2sd(\hat{p})] = [\hat{p} - 2\sqrt{\frac{p(1-p)}{n}}, \hat{p} + 2\sqrt{\frac{p(1-p)}{n}}]$$

# Confidence Intervals for a binomial parameter

- An approximate 95% CI for the binomial parameter is given by

$$[\hat{p} - 2sd(\hat{p}), \hat{p} + 2sd(\hat{p})] = [\hat{p} - 2\sqrt{\frac{p(1-p)}{n}}, \hat{p} + 2\sqrt{\frac{p(1-p)}{n}}]$$

- Unfortunately we do not know  $p$  so we replace it with  $\hat{p}$  in the standard deviation calculation (just like replacing  $\sigma$  with  $s$ ).

# Confidence Intervals for a binomial parameter

- An approximate 95% CI for the binomial parameter is given by

$$[\hat{p} - 2sd(\hat{p}), \hat{p} + 2sd(\hat{p})] = [\hat{p} - 2\sqrt{\frac{p(1-p)}{n}}, \hat{p} + 2\sqrt{\frac{p(1-p)}{n}}]$$

- Unfortunately we do not know  $p$  so we replace it with  $\hat{p}$  in the standard deviation calculation (just like replacing  $\sigma$  with  $s$ ).
- So the 95% confidence interval for  $p$  is

$$[\hat{p} - 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}].$$

# Confidence Intervals for a binomial parameter

- An approximate 95% CI for the binomial parameter is given by

$$[\hat{p} - 2sd(\hat{p}), \hat{p} + 2sd(\hat{p})] = [\hat{p} - 2\sqrt{\frac{p(1-p)}{n}}, \hat{p} + 2\sqrt{\frac{p(1-p)}{n}}]$$

- Unfortunately we do not know  $p$  so we replace it with  $\hat{p}$  in the standard deviation calculation (just like replacing  $\sigma$  with  $s$ ).
- So the 95% confidence interval for  $p$  is

$$[\hat{p} - 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}].$$

- In general, the  $(1 - \alpha)$  confidence interval for  $p$  is

$$[\hat{p} - z_{\alpha/2}\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + z_{\alpha/2}\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}].$$

# Example

- A pharmaceutical company needs to compare the performance of its clinical trials to an industry benchmark.
- One way of measuring this is to look at the proportion of trials that move from Phase I to Phase II. The *attrition rate* measures the proportion of trials that fail to make it to Phase II.
- The industry benchmark is 45%.
- The company conducted  $n = 48$  trials, and 30 of them failed.
- Provide a 95% CI for the attrition rate and comment on whether there appears to be a concern with this particular company's trials.

# The attrition data

- There were  $n = 48$  trails.
- Of these, 30 failed, so  $\hat{p} = 0.625$ .
- $n = 48 > 30$ ,  $n\hat{p}$ ,  $n(1 - \hat{p}) > 10$ .



# The attrition data

- There were  $n = 48$  trials.
- Of these, 30 failed, so  $\hat{p} = 0.625$ .
- $n = 48 > 30$ ,  $n\hat{p}$ ,  $n(1 - \hat{p}) > 10$ .
- The approximate 95% CI is given by

$$0.625 \pm 2\sqrt{\frac{0.625 \times (1 - 0.625)}{48}} = 0.625 \pm 2 \times 0.07 = (0.485, 0.765).$$

# The attrition data

- There were  $n = 48$  trials.
- Of these, 30 failed, so  $\hat{p} = 0.625$ .
- $n = 48 > 30$ ,  $n\hat{p}$ ,  $n(1 - \hat{p}) > 10$ .
- The approximate 95% CI is given by

$$0.625 \pm 2\sqrt{\frac{0.625 \times (1 - 0.625)}{48}} = 0.625 \pm 2 \times 0.07 = (0.485, 0.765).$$

- Reporting the interval on a percentage basis gives (48.5%, 76.5%).
- Notice that 45% is not in this interval.
- There is clear evidence that this company is not keeping up with the industry benchmark. Its attrition rate is significantly higher than the benchmark.

# Manipulating confidence intervals

- In many medical settings clinicians like to report results in terms of odds ratios rather than probabilities.

# Manipulating confidence intervals

- In many medical settings clinicians like to report results in terms of odds ratios rather than probabilities.
- If the probability is given by  $p$  then the odds ratio is  $p/(1 - p)$ . A study has found the 95% confidence interval for  $p$  to be  $(0.485, 0.765)$ .

# Manipulating confidence intervals

- In many medical settings clinicians like to report results in terms of odds ratios rather than probabilities.
- If the probability is given by  $p$  then the odds ratio is  $p/(1 - p)$ . A study has found the 95% confidence interval for  $p$  to be  $(0.485, 0.765)$ .
- Find the 95% CI for the odds ratio.

# Manipulating confidence intervals

- In many medical settings clinicians like to report results in terms of odds ratios rather than probabilities.
- If the probability is given by  $p$  then the odds ratio is  $p/(1 - p)$ . A study has found the 95% confidence interval for  $p$  to be (0.485, 0.765).
- Find the 95% CI for the odds ratio.
- Answer:  $(0.485/0.515, 0.765/0.235) = (0.942, 3.255)$ .

# Manipulating confidence intervals

- In many medical settings clinicians like to report results in terms of odds ratios rather than probabilities.
- If the probability is given by  $p$  then the odds ratio is  $p/(1 - p)$ . A study has found the 95% confidence interval for  $p$  to be (0.485, 0.765).
- Find the 95% CI for the odds ratio.
- **Answer:**  $(0.485/0.515, 0.765/0.235) = (0.942, 3.255)$ .
- Remark: the point estimation for the odds ratio is  $\hat{p}/(1 - \hat{p}) = 0.625/0.375 = 1.67$ . CI is not necessarily symmetric around the point estimator.

# Manipulating confidence intervals

- The key idea: you are allowed to transform the ends of the confidence interval to obtain a new confidence interval for the transformed parameter (the transformation must be **monotone** though).



# Manipulating confidence intervals

- The key idea: you are allowed to transform the ends of the confidence interval to obtain a new confidence interval for the transformed parameter (the transformation must be **monotone** though).
- Example: Let  $X$  be the temperature at 9am tomorrow in degree Celsius. You have a 95% CI for  $\mu = \mathbb{E}(X)$ :  $(12^{\circ}\text{C}, 14^{\circ}\text{C})$ .

# Manipulating confidence intervals

- The key idea: you are allowed to transform the ends of the confidence interval to obtain a new confidence interval for the transformed parameter (the transformation must be **monotone** though).
- Example: Let  $X$  be the temperature at 9am tomorrow in degree Celsius. You have a 95% CI for  $\mu = \mathbb{E}(X)$ :  $(12^{\circ}\text{C}, 14^{\circ}\text{C})$ .
- Recall the formula that converts Celsius to Fahrenheit,  $F = 32 + 9C$ .

# Manipulating confidence intervals

- The key idea: you are allowed to transform the ends of the confidence interval to obtain a new confidence interval for the transformed parameter (the transformation must be **monotone** though).
- Example: Let  $X$  be the temperature at 9am tomorrow in degree Celsius. You have a 95% CI for  $\mu = \mathbb{E}(X)$ :  $(12^{\circ}\text{C}, 14^{\circ}\text{C})$ .
- Recall the formula that converts Celsius to Fahrenheit,  $F = 32 + 9C$ .
- Find a 95% CI for  $\mu$  when it is measured in degree Fahrenheit.

# Manipulating confidence intervals

- The key idea: you are allowed to transform the ends of the confidence interval to obtain a new confidence interval for the transformed parameter (the transformation must be **monotone** though).
- Example: Let  $X$  be the temperature at 9am tomorrow in degree Celsius. You have a 95% CI for  $\mu = \mathbb{E}(X)$ :  $(12^{\circ}\text{C}, 14^{\circ}\text{C})$ .
- Recall the formula that converts Celsius to Fahrenheit,  $F = 32 + 9C$ .
- Find a 95% CI for  $\mu$  when it is measured in degree Fahrenheit.
- The interval on the transformed scale is:  
 $(12 \times 9 + 32, 14 \times 9 + 32)^{\circ}\text{F} = (140^{\circ}\text{F}, 158^{\circ}\text{F})$ .

# Example

## Example

A novel coronavirus (2019-nCov) outbreak spreads around the world, especially in Wuhan, China. As of last Saturday, Japan confirmed 8 among 565 citizens evacuated from Wuhan test positive for coronavirus, and Singapore confirmed 2 among 92 evacuees test positive. Assuming the cases are *i.i.d.*, what is the 95% CI for the infection rate of this virus?

# Example

## Example

A novel coronavirus (2019-nCov) outbreak spreads around the world, especially in Wuhan, China. As of last Saturday, Japan confirmed 8 among 565 citizens evacuated from Wuhan test positive for coronavirus, and Singapore confirmed 2 among 92 evacuees test positive. Assuming the cases are *i.i.d.*, what is the 95% CI for the infection rate of this virus?

## Answer

- $\hat{p} = (8 + 2)/(565 + 92) = 1.52\%$

# Example

## Example

A novel coronavirus (2019-nCov) outbreak spreads around the world, especially in Wuhan, China. As of last Saturday, Japan confirmed 8 among 565 citizens evacuated from Wuhan test positive for coronavirus, and Singapore confirmed 2 among 92 evacuees test positive. Assuming the cases are *i.i.d.*, what is the 95% CI for the infection rate of this virus?

## Answer

- $\hat{p} = (8 + 2)/(565 + 92) = 1.52\%$
- $n = 657 > 30, n\hat{p}, n(1 - \hat{p}) \geq 10.$

# Example

## Example

A novel coronavirus (2019-nCov) outbreak spreads around the world, especially in Wuhan, China. As of last Saturday, Japan confirmed 8 among 565 citizens evacuated from Wuhan test positive for coronavirus, and Singapore confirmed 2 among 92 evacuees test positive. Assuming the cases are *i.i.d.*, what is the 95% CI for the infection rate of this virus?

## Answer

- $\hat{p} = (8 + 2)/(565 + 92) = 1.52\%$
- $n = 657 > 30$ ,  $n\hat{p}$ ,  $n(1 - \hat{p}) \geq 10$ .
- The approximate 95% CI is given by

$$1.52\% \pm 2\sqrt{\frac{1.52\% \times (1 - 1.52\%)}{657}} = (0.57\%, 2.47\%).$$



# Example

## Example

A novel coronavirus (2019-nCov) outbreak spreads around the world, especially in Wuhan, China. As of last Saturday, Japan confirmed 8 among 565 citizens evacuated from Wuhan test positive for coronavirus, and Singapore confirmed 2 among 92 evacuees test positive. Assuming the cases are *i.i.d.*, what is the 95% CI for the infection rate of this virus?

## Answer

- $\hat{p} = (8 + 2)/(565 + 92) = 1.52\%$
- $n = 657 > 30$ ,  $n\hat{p}$ ,  $n(1 - \hat{p}) \geq 10$ .
- The approximate 95% CI is given by

$$1.52\% \pm 2\sqrt{\frac{1.52\% \times (1 - 1.52\%)}{657}} = (0.57\%, 2.47\%).$$

- Remark: *i.i.d.* assumption

# A conservative 95% CI for a binomial parameter

- Recall that the 95% confidence interval for  $p$  is

$$[\hat{p} - 2\sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, \hat{p} + 2\sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}]$$

# A conservative 95% CI for a binomial parameter

- Recall that the 95% confidence interval for  $p$  is

$$[\hat{p} - 2\sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, \hat{p} + 2\sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}]$$

- Since  $0 \leq \hat{p} \leq 1$ , we then have  $\hat{p}(1 - \hat{p}) \leq \frac{1}{4}$ , and the equality holds if and only if  $\hat{p} = \frac{1}{2}$ .

# A conservative 95% CI for a binomial parameter

- Recall that the 95% confidence interval for  $p$  is

$$[\hat{p} - 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}]$$

- Since  $0 \leq \hat{p} \leq 1$ , we then have  $\hat{p}(1-\hat{p}) \leq \frac{1}{4}$ , and the equality holds if and only if  $\hat{p} = \frac{1}{2}$ .
- This means  $2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq 2\sqrt{\frac{1}{4n}} = \sqrt{\frac{1}{n}}$ .

# A conservative 95% CI for a binomial parameter

- Recall that the 95% confidence interval for  $p$  is

$$[\hat{p} - 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}]$$

- Since  $0 \leq \hat{p} \leq 1$ , we then have  $\hat{p}(1-\hat{p}) \leq \frac{1}{4}$ , and the equality holds if and only if  $\hat{p} = \frac{1}{2}$ .
- This means  $2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq 2\sqrt{\frac{1}{4n}} = \sqrt{\frac{1}{n}}$ .
- Therefore, a conservative 95% confidence interval for  $p$  is

$$[\hat{p} - \sqrt{\frac{1}{n}}, \hat{p} + \sqrt{\frac{1}{n}}].$$

# Sample size for estimating a binomial parameter

- In the conservative 95% confidence interval for  $p$ ,  $MoE = \sqrt{\frac{1}{n}}$ .

# Sample size for estimating a binomial parameter

- In the conservative 95% confidence interval for  $p$ ,  $MoE = \sqrt{\frac{1}{n}}$ .
- Therefore, a conservative approximation to the sample size formula becomes

$$n = \left(\frac{1}{MoE}\right)^2.$$

# Sample size for estimating a binomial parameter

- In the conservative 95% confidence interval for  $p$ ,  $MoE = \sqrt{\frac{1}{n}}$ .
- Therefore, a conservative approximation to the sample size formula becomes

$$n = \left(\frac{1}{MoE}\right)^2.$$

- You can rely on this approximation if:



# Sample size for estimating a binomial parameter

- In the conservative 95% confidence interval for  $p$ ,  $MoE = \sqrt{\frac{1}{n}}$ .
- Therefore, a conservative approximation to the sample size formula becomes

$$n = \left(\frac{1}{MoE}\right)^2.$$

- You can rely on this approximation if:
  - 1 Your data is *i.i.d.*

# Sample size for estimating a binomial parameter

- In the conservative 95% confidence interval for  $p$ ,  $MoE = \sqrt{\frac{1}{n}}$ .
- Therefore, a conservative approximation to the sample size formula becomes

$$n = \left(\frac{1}{MoE}\right)^2.$$

- You can rely on this approximation if:
  - 1 Your data is *i.i.d.*
  - 2 You want a 95% CI.

# Sample size for estimating a binomial parameter

- In the conservative 95% confidence interval for  $p$ ,  $MoE = \sqrt{\frac{1}{n}}$ .
- Therefore, a conservative approximation to the sample size formula becomes

$$n = \left(\frac{1}{MoE}\right)^2.$$

- You can rely on this approximation if:
  - 1 Your data is *i.i.d.*
  - 2 You want a 95% CI.
  - 3  $p$  lies between 0.25 and 0.75.

## Example

A pollster wants to estimate the proportion of voters who will vote for a given presidential candidate. Assume that the behaviors of voters are *i.i.d.*. How large a sample is required to produce a 95% confidence interval with a margin of error of 3%?

## Example

A pollster wants to estimate the proportion of voters who will vote for a given presidential candidate. Assume that the behaviors of voters are *i.i.d.*. How large a sample is required to produce a 95% confidence interval with a margin of error of 3%?

### Answer

$$n = \left(\frac{1}{MoE}\right)^2 = \left(\frac{1}{3\%}\right)^2 = 1111.$$

## Example

A market research company has estimated the proportion of physicians who say they will prescribe a new drug as 30%. Unfortunately they did not provide a confidence interval. You have since learned that they used a sample size of 100. What was the Margin of Error for a conservative 95% CI? What's the 95% CI?

## Example

A market research company has estimated the proportion of physicians who say they will prescribe a new drug as 30%. Unfortunately they did not provide a confidence interval. You have since learned that they used a sample size of 100. What was the Margin of Error for a conservative 95% CI? What's the 95% CI?

### Answer

The  $MoE = 1/\sqrt{100} = 0.1$ , so the confidence interval is approximately  $(30\% - 10\%, 30\% + 10\%) = (20\%, 40\%)$ .

# Confidence Intervals with unknown $\sigma$

For the CI of  $\mu$ : when the sample size  $n$  is large

- If  $X_1, \dots, X_n$  is an *i.i.d.* sample from a population, the 95% confidence interval (CI) for the population mean  $\mu$  is

$$\left[ \bar{X} - 2 \frac{\sigma}{\sqrt{n}}, \bar{X} + 2 \frac{\sigma}{\sqrt{n}} \right]$$



# Confidence Intervals with unknown $\sigma$

For the CI of  $\mu$ : when the sample size  $n$  is large

- If  $X_1, \dots, X_n$  is an *i.i.d.* sample from a population, the 95% confidence interval (CI) for the population mean  $\mu$  is

$$[\bar{X} - 2\frac{\sigma}{\sqrt{n}}, \bar{X} + 2\frac{\sigma}{\sqrt{n}}]$$

- For large  $n$ , you can substitute  $s$  for  $\sigma$  to obtain a 95% confidence interval

$$[\bar{X} - 2\frac{s}{\sqrt{n}}, \bar{X} + 2\frac{s}{\sqrt{n}}],$$

$$\text{where } s = \sqrt{\frac{(X_1 - \hat{\mu})^2 + \dots + (X_n - \hat{\mu})^2}{n-1}}.$$

## Small sample size: confidence interval for the mean.

- What if you are out of luck and your sample size is not large?

## Small sample size: confidence interval for the mean.

- What if you are out of luck and your sample size is not large?
- Answer: If  $n \leq 30$  then the estimation of  $\sigma$  is not accurate, and the normal approximation is not really appropriate since the CLT speaks only about large sample sizes.

## Small sample size: confidence interval for the mean.

- What if you are out of luck and your sample size is not large?
- Answer: If  $n \leq 30$  then the estimation of  $\sigma$  is not accurate, and the normal approximation is not really appropriate since the CLT speaks only about large sample sizes.
- For large sample size, the Central Limit Theorem told us that

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right).$$

## Small sample size: confidence interval for the mean.

- What if you are out of luck and your sample size is not large?
- Answer: If  $n \leq 30$  then the estimation of  $\sigma$  is not accurate, and the normal approximation is not really appropriate since the CLT speaks only about large sample sizes.
- For large sample size, the Central Limit Theorem told us that

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right).$$

- Equivalently,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1).$$

## Small sample size: confidence interval for the mean.

- For large sample size,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1).$$

## Small sample size: confidence interval for the mean.

- For large sample size,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1).$$

- When we substitute  $s$  instead of  $\sigma$ , we get

$$\frac{\bar{X} - \mu}{s/\sqrt{n}}$$

## Small sample size: confidence interval for the mean.

- For large sample size,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1).$$

- When we substitute  $s$  instead of  $\sigma$ , we get

$$\frac{\bar{X} - \mu}{s/\sqrt{n}}$$

- For large sample sizes, this is close to the standard normal random variable, but not for small  $n$ .



## Small sample size: confidence interval for the mean.

- For large sample size,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1).$$

- When we substitute  $s$  instead of  $\sigma$ , we get

$$\frac{\bar{X} - \mu}{s/\sqrt{n}}$$

- For large sample sizes, this is close to the standard normal random variable, but not for small  $n$ .
- This is because  $s$  is a random variable too and for small  $n$ ,  $s$  is a bad estimate of  $\sigma$ .

## Small sample size: confidence interval for the mean.

- Consider  $\frac{\bar{X} - \mu}{s/\sqrt{n}}$ , where  $X_1, \dots, X_n$  i.i.d.  $\sim P$ .
- For large sample size  $n$ , it's approximately normal distributed
- For small sample size  $n$ ,
  - If the population  $P$  is not a normal distribution, there is nothing we can do at the moment for small sample sizes.
  - We only consider the case when  $P$  is normal.

# Student's t-distribution

- Fact: if  $X_1, \dots, X_n$  *i.i.d.*  $\sim N(\mu, \sigma^2)$ , then

$$T_{n-1} = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{n-1}$$

has a t-distribution (also called Student's t-distribution).

# Student's t-distribution

- Fact: if  $X_1, \dots, X_n$  *i.i.d.*  $\sim N(\mu, \sigma^2)$ , then

$$T_{n-1} = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{n-1}$$

has a t-distribution (also called Student's t-distribution).

- Note that  $t_{n-1}$  depends on  $n$ . The subscript  $n - 1$  denotes the  $n - 1$  **degrees of freedom (d.f.)**, which controls the shape of  $t_{n-1}$ .

# Student's t-distribution

- Fact: if  $X_1, \dots, X_n$  *i.i.d.*  $\sim N(\mu, \sigma^2)$ , then

$$T_{n-1} = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{n-1}$$

has a t-distribution (also called Student's t-distribution).

- Note that  $t_{n-1}$  depends on  $n$ . The subscript  $n - 1$  denotes the  $n - 1$  **degrees of freedom (d.f.)**, which controls the shape of  $t_{n-1}$ .
- The reason why the  $T_{n-1}$  is different from a standard normal  $Z$  is that the  $s$  in the denominator will vary from sample to sample, whereas the  $\sigma$  in the  $Z$  is just a fixed number.

# Student's t-distribution

- Fact: if  $X_1, \dots, X_n$  *i.i.d.*  $\sim N(\mu, \sigma^2)$ , then


$$T_{n-1} = \frac{\bar{X} - \mu}{s/\sqrt{n}} \sim t_{n-1}$$

has a t-distribution (also called Student's t-distribution).

- Note that  $t_{n-1}$  depends on  $n$ . The subscript  $n - 1$  denotes the  $n - 1$  **degrees of freedom (d.f.)**, which controls the shape of  $t_{n-1}$ .
- The reason why the  $T_{n-1}$  is different from a standard normal  $Z$  is that the  $s$  in the denominator will vary from sample to sample, whereas the  $\sigma$  in the  $Z$  is just a fixed number.
- In fact, as  $n$  becomes large,  $t_{n-1}$  converges to a  $N(0, 1)$ .


# Pictures of the Students t-distribution.

**Figure 1:** A t-distribution with 3 degrees of freedom and the standard normal compared in red. The 97.5 percentile is identified



# Pictures of the Students t-distribution.


**Figure 2:** A t-distribution with 10 degrees of freedom and the standard normal compared in red. The 97.5 percentile is identified





# Pictures of the Students t-distribution.

**Figure 3:** A t-distribution with 30 degrees of freedom and the standard normal compared in red. The 97.5 percentile is identified



# Summary of the t-distribution

- $t_{n-1}$  looks like  $N(0, 1)$ , but with fatter tails.

# Summary of the t-distribution

- $t_{n-1}$  looks like  $N(0, 1)$ , but with fatter tails.
- Thus, to get 95% on this fatter distribution, you need to go out further than on the normal distribution.

# Summary of the t-distribution

- $t_{n-1}$  looks like  $N(0, 1)$ , but with fatter tails.
- Thus, to get 95% on this fatter distribution, you need to go out further than on the normal distribution.
- The 97.5 percentile cut-off for the  $t_{n-1}$  is greater than 1.96 (that for  $N(0, 1)$ ).

# Summary of the t-distribution

- $t_{n-1}$  looks like  $N(0, 1)$ , but with fatter tails.
- Thus, to get 95% on this fatter distribution, you need to go out further than on the normal distribution.
- The 97.5 percentile cut-off for the  $t_{n-1}$  is greater than 1.96 (that for  $N(0, 1)$ ).
- t-tables to find these cut-offs are in the back of the textbook.

# Summary of the t-distribution

- $t_{n-1}$  looks like  $N(0, 1)$ , but with fatter tails.
- Thus, to get 95% on this fatter distribution, you need to go out further than on the normal distribution.
- The 97.5 percentile cut-off for the  $t_{n-1}$  is greater than 1.96 (that for  $N(0, 1)$ ).
- t-tables to find these cut-offs are in the back of the textbook.
- For small degrees of freedom, e.g. 3, there can be a large difference between the cutoffs (3.18 v.s. 1.96).

# Summary of the t-distribution

- $t_{n-1}$  looks like  $N(0, 1)$ , but with fatter tails.
- Thus, to get 95% on this fatter distribution, you need to go out further than on the normal distribution.
- The 97.5 percentile cut-off for the  $t_{n-1}$  is greater than 1.96 (that for  $N(0, 1)$ ).
- t-tables to find these cut-offs are in the back of the textbook.
- For small degrees of freedom, e.g. 3, there can be a large difference between the cutoffs (3.18 v.s. 1.96).
- By the time the degrees of freedom reach 30, the difference between cutoffs is very small (2.04 v.s. 1.96).

# Summary of the t-distribution

- $t_{n-1}$  looks like  $N(0, 1)$ , but with fatter tails.
- Thus, to get 95% on this fatter distribution, you need to go out further than on the normal distribution.
- The 97.5 percentile cut-off for the  $t_{n-1}$  is greater than 1.96 (that for  $N(0, 1)$ ).
- t-tables to find these cut-offs are in the back of the textbook.
- For small degrees of freedom, e.g. 3, there can be a large difference between the cutoffs (3.18 v.s. 1.96).
- By the time the degrees of freedom reach 30, the difference between cutoffs is very small (2.04 v.s. 1.96).
- In most practical problems with large sample size, this difference is not important and simply rounding the cut-off to 2 is a reasonable rule of thumb.



# Summary of the t-distribution

- $t_{n-1}$  looks like  $N(0, 1)$ , but with fatter tails.
- Thus, to get 95% on this fatter distribution, you need to go out further than on the normal distribution.
- The 97.5 percentile cut-off for the  $t_{n-1}$  is greater than 1.96 (that for  $N(0, 1)$ ).
- t-tables to find these cut-offs are in the back of the textbook.
- For small degrees of freedom, e.g. 3, there can be a large difference between the cutoffs (3.18 v.s. 1.96).
- By the time the degrees of freedom reach 30, the difference between cutoffs is very small (2.04 v.s. 1.96).
- In most practical problems with large sample size, this difference is not important and simply rounding the cut-off to 2 is a reasonable rule of thumb.
- When software is used to do the calculations, exact cut-off values from the relevant t-distribution are used.

# Summary of the t-distribution

- **Important:** only use the  $t$ -interval if the population distribution (of  $X_i$ ) is normal or approximately normal.

# Summary of the $t$ -distribution

- **Important:** only use the  $t$ -interval if the population distribution (of  $X_i$ ) is normal or approximately normal.
- **Key fact:** for  $n > 30$ ,  $t$ -distribution is very close to the standard normal. For smaller sample size, there is difference, which becomes more and more noticeable for smaller  $n$ .

# Summary of the $t$ -distribution

- **Important:** only use the  $t$ -interval if the population distribution (of  $X_i$ ) is normal or approximately normal.
- **Key fact:** for  $n > 30$ ,  $t$ -distribution is very close to the standard normal. For smaller sample size, there is difference, which becomes more and more noticeable for smaller  $n$ .
- **Rule:** use  $t$ -values whenever  $n < 30$  when you substitute  $s$  for  $\sigma$ , if the population distribution is normal.

# General confidence interval based on the t-distribution

- t-distribution is bell-shaped but has fatter tails.

# General confidence interval based on the t-distribution

- t-distribution is bell-shaped but has fatter tails.
- Recall the  $(1 - \alpha)$  confidence interval for known  $\sigma$ :

$$[\bar{X} - z_{\alpha/2} \cdot \sigma / \sqrt{n}, \bar{X} + z_{\alpha/2} \cdot \sigma / \sqrt{n}].$$

We will call this a z-interval because we rely on the normal distribution.

# General confidence interval based on the t-distribution

- t-distribution is bell-shaped but has fatter tails.
- Recall the  $(1 - \alpha)$  confidence interval for known  $\sigma$ :

$$[\bar{X} - z_{\alpha/2} \cdot \sigma / \sqrt{n}, \bar{X} + z_{\alpha/2} \cdot \sigma / \sqrt{n}].$$

We will call this a z-interval because we rely on the normal distribution.

- When we replace  $\sigma$  with  $s$ , the  $(1 - \alpha)$  confidence interval (which we now call t-interval ) becomes

$$[\bar{X} - t_{\alpha/2, n-1} \cdot s / \sqrt{n}, \bar{X} + t_{\alpha/2, n-1} \cdot s / \sqrt{n}],$$

where  $t_{\alpha/2, n-1}$  is the value such that  $\mathbb{P}(T_{n-1} > t_{\alpha/2, n-1}) = \alpha/2$  for the  $t$ -random variable  $T_{n-1}$  with  $n - 1$  degrees of freedom.

# General confidence interval based on the t-distribution

- t-distribution is bell-shaped but has fatter tails.
- Recall the  $(1 - \alpha)$  confidence interval for known  $\sigma$ :

$$[\bar{X} - z_{\alpha/2} \cdot \sigma / \sqrt{n}, \bar{X} + z_{\alpha/2} \cdot \sigma / \sqrt{n}].$$

We will call this a z-interval because we rely on the normal distribution.

- When we replace  $\sigma$  with  $s$ , the  $(1 - \alpha)$  confidence interval (which we now call t-interval ) becomes

$$[\bar{X} - t_{\alpha/2, n-1} \cdot s / \sqrt{n}, \bar{X} + t_{\alpha/2, n-1} \cdot s / \sqrt{n}],$$

where  $t_{\alpha/2, n-1}$  is the value such that  $\mathbb{P}(T_{n-1} > t_{\alpha/2, n-1}) = \alpha/2$  for the  $t$ -random variable  $T_{n-1}$  with  $n - 1$  degrees of freedom.

- Looks scary, but it's not. The only thing that's different is  $t_{\alpha/2, n-1}$  instead of  $z_{\alpha/2}$ . I will provide you with  $t_{\alpha/2, n-1}$ , so no worries!



# Key Take-away

- For large  $n$ ,  $t_{\alpha/2, n-1} \approx z_{\alpha/2}$ , so we are justified in using  $z$ -values for the case of unknown  $\sigma$ .
- For small  $n$ , however, the  $t$ -value is larger than the corresponding  $z$ -value. Indeed, the tails of the  $t$ -distribution are fatter and the bump at the center is smaller, so you need to go further away from the center to get the same probability.
- Keep in mind that  $t$ -distribution can only be used if the original population (of  $X_i$ 's) is roughly normal.

# Practice Problem

A list of t-values are given below:  $t_{0.05,9} = 1.833$ ,  $t_{0.025,9} = 2.262$ ,  $t_{0.01,9} = 2.821$ .

A sample of 10 observations is drawn from a normal distribution with unknown mean and variance. The sample mean is 12.8 and sample variance  $s^2$  is 4.1. What is a 95% CI for the true mean?

## Practice Problem

A list of t-values are given below:  $t_{0.05,9} = 1.833$ ,  $t_{0.025,9} = 2.262$ ,  $t_{0.01,9} = 2.821$ .

A sample of 10 observations is drawn from a normal distribution with unknown mean and variance. The sample mean is 12.8 and sample variance  $s^2$  is 4.1. What is a 95% CI for the true mean?

### Answer

- $[\bar{X} - t_{\alpha/2, n-1} \cdot s/\sqrt{n}, \bar{X} + t_{\alpha/2, n-1} \cdot s/\sqrt{n}]$

## Practice Problem

A list of t-values are given below:  $t_{0.05,9} = 1.833$ ,  $t_{0.025,9} = 2.262$ ,  $t_{0.01,9} = 2.821$ .

A sample of 10 observations is drawn from a normal distribution with unknown mean and variance. The sample mean is 12.8 and sample variance  $s^2$  is 4.1. What is a 95% CI for the true mean?

### Answer

- $[\bar{X} - t_{\alpha/2, n-1} \cdot s/\sqrt{n}, \bar{X} + t_{\alpha/2, n-1} \cdot s/\sqrt{n}]$
- $12.8 \pm 2.262 \cdot \sqrt{4.1}/\sqrt{10} = [11.35, 14.25]$ .

# Summary of Confidence intervals

- A general formula for CI of mean-based parameters. For a parameter  $\theta$ , suppose we estimate it by  $\hat{\theta}$  and  $\hat{\theta}$  is approximated normal. Then a  $(1 - \alpha)$  confidence interval for  $\theta$  is

$$[\hat{\theta} - z_{\alpha/2} \cdot \widehat{\text{sd}}(\hat{\theta}), \quad \hat{\theta} + z_{\alpha/2} \cdot \widehat{\text{sd}}(\hat{\theta})]$$

where  $\text{sd}(\hat{\theta})$  is the standard deviation of  $\hat{\theta}$ , and  $\widehat{\text{sd}}(\hat{\theta})$  is the estimate of  $\text{sd}(\hat{\theta})$ .

- In the setting where the sample size is small and the observations are normally distributed, we replace  $z_{\alpha/2}$  with  $t_{\alpha/2, n-1}$ .

# Confidence intervals

- When  $n \geq 30$ , and  $n\hat{p}, n(1 - \hat{p}) > 10$ , if  $X$  has a binomial distribution  $\text{Bin}(n, p)$ , then the estimate of the binomial parameter  $p$  is  $\hat{p} = \frac{X}{n}$

# Confidence intervals

- When  $n \geq 30$ , and  $n\hat{p}, n(1 - \hat{p}) > 10$ , if  $X$  has a binomial distribution  $\text{Bin}(n, p)$ , then the estimate of the binomial parameter  $p$  is  $\hat{p} = \frac{X}{n}$
- $sd(\hat{p}) = \frac{p(1-p)}{n}$

# Confidence intervals

- When  $n \geq 30$ , and  $n\hat{p}, n(1 - \hat{p}) > 10$ , if  $X$  has a binomial distribution  $\text{Bin}(n, p)$ , then the estimate of the binomial parameter  $p$  is  $\hat{p} = \frac{X}{n}$
- $sd(\hat{p}) = \frac{p(1-p)}{n}$
- $\widehat{sd(\hat{p})} = \frac{\hat{p}(1-\hat{p})}{n}$



# Confidence intervals

- When  $n \geq 30$ , and  $n\hat{p}, n(1 - \hat{p}) > 10$ , if  $X$  has a binomial distribution  $Bin(n, p)$ , then the estimate of the binomial parameter  $p$  is  $\hat{p} = \frac{X}{n}$
- $sd(\hat{p}) = \frac{p(1-p)}{n}$
- $\widehat{sd}(\hat{p}) = \frac{\hat{p}(1-\hat{p})}{n}$
- An approximate 95% confidence interval for  $p$  is

$$\left[ \hat{p} - 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \right]$$

# Confidence intervals

- When  $n \geq 30$ , and  $n\hat{p}, n(1 - \hat{p}) > 10$ , if  $X$  has a binomial distribution  $\text{Bin}(n, p)$ , then the estimate of the binomial parameter  $p$  is  $\hat{p} = \frac{X}{n}$
- $sd(\hat{p}) = \frac{p(1-p)}{n}$
- $\widehat{sd}(\hat{p}) = \frac{\hat{p}(1-\hat{p})}{n}$
- An approximate 95% confidence interval for  $p$  is

$$[\hat{p} - 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + 2\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}]$$

- A conservative 95% confidence interval for  $p$  is

$$[\hat{p} - \sqrt{\frac{1}{n}}, \hat{p} + \sqrt{\frac{1}{n}}].$$

# Confidence intervals

- When the sample size  $n > 30$ , if  $X_1, X_2, \dots, X_n$  are i.i.d. with mean  $\mu$  and variance  $\sigma^2$ , then an estimate of  $\mu$  is  $\hat{\mu} = \bar{X} = \frac{X_1 + \dots + X_n}{n}$ .
- $sd(\bar{X}) = \frac{\sigma}{\sqrt{n}}$
- $\widehat{sd}(\bar{X}) = \frac{s}{\sqrt{n}}$ , where  $s^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$
- An approximate 95% confidence interval for  $\mu$  is

$$[\bar{X} - 2\frac{s}{\sqrt{n}}, \bar{X} + 2\frac{s}{\sqrt{n}}].$$

# Confidence intervals

- When the sample size is small, if  $X_1, X_2, \dots, X_n$  are i.i.d.  $N(\mu, \sigma^2)$ , then an estimate of  $\mu$  is still  $\hat{\mu} = \bar{X} = \frac{X_1 + \dots + X_n}{n}$ .

# Confidence intervals

- When the sample size is small, if  $X_1, X_2, \dots, X_n$  are i.i.d.  $N(\mu, \sigma^2)$ , then an estimate of  $\mu$  is still  $\hat{\mu} = \bar{X} = \frac{X_1 + \dots + X_n}{n}$ .
- $sd(\bar{X}) = \frac{\sigma}{\sqrt{n}}$

# Confidence intervals

- When the sample size is small, if  $X_1, X_2, \dots, X_n$  are i.i.d.  $N(\mu, \sigma^2)$ , then an estimate of  $\mu$  is still  $\hat{\mu} = \bar{X} = \frac{X_1 + \dots + X_n}{n}$ .
- $sd(\bar{X}) = \frac{\sigma}{\sqrt{n}}$
- $\widehat{sd}(\bar{X}) = \frac{s}{\sqrt{n}}$ , where  $s^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$

# Confidence intervals

- When the sample size is small, if  $X_1, X_2, \dots, X_n$  are i.i.d.  $N(\mu, \sigma^2)$ , then an estimate of  $\mu$  is still  $\hat{\mu} = \bar{X} = \frac{X_1 + \dots + X_n}{n}$ .
- $sd(\bar{X}) = \frac{\sigma}{\sqrt{n}}$
- $\widehat{sd}(\bar{X}) = \frac{s}{\sqrt{n}}$ , where  $s^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$
- A 95% confidence interval for  $\mu$  is

$$[\bar{X} - t_{0.025, n-1} \cdot \frac{s}{\sqrt{n}}, \bar{X} + t_{0.025, n-1} \cdot \frac{s}{\sqrt{n}}].$$

# CI for the difference between two binomial parameters

- When  $n, m$  are sufficiently large, if  $X$  has a binomial distribution  $Bi(n, p_1)$ , and  $Y$  has a binomial distribution  $Bi(m, p_2)$ .  $X$  and  $Y$  are independent.



# CI for the difference between two binomial parameters

- When  $n, m$  are sufficiently large, if  $X$  has a binomial distribution  $Bi(n, p_1)$ , and  $Y$  has a binomial distribution  $Bi(m, p_2)$ .  $X$  and  $Y$  are independent.
- An estimate of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = \frac{X}{n} - \frac{Y}{m}$ .

# CI for the difference between two binomial parameters

- When  $n, m$  are sufficiently large, if  $X$  has a binomial distribution  $Bi(n, p_1)$ , and  $Y$  has a binomial distribution  $Bi(m, p_2)$ .  $X$  and  $Y$  are independent.
- An estimate of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = \frac{X}{n} - \frac{Y}{m}$ .
- $sd(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{p_1(1-p_1)}{n} + \frac{p_2(1-p_2)}{m}}$

# CI for the difference between two binomial parameters

- When  $n, m$  are sufficiently large, if  $X$  has a binomial distribution  $Bi(n, p_1)$ , and  $Y$  has a binomial distribution  $Bi(m, p_2)$ .  $X$  and  $Y$  are independent.
- An estimate of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = \frac{X}{n} - \frac{Y}{m}$ .
- $sd(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{p_1(1-p_1)}{n} + \frac{p_2(1-p_2)}{m}}$
- $\widehat{sd(\hat{p}_1 - \hat{p}_2)} = \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}$

# CI for the difference between two binomial parameters

- When  $n, m$  are sufficiently large, if  $X$  has a binomial distribution  $Bi(n, p_1)$ , and  $Y$  has a binomial distribution  $Bi(m, p_2)$ .  $X$  and  $Y$  are independent.
- An estimate of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = \frac{X}{n} - \frac{Y}{m}$ .
- $sd(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{p_1(1-p_1)}{n} + \frac{p_2(1-p_2)}{m}}$
- $\widehat{sd(\hat{p}_1 - \hat{p}_2)} = \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}$
- An approximate 95% confidence interval for  $p_1 - p_2$  is

$$[\hat{p}_1 - \hat{p}_2 - 2\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}, \hat{p}_1 - \hat{p}_2 + 2\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}].$$

# CI for the difference between two binomial parameters

- When  $n, m$  are sufficiently large, if  $X$  has a binomial distribution  $Bi(n, p_1)$ , and  $Y$  has a binomial distribution  $Bi(m, p_2)$ .  $X$  and  $Y$  are independent.

- An estimate of  $p_1 - p_2$  is  $\hat{p}_1 - \hat{p}_2 = \frac{X}{n} - \frac{Y}{m}$ .

- $sd(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{p_1(1-p_1)}{n} + \frac{p_2(1-p_2)}{m}}$

- $\widehat{sd}(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}$

- An approximate 95% confidence interval for  $p_1 - p_2$  is

$$[\hat{p}_1 - \hat{p}_2 - 2\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}, \hat{p}_1 - \hat{p}_2 + 2\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}].$$

- A conservative 95% confidence interval for  $p_1 - p_2$  is

$$[\hat{p}_1 - \hat{p}_2 - \sqrt{\frac{1}{n} + \frac{1}{m}}, \hat{p}_1 - \hat{p}_2 + \sqrt{\frac{1}{n} + \frac{1}{m}}].$$

- $X \sim \text{Bi}(n, p_1)$ ,  $Y \sim \text{Bi}(m, p_2)$

$$\begin{aligned}sd(\hat{p}_1 - \hat{p}_2) &= \sqrt{\text{Var}(\hat{p}_1 - \hat{p}_2)} = \sqrt{\text{Var}\left(\frac{X}{n} - \frac{Y}{m}\right)} \\&= \sqrt{\text{Var}\left(\frac{X}{n}\right) + \text{Var}\left(\frac{Y}{m}\right)} \\&= \sqrt{\frac{1}{n^2} \cdot np_1(1 - p_1) + \frac{1}{m^2} \cdot mp_2(1 - p_2)} \\&= \sqrt{\frac{p_1(1 - p_1)}{n} + \frac{p_2(1 - p_2)}{m}}\end{aligned}$$

- $2\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}} \leq 2\sqrt{\frac{1}{4n} + \frac{1}{4m}} = \sqrt{\frac{1}{n} + \frac{1}{m}}.$

# Practice problem

## Question

We have two drugs, drug  $A$  and drug  $B$ , both aimed at curing a certain illness. We wish to compare the two drugs. Define  $p_1$  as the probability that a person with this illness will be cured if he/she takes drug  $A$ , and  $p_2$  as the probability that a person with this illness will be cured if he/she takes drug  $B$ . We give drug  $A$  to 1,000 people and it cures 840 of them. We give drug  $B$  to 1,200 people and it cures 970 of them. Find the 95% CI for  $p_1 - p_2$ .

# Practice problem

## Question

We have two drugs, drug  $A$  and drug  $B$ , both aimed at curing a certain illness. We wish to compare the two drugs. Define  $p_1$  as the probability that a person with this illness will be cured if he/she takes drug  $A$ , and  $p_2$  as the probability that a person with this illness will be cured if he/she takes drug  $B$ . We give drug  $A$  to 1,000 people and it cures 840 of them. We give drug  $B$  to 1,200 people and it cures 970 of them. Find the 95% CI for  $p_1 - p_2$ .

## Solution

$\hat{p}_1 = 0.84$ ,  $\hat{p}_2 = 0.81$ ,  $n = 1000$ ,  $m = 1200$ . Hence the 95% confidence interval is  $0.03 \pm \sqrt{\frac{1}{1200} + \frac{1}{1000}}$ , which is -0.01 to 0.07.



## CI for $\mu_X - \mu_Y$ when $n, m$ are large

If  $X_1, X_2, \dots, X_n$  are i.i.d. with mean  $\mu_X$  and variance  $\sigma_X^2$ ,  $Y_1, Y_2, \dots, Y_m$  are i.i.d. with mean  $\mu_Y$  and variance  $\sigma_Y^2$ .  $\{X_1, \dots, X_n\}$  and  $\{Y_1, \dots, Y_m\}$  are independent.

- We estimate  $\mu_X - \mu_Y$  by  $\bar{X} - \bar{Y}$ , where  $\bar{X} = \frac{X_1 + \dots + X_n}{n}$ ,  $\bar{Y} = \frac{Y_1 + \dots + Y_m}{m}$ .

## CI for $\mu_X - \mu_Y$ when $n, m$ are large

If  $X_1, X_2, \dots, X_n$  are i.i.d. with mean  $\mu_X$  and variance  $\sigma_X^2$ ,  $Y_1, Y_2, \dots, Y_m$  are i.i.d. with mean  $\mu_Y$  and variance  $\sigma_Y^2$ .  $\{X_1, \dots, X_n\}$  and  $\{Y_1, \dots, Y_m\}$  are independent.

- We estimate  $\mu_X - \mu_Y$  by  $\bar{X} - \bar{Y}$ , where  $\bar{X} = \frac{X_1 + \dots + X_n}{n}$ ,  $\bar{Y} = \frac{Y_1 + \dots + Y_m}{m}$ .
- The variance of  $\bar{X} - \bar{Y}$  is  $\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}$

## CI for $\mu_X - \mu_Y$ when $n, m$ are large

If  $X_1, X_2, \dots, X_n$  are i.i.d. with mean  $\mu_X$  and variance  $\sigma_X^2$ ,  $Y_1, Y_2, \dots, Y_m$  are i.i.d. with mean  $\mu_Y$  and variance  $\sigma_Y^2$ .  $\{X_1, \dots, X_n\}$  and  $\{Y_1, \dots, Y_m\}$  are independent.

- We estimate  $\mu_X - \mu_Y$  by  $\bar{X} - \bar{Y}$ , where  $\bar{X} = \frac{X_1 + \dots + X_n}{n}$ ,  $\bar{Y} = \frac{Y_1 + \dots + Y_m}{m}$ .
- The variance of  $\bar{X} - \bar{Y}$  is  $\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}$
- We estimate  $\sigma_X^2$  and  $\sigma_Y^2$  by  $s_X^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$ ,  $s_Y^2 = \frac{(Y_1 - \bar{Y})^2 + \dots + (Y_m - \bar{Y})^2}{m-1}$ .

## CI for $\mu_X - \mu_Y$ when $n, m$ are large

If  $X_1, X_2, \dots, X_n$  are i.i.d. with mean  $\mu_X$  and variance  $\sigma_X^2$ ,  $Y_1, Y_2, \dots, Y_m$  are i.i.d. with mean  $\mu_Y$  and variance  $\sigma_Y^2$ .  $\{X_1, \dots, X_n\}$  and  $\{Y_1, \dots, Y_m\}$  are independent.

- We estimate  $\mu_X - \mu_Y$  by  $\bar{X} - \bar{Y}$ , where  $\bar{X} = \frac{X_1 + \dots + X_n}{n}$ ,  $\bar{Y} = \frac{Y_1 + \dots + Y_m}{m}$ .
- The variance of  $\bar{X} - \bar{Y}$  is  $\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}$
- We estimate  $\sigma_X^2$  and  $\sigma_Y^2$  by  $s_X^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$ ,  $s_Y^2 = \frac{(Y_1 - \bar{Y})^2 + \dots + (Y_m - \bar{Y})^2}{m-1}$ .
- A  $(1 - \alpha)$  confidence interval for  $\mu_X - \mu_Y$  is

$$[\bar{X} - \bar{Y} - z_{\alpha/2} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}, \bar{X} - \bar{Y} + z_{\alpha/2} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}].$$

$$\begin{aligned}sd(\hat{\mu}_X - \hat{\mu}_Y) &= \sqrt{\text{Var}(\hat{\mu}_X - \hat{\mu}_Y)} = \sqrt{\text{Var}(\bar{X} - \bar{Y})} \\&= \sqrt{\text{Var}(\bar{X}) + \text{Var}(\bar{Y})} \\&= \sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}\end{aligned}$$

# Practice problem

A psychologist was interested in exploring whether or not male and female college students have different driving behaviors. She focused on the fastest speed ever driven by an individual. She conducted a survey of a random  $n = 34$  male college students and a random  $m = 30$  female college students. Here is a descriptive summary of the results of her survey: for male students, the average of their fastest speeds is 105 miles per hour (mph), and the standard deviation is 20.1 mph; for female students, the average is 90.9 mph and the standard deviation is 12.2 mph. Construct a 95% CI for the difference of the fastest speed ever driven by male and female college students.

## Practice problem

A psychologist was interested in exploring whether or not male and female college students have different driving behaviors. She focused on the fastest speed ever driven by an individual. She conducted a survey of a random  $n = 34$  male college students and a random  $m = 30$  female college students. Here is a descriptive summary of the results of her survey: for male students, the average of their fastest speeds is 105 miles per hour (mph), and the standard deviation is 20.1 mph; for female students, the average is 90.9 mph and the standard deviation is 12.2 mph. Construct a 95% CI for the difference of the fastest speed ever driven by male and female college students.

### Answer

- $$[\bar{X} - \bar{Y} - z_{\alpha/2} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}, \quad \bar{X} - \bar{Y} + z_{\alpha/2} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}]$$

## Practice problem

A psychologist was interested in exploring whether or not male and female college students have different driving behaviors. She focused on the fastest speed ever driven by an individual. She conducted a survey of a random  $n = 34$  male college students and a random  $m = 30$  female college students. Here is a descriptive summary of the results of her survey: for male students, the average of their fastest speeds is 105 miles per hour (mph), and the standard deviation is 20.1 mph; for female students, the average is 90.9 mph and the standard deviation is 12.2 mph. Construct a 95% CI for the difference of the fastest speed ever driven by male and female college students.

### Answer

- $[\bar{X} - \bar{Y} - z_{\alpha/2} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}, \bar{X} - \bar{Y} + z_{\alpha/2} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}]$
- $(105 - 90.9) \pm 2 \sqrt{\frac{20.1^2}{34} + \frac{12.2^2}{30}} = [5.89, 22.31] \text{ mph}$



## CI for $\mu_X - \mu_Y$ when $n, m$ are small

When  $n, m$  are small. If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ .  $\{X_1, \dots, X_n\}$  and  $\{Y_1, \dots, Y_m\}$  are independent.

- We estimate  $\mu_X - \mu_Y$  by  $\bar{X} - \bar{Y}$ , where  $\bar{X} = \frac{X_1 + \dots + X_n}{n}$ ,  $\bar{Y} = \frac{Y_1 + \dots + Y_m}{m}$ .

## CI for $\mu_X - \mu_Y$ when $n, m$ are small

When  $n, m$  are small. If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ .  $\{X_1, \dots, X_n\}$  and  $\{Y_1, \dots, Y_m\}$  are independent.

- We estimate  $\mu_X - \mu_Y$  by  $\bar{X} - \bar{Y}$ , where  $\bar{X} = \frac{X_1 + \dots + X_n}{n}$ ,  $\bar{Y} = \frac{Y_1 + \dots + Y_m}{m}$ .
- The variance of  $\bar{X} - \bar{Y}$  is  $\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}$ , and we estimate  $\sigma_X^2$  and  $\sigma_Y^2$  by  $s_X^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$ ,  $s_Y^2 = \frac{(Y_1 - \bar{Y})^2 + \dots + (Y_m - \bar{Y})^2}{m-1}$ .

## CI for $\mu_X - \mu_Y$ when $n, m$ are small

When  $n, m$  are small. If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ .  $\{X_1, \dots, X_n\}$  and  $\{Y_1, \dots, Y_m\}$  are independent.

- We estimate  $\mu_X - \mu_Y$  by  $\bar{X} - \bar{Y}$ , where  $\bar{X} = \frac{X_1 + \dots + X_n}{n}$ ,  $\bar{Y} = \frac{Y_1 + \dots + Y_m}{m}$ .
- The variance of  $\bar{X} - \bar{Y}$  is  $\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}$ , and we estimate  $\sigma_X^2$  and  $\sigma_Y^2$  by  $s_X^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$ ,  $s_Y^2 = \frac{(Y_1 - \bar{Y})^2 + \dots + (Y_m - \bar{Y})^2}{m-1}$ .
- We then have  $\frac{\bar{X} - \bar{Y} - (\mu_X - \mu_Y)}{\sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}} \approx T_k$ , where  $k = \min\{n, m\} - 1$

## CI for $\mu_X - \mu_Y$ when $n, m$ are small

When  $n, m$  are small. If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ .  $\{X_1, \dots, X_n\}$  and  $\{Y_1, \dots, Y_m\}$  are independent.

- We estimate  $\mu_X - \mu_Y$  by  $\bar{X} - \bar{Y}$ , where  $\bar{X} = \frac{X_1 + \dots + X_n}{n}$ ,  $\bar{Y} = \frac{Y_1 + \dots + Y_m}{m}$ .
- The variance of  $\bar{X} - \bar{Y}$  is  $\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}$ , and we estimate  $\sigma_X^2$  and  $\sigma_Y^2$  by  $s_X^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$ ,  $s_Y^2 = \frac{(Y_1 - \bar{Y})^2 + \dots + (Y_m - \bar{Y})^2}{m-1}$ .
- We then have  $\frac{\bar{X} - \bar{Y} - (\mu_X - \mu_Y)}{\sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}} \approx T_k$ , where  $k = \min\{n, m\} - 1$
- A  $(1 - \alpha)$  confidence interval for  $\mu_X - \mu_Y$  is

$$[\bar{X} - \bar{Y} - t_{\alpha/2, k} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}, \bar{X} - \bar{Y} + t_{\alpha/2, k} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}].$$

# Practice problem

We are interested in investigating any potential difference between the mean blood sugar level of diabetics ( $\mu_X$ ) and that of non-diabetics ( $\mu_Y$ ). To do this we took a sample of six diabetics and found the following blood sugar levels: 127, 144, 140, 136, 118, 138. We also took a sample of eight non-diabetics and found the following blood sugar levels: 125, 128, 133, 141, 109, 125, 126, 122. (a) Estimate  $\mu_X - \mu_Y$ . (b) Find the 95% CI for  $\mu_X - \mu_Y$ .

$$(s_X^2 = 93.24, s_Y^2 = 83.55, t_{0.05,5} = 2.015, t_{0.025,5} = 2.571, t_{0.05,7} = 1.895, t_{0.025,7} = 2.365)$$

## Practice problem

We are interested in investigating any potential difference between the mean blood sugar level of diabetics ( $\mu_X$ ) and that of non-diabetics ( $\mu_Y$ ). To do this we took a sample of six diabetics and found the following blood sugar levels: 127, 144, 140, 136, 118, 138. We also took a sample of eight non-diabetics and found the following blood sugar levels: 125, 128, 133, 141, 109, 125, 126, 122. (a) Estimate  $\mu_X - \mu_Y$ . (b) Find the 95% CI for  $\mu_X - \mu_Y$ .

$$(s_X^2 = 93.24, s_Y^2 = 83.55, t_{0.05,5} = 2.015, t_{0.025,5} = 2.571, t_{0.05,7} = 1.895, t_{0.025,7} = 2.365)$$

### Solution

$$\hat{\mu}_X = \bar{X} = \frac{1}{6}(127 + 144 + 140 + 136 + 118 + 138) = 133.83.$$

$$\hat{\sigma}_X^2 = s_X^2 = \frac{1}{6-1}(127^2 + 144^2 + 140^2 + 136^2 + 118^2 + 138^2 - 6 \times 133.83^2) = 93.24.$$

$$\hat{\mu}_Y = \bar{Y} = \frac{1}{8}(125 + 128 + 133 + 141 + 109 + 125 + 126 + 122) = 126.13.$$

$$\hat{\sigma}_Y^2 = s_Y^2 = \frac{1}{8-1}(125^2 + 128^2 + 133^2 + 141^2 + 109^2 + 125^2 + 126^2 + 122^2 - 8 \times 126.13^2) = 83.55.$$

# Practice problem

We are interested in investigating any potential difference between the mean blood sugar level of diabetics ( $\mu_X$ ) and that of non-diabetics ( $\mu_Y$ ). To do this we took a sample of six diabetics and found the following blood sugar levels: 127, 144, 140, 136, 118, 138. We also took a sample of eight non-diabetics and found the following blood sugar levels: 125, 128, 133, 141, 109, 125, 126, 122. (a) Estimate  $\mu_X - \mu_Y$ . (b) Find the 95% CI for  $\mu_X - \mu_Y$ .

$$(s_X^2 = 93.24, s_Y^2 = 83.55, t_{0.05,5} = 2.015, t_{0.025,5} = 2.571, t_{0.05,7} = 1.895, t_{0.025,7} = 2.365)$$

## Solution

$$\hat{\mu}_X = \bar{X} = \frac{1}{6}(127 + 144 + 140 + 136 + 118 + 138) = 133.83.$$

$$\hat{\sigma}_X^2 = s_X^2 = \frac{1}{6-1}(127^2 + 144^2 + 140^2 + 136^2 + 118^2 + 138^2 - 6 \times 133.83^2) = 93.24.$$

$$\hat{\mu}_Y = \bar{Y} = \frac{1}{8}(125 + 128 + 133 + 141 + 109 + 125 + 126 + 122) = 126.13.$$

$$\hat{\sigma}_Y^2 = s_Y^2 = \frac{1}{8-1}(125^2 + 128^2 + 133^2 + 141^2 + 109^2 + 125^2 + 126^2 + 122^2 - 8 \times 126.13^2) = 83.55.$$

The 95% confidence interval is given as

$$\bar{X} - \bar{Y} \pm t_{0.025,5} \sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}} = (133.83 - 126.13) \pm 2.571 \sqrt{\frac{93.24}{6} + \frac{83.55}{8}} = [-5.41, 20.81]$$

$$s^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n - 1} = \frac{X_1^2 + \dots + X_n^2 - n(\bar{X})^2}{n - 1}$$

Proof:

$$\begin{aligned} s^2 &= \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n - 1} = \frac{\sum_{i=1}^n (X_i^2 + (\bar{X})^2 - 2X_i \cdot \bar{X})}{n - 1} \\ &= \frac{\sum_{i=1}^n X_i^2 + \sum_{i=1}^n (\bar{X})^2 - \sum_{i=1}^n 2X_i \cdot \bar{X}}{n - 1} \\ &= \frac{\sum_{i=1}^n X_i^2 + n(\bar{X})^2 - 2n(\bar{X})^2}{n - 1} \\ &= \frac{X_1^2 + \dots + X_n^2 - n(\bar{X})^2}{n - 1}. \end{aligned}$$



## When the variances are the same

If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma^2)$ ,  $Y_1, Y_2, \dots, Y_m$  i.i.d.  $\sim N(\mu_Y, \sigma^2)$

- If it is known that both populations have the same variance, then we can leverage this information to get a more accurate estimate of by combining both  $s_X$  and  $s_Y$  to create what is called a pooled estimate of the variance:

$$s_{pool}^2 = \frac{(n-1)s_X^2 + (m-1)s_Y^2}{n+m-2},$$

$$\text{where } s_X^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}, \quad s_Y^2 = \frac{(Y_1 - \bar{Y})^2 + \dots + (Y_m - \bar{Y})^2}{m-1}.$$

# When the variances are the same

If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma^2)$ ,  $Y_1, Y_2, \dots, Y_n$  i.i.d.  $\sim N(\mu_Y, \sigma^2)$

- If it is known that both populations have the same variance, then we can leverage this information to get a more accurate estimate of by combining both  $s_X$  and  $s_Y$  to create what is called a pooled estimate of the variance:

$$s_{pool}^2 = \frac{(n-1)s_X^2 + (m-1)s_Y^2}{n+m-2},$$

where  $s_X^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$ ,  $s_Y^2 = \frac{(Y_1 - \bar{Y})^2 + \dots + (Y_m - \bar{Y})^2}{m-1}$ .

- The pooled estimate is a weighted average of the individual sample estimates. This increases the sample size, which is almost always better!

# When the variances are the same

If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma^2)$ ,  $Y_1, Y_2, \dots, Y_n$  i.i.d.  $\sim N(\mu_Y, \sigma^2)$

- If it is known that both populations have the same variance, then we can leverage this information to get a more accurate estimate of by combining both  $s_X$  and  $s_Y$  to create what is called a pooled estimate of the variance:

$$s_{pool}^2 = \frac{(n-1)s_X^2 + (m-1)s_Y^2}{n+m-2},$$

where  $s_X^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$ ,  $s_Y^2 = \frac{(Y_1 - \bar{Y})^2 + \dots + (Y_m - \bar{Y})^2}{m-1}$ .

- The pooled estimate is a weighted average of the individual sample estimates. This increases the sample size, which is almost always better!
- The formula for confidence intervals now becomes:  $\bar{X} - \bar{Y} \pm t_{\alpha/2, k} \sqrt{\frac{s_p^2}{n} + \frac{s_p^2}{m}}$ .  
where  $k = n + m - 2$ .

# Practice problem

We are interested in investigating any potential difference between the mean blood sugar level of diabetics ( $\mu_X$ ) and that of non-diabetics ( $\mu_Y$ ). To do this we took a sample of six diabetics and found the following blood sugar levels: 127, 144, 140, 136, 118, 138. We also took a sample of eight non-diabetics and found the following blood sugar levels: 125, 128, 133, 141, 109, 125, 126, 122. **Assuming the variance for these two populations are the same.** (a) Estimate  $\mu_X - \mu_Y$ . (b) Find the 95% CI for  $\mu_X - \mu_Y$ .

( $s_X^2 = 93.24$ ,  $s_Y^2 = 83.55$ ,  $t_{0.05,12} = 1.782$ ,  $t_{0.025,12} = 2.179$ ,  $t_{0.05,14} = 1.761$ ,  $t_{0.025,14} = 2.145$ )

## Practice problem

We are interested in investigating any potential difference between the mean blood sugar level of diabetics ( $\mu_X$ ) and that of non-diabetics ( $\mu_Y$ ). To do this we took a sample of six diabetics and found the following blood sugar levels: 127, 144, 140, 136, 118, 138. We also took a sample of eight non-diabetics and found the following blood sugar levels: 125, 128, 133, 141, 109, 125, 126, 122. **Assuming the variance for these two populations are the same.** (a) Estimate  $\mu_X - \mu_Y$ . (b) Find the 95% CI for  $\mu_X - \mu_Y$ .

( $s_X^2 = 93.24$ ,  $s_Y^2 = 83.55$ ,  $t_{0.05,12} = 1.782$ ,  $t_{0.025,12} = 2.179$ ,  $t_{0.05,14} = 1.761$ ,  $t_{0.025,14} = 2.145$ )

### Solution

$$\hat{\mu}_X = \bar{X} = \frac{1}{6}(127 + 144 + 140 + 136 + 118 + 138) = 133.83.$$

$$\hat{\mu}_Y = \bar{Y} = \frac{1}{8}(125 + 128 + 133 + 141 + 109 + 125 + 126 + 122) = 126.13.$$

$$s_{pool}^2 = \frac{1}{6+8-2}((6-1) \times 93.24 + (8-1) \times 83.55) = 87.59$$

# Practice problem

We are interested in investigating any potential difference between the mean blood sugar level of diabetics ( $\mu_X$ ) and that of non-diabetics ( $\mu_Y$ ). To do this we took a sample of six diabetics and found the following blood sugar levels: 127, 144, 140, 136, 118, 138. We also took a sample of eight non-diabetics and found the following blood sugar levels: 125, 128, 133, 141, 109, 125, 126, 122. **Assuming the variance for these two populations are the same.** (a) Estimate  $\mu_X - \mu_Y$ . (b) Find the 95% CI for  $\mu_X - \mu_Y$ .

$$(s_X^2 = 93.24, s_Y^2 = 83.55, t_{0.05,12} = 1.782, t_{0.025,12} = 2.179, t_{0.05,14} = 1.761, t_{0.025,14} = 2.145)$$

## Solution

$$\hat{\mu}_X = \bar{X} = \frac{1}{6}(127 + 144 + 140 + 136 + 118 + 138) = 133.83.$$

$$\hat{\mu}_Y = \bar{Y} = \frac{1}{8}(125 + 128 + 133 + 141 + 109 + 125 + 126 + 122) = 126.13.$$

$$s_{pool}^2 = \frac{1}{6+8-2}((6-1) \times 93.24 + (8-1) \times 83.55) = 87.59$$

The 95% confidence interval is given as

$$\bar{X} - \bar{Y} \pm t_{0.025,12} \sqrt{s_{pool}^2} = (133.83 - 126.13) \pm 2.179 \sqrt{87.59} = [-6.45, 21.85]$$

# Paired Sample

- Sometimes it appears we have data from two samples with the further feature that there is a natural “pairing” of the data between the two samples.

# Paired Sample

- Sometimes it appears we have data from two samples with the further feature that there is a natural “pairing” of the data between the two samples.
- For example, suppose that the data consists on  $n$  brother-sister pairs, with blood pressures  $X_1, \dots, X_n$  for the  $n$  sisters and blood pressures  $Y_1, \dots, Y_n$  for their respective brothers.



# Paired Sample

- Sometimes it appears we have data from two samples with the further feature that there is a natural “pairing” of the data between the two samples.
- For example, suppose that the data consists on  $n$  brother-sister pairs, with blood pressures  $X_1, \dots, X_n$  for the  $n$  sisters and blood pressures  $Y_1, \dots, Y_n$  for their respective brothers.
- Thus  $X_1$  and  $Y_1$  are the blood pressures from the the sister and brother in family 1,  $X_2$  and  $Y_2$  are the blood pressures from the the sister and brother in family 2, and so on. The natural pairing is between the sister and the brother in the same family.

# Paired Sample

- Sometimes it appears we have data from two samples with the further feature that there is a natural “pairing” of the data between the two samples.
- For example, suppose that the data consists on  $n$  brother-sister pairs, with blood pressures  $X_1, \dots, X_n$  for the  $n$  sisters and blood pressures  $Y_1, \dots, Y_n$  for their respective brothers.
- Thus  $X_1$  and  $Y_1$  are the blood pressures from the the sister and brother in family 1,  $X_2$  and  $Y_2$  are the blood pressures from the the sister and brother in family 2, and so on. The natural pairing is between the sister and the brother in the same family.
- We could not use the previous CI formula for  $\mu_X - \mu_Y$ , because  $\{X_i\}$  and  $\{Y_i\}$  are not independent.

# CI for the difference between two means

Let's first consider the case where  $n$  is small.

- If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ . In the paired samples' case, it's not realistic to assume that  $\{X_i\}$  and  $\{Y_i\}$  are independent.

# CI for the difference between two means

Let's first consider the case where  $n$  is small.

- If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ . In the paired samples' case, it's not realistic to assume that  $\{X_i\}$  and  $\{Y_i\}$  are independent.
- However, it's natural to assume  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$  are  $n$  independent pairs.

# CI for the difference between two means

Let's first consider the case where  $n$  is small.

- If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ . In the paired samples' case, it's not realistic to assume that  $\{X_i\}$  and  $\{Y_i\}$  are independent.
- However, it's natural to assume  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$  are  $n$  independent pairs.
- We denote  $D_i = X_i - Y_i$ , then  $D_i$  i.i.d.  $\sim N(\mu_X - \mu_Y, \sigma_D^2)$

# CI for the difference between two means

Let's first consider the case where  $n$  is small.

- If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ . In the paired samples' case, it's not realistic to assume that  $\{X_i\}$  and  $\{Y_i\}$  are independent.
- However, it's natural to assume  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$  are  $n$  independent pairs.
- We denote  $D_i = X_i - Y_i$ , then  $D_i$  i.i.d.  $\sim N(\mu_X - \mu_Y, \sigma_D^2)$
- We estimate  $\mu_X - \mu_Y$  by  $\bar{D} = \bar{X} - \bar{Y}$

# CI for the difference between two means

Let's first consider the case where  $n$  is small.

- If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ . In the paired samples' case, it's not realistic to assume that  $\{X_i\}$  and  $\{Y_i\}$  are independent.
- However, it's natural to assume  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$  are  $n$  independent pairs.
- We denote  $D_i = X_i - Y_i$ , then  $D_i$  i.i.d.  $\sim N(\mu_X - \mu_Y, \sigma_D^2)$
- We estimate  $\mu_X - \mu_Y$  by  $\bar{D} = \bar{X} - \bar{Y}$
- A  $(1 - \alpha)$  confidence interval for  $\mu_X - \mu_Y$  is

$$\left[ \bar{D} - t_{\alpha/2, n-1} \frac{s_D}{\sqrt{n}}, \quad \bar{D} + t_{\alpha/2, n-1} \frac{s_D}{\sqrt{n}} \right],$$

$$\text{where } s_D = \sqrt{\frac{(D_1 - \bar{D})^2 + \dots + (D_n - \bar{D})^2}{n}}.$$

# CI for the difference between two means

Let's first consider the case where  $n$  is small.

- If  $X_1, X_2, \dots, X_n$  i.i.d.  $\sim N(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n$  i.i.d.  $\sim N(\mu_Y, \sigma_Y^2)$ . In the paired samples' case, it's not realistic to assume that  $\{X_i\}$  and  $\{Y_i\}$  are independent.
- However, it's natural to assume  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$  are  $n$  independent pairs.
- We denote  $D_i = X_i - Y_i$ , then  $D_i$  i.i.d.  $\sim N(\mu_X - \mu_Y, \sigma_D^2)$
- We estimate  $\mu_X - \mu_Y$  by  $\bar{D} = \bar{X} - \bar{Y}$
- A  $(1 - \alpha)$  confidence interval for  $\mu_X - \mu_Y$  is

$$\left[ \bar{D} - t_{\alpha/2, n-1} \frac{s_D}{\sqrt{n}}, \quad \bar{D} + t_{\alpha/2, n-1} \frac{s_D}{\sqrt{n}} \right],$$

where  $s_D = \sqrt{\frac{(D_1 - \bar{D})^2 + \dots + (D_n - \bar{D})^2}{n}}$ .

- If  $n$  is large, we can simply replace  $t_{\alpha/2, n-1}$  with  $z_{\alpha/2}$  and remove the normality assumptions.



# Example

Suppose that we take the blood pressures of  $n = 12$  women and their brothers, and get the following blood pressure reading:

Family	1	2	3	4	5	6	7	8	9	10	11	12
Sister	107	134	111	141	121	118	145	110	164	126	148	132
Brother	110	136	115	140	124	119	148	113	168	129	148	137

Construct the 95% CI for the difference of the mean blood pressure of men and women.

( $s_{sister}^2 = 307$ ,  $s_{brother}^2 = 299$ ,  $s_{diff}^2 = 3$ ,  $t_{0.025,11} = 2.306$ ,  $t_{0.05,11} = 1.860$ ,  $t_{0.025,12} = 2.262$ ,  $t_{0.05,12} = 1.833$ )

## Example

Suppose that we take the blood pressures of  $n = 12$  women and their brothers, and get the following blood pressure reading:

Family	1	2	3	4	5	6	7	8	9	10	11	12
Sister	107	134	111	141	121	118	145	110	164	126	148	132
Brother	110	136	115	140	124	119	148	113	168	129	148	137

Construct the 95% CI for the difference of the mean blood pressure of men and women.

( $s_{sister}^2 = 307$ ,  $s_{brother}^2 = 299$ ,  $s_{diff}^2 = 3$ ,  $t_{0.025,11} = 2.306$ ,  $t_{0.05,11} = 1.860$ ,  $t_{0.025,12} = 2.262$ ,  $t_{0.05,12} = 1.833$ )

## Answer

Family	1	2	3	4	5	6	7	8	9	10	11	12
Difference	3	2	4	-1	3	1	3	3	4	3	0	5

•  $\bar{D} = 2.5$ ,  $s_D^2 = 3$ ,  $t_{0.025,11} = 2.306$

## Example

Suppose that we take the blood pressures of  $n = 12$  women and their brothers, and get the following blood pressure reading:

Family	1	2	3	4	5	6	7	8	9	10	11	12
Sister	107	134	111	141	121	118	145	110	164	126	148	132
Brother	110	136	115	140	124	119	148	113	168	129	148	137

Construct the 95% CI for the difference of the mean blood pressure of men and women.

( $s_{sister}^2 = 307$ ,  $s_{brother}^2 = 299$ ,  $s_{diff}^2 = 3$ ,  $t_{0.025,11} = 2.306$ ,  $t_{0.05,11} = 1.860$ ,  $t_{0.025,12} = 2.262$ ,  $t_{0.05,12} = 1.833$ )

## Answer

Family	1	2	3	4	5	6	7	8	9	10	11	12
Difference	3	2	4	-1	3	1	3	3	4	3	0	5

- $\bar{D} = 2.5$ ,  $s_D^2 = 3$ ,  $t_{0.025,11} = 2.306$
- $2.5 \pm 2.306 \cdot \frac{\sqrt{3}}{\sqrt{12}} = [1.35, 3.65]$ .

# Summary

- CI for the binomial parameter  $p$  when the sample size  $n$  is large
- CI for the mean  $\mu$  when the sample size  $n$  is small and **samples are normal**
- CI for the difference between two binomial parameters  $p_1 - p_2$
- CI for the difference between two means  $\mu_X - \mu_Y$ 
  - Large sample size
  - Small sample size
  - When we have additional information that the variances are the same
  - Paired samples

# Homework

8.56, 8.60, 8.66, 8.70, 8.82, 8.85

# Other confidence intervals

- A useful method for deriving confidence intervals is to use a **pivotal quantity**
- A pivotal quantity
  - is a function of the sample data, the unknown target parameter, and **no other quantities**
  - has a distribution that does not depend on the target parameter
- Example:
- We randomly sample an observation from an exponential distribution with unknown mean  $\theta$ . Find a formula for a 90% CI for  $\theta$ .
- If  $Y \sim \text{Exp}(\theta)$ , then  $f_Y(y) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .
- Let  $U = \frac{Y}{\theta}$ , we have  $f_U(u) = f_Y(u\theta) \cdot \frac{dy}{du} = \frac{1}{\theta} e^{-u} \cdot \theta = e^{-u}$  for  $u > 0$ , that is,  $U \sim \text{Exp}(1)$
- Thus, we can use  $U = \frac{Y}{\theta}$  as a pivotal quantity


# Example

- We need to find two numbers  $a$  and  $b$  such that

$$\mathbb{P}(a \leq U \leq b) = 90\%$$

- One way to do this is to choose  $a$  and  $b$  to satisfy

$$\mathbb{P}(U < a) = \mathbb{P}(U > b) = 5\%$$



- $1 - e^{-a} = .05$  and  $e^{-b} = .05$ , equivalently,  $a = 0.051$ ,  $b = 2.996$
- $0.9 = \mathbb{P}(0.051 \leq \frac{Y}{\theta} \leq 2.996) = \mathbb{P}(\frac{Y}{2.996} \leq \theta \leq \frac{Y}{0.051})$

# Example

## Example

We randomly sample 10 observations  $X_1, \dots, X_{10}$  from an exponential distribution with unknown mean  $\theta$ . Find a formula for a 90% CI for  $\theta$ .

## Answer

- Hint: If  $X_i \sim \text{Exp}(\theta_i)$  then  $\min\{X_1, \dots, X_n\} \sim \text{Exp}(\frac{1}{\frac{1}{\theta_1} + \dots + \frac{1}{\theta_n}})$ .
- $\frac{X_1}{\theta}, \dots, \frac{X_{10}}{\theta} \stackrel{i.i.d.}{\sim} \text{Exp}(1)$
- $U = \min\{\frac{X_1}{\theta}, \dots, \frac{X_{10}}{\theta}\} \sim \text{Exp}(\frac{1}{10})$
- Find  $a, b$  such that  $\mathbb{P}(U < a) = 1 - e^{-10a} = .05$  and  $\mathbb{P}(U > b) = e^{-10b} = .05$ , equivalently,  $a = 0.005$ ,  $b = 0.300$
- $0.9 = \mathbb{P}(0.005 < U < 0.300) = \mathbb{P}(0.005 < \frac{\min\{X_1, \dots, X_{10}\}}{\theta} < 0.300) = \mathbb{P}(\frac{\min\{X_1, \dots, X_{10}\}}{0.300} < \theta < \frac{\min\{X_1, \dots, X_{10}\}}{0.005})$



# Example

## Example

Suppose that we take a sample  $Y$  from a uniform distribution defined on the interval  $[0, \theta]$ , where  $\theta$  is unknown. Find a 95% lower confidence bound for  $\theta$ , that is, find  $L(Y)$ , such that  $\mathbb{P}(\theta \geq L(Y)) = 0.95$ .

## Answer

- Consider  $U = \frac{Y}{\theta}$
- $U \sim U(0, 1)$
- $\mathbb{P}(U < 0.95) = 0.95$
- $0.95 = \mathbb{P}(U < 0.95) = \mathbb{P}\left(\frac{Y}{\theta} < 0.95\right) = \mathbb{P}\left(\frac{Y}{0.95} \leq \theta\right)$

# Confidence interval for the variance

- Suppose  $Y_1, \dots, Y_n \stackrel{i.i.d.}{\sim} N(\mu, \sigma^2)$  with  $\mu$  and  $\sigma^2$  unknown. We seek a  $100(1 - \alpha)\%$  CI for  $\sigma^2$
- Recall that, for  $S^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2$ , we have

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2.$$

- As a result,

$$\mathbb{P}(\chi_{1-(\alpha/2), n-1}^2 \leq \frac{(n-1)S^2}{\sigma^2} \leq \chi_{\alpha/2, n-1}^2) = 1 - \alpha$$

- A  $100(1 - \alpha)\%$  Confidence Interval for  $\sigma^2$ :

$$\mathbb{P}\left(\frac{(n-1)S^2}{\chi_{\alpha/2, n-1}^2} \leq \sigma^2 \leq \frac{(n-1)S^2}{\chi_{1-(\alpha/2), n-1}^2}\right) = 1 - \alpha.$$

# Example

## Example

Suppose the maturation times for a flower species are  $N(\mu, \sigma^2)$ . If a random sample of  $n = 13$  seeds yielded  $s^2 = 10.7$ , then what is a 90% CI for  $\sigma^2$ ?

( $\chi_{0.05,12}^2 = 21.03$ ,  $\chi_{0.95,12}^2 = 5.23$ )

## Answer

$$\mathbb{P}\left(\frac{(n-1)S^2}{\chi_{\alpha/2,n-1}^2} \leq \sigma^2 \leq \frac{(n-1)S^2}{\chi_{1-(\alpha/2),n-1}^2}\right) = 1 - \alpha.$$

- $\frac{(n-1)S^2}{\chi_{\alpha/2,n-1}^2} = \frac{12 \times 10.7}{21.03} = 6.11$
- $\frac{(n-1)S^2}{\chi_{1-\alpha/2,n-1}^2} = \frac{12 \times 10.7}{5.23} = 24.55$
- The 90% CI is  $[6.11, 24.55]$ .

# Homework

