

MSA HW2

Problem 1

a.

$$\|\mathbf{w}\|_1 = |2| + |3| + |-1| + |0| = 6$$

b.

$$\|\mathbf{w}\|_2 = \sqrt{2^2 + 3^2 + (-1)^2 + 0^2} = \sqrt{14}$$

c.

$$\|\mathbf{w}\|_\infty = \max\{|2|, |3|, |-1|, |0|\} = 3$$

d.

$$\text{supp}(\mathbf{w}) = \{1, 2, 3\}$$

e.

$$\text{card}(\text{supp}(\mathbf{w})) = 3$$

Problem 2

According to [FR2013], a non-negative function is called a norm if

(a) $\|\mathbf{x}\| = 0$ if and only if $\mathbf{x} = 0$

(b) $\|\lambda\mathbf{x}\| = |\lambda| \times \|\mathbf{x}\|$ for all scalar λ and \mathbf{x}

(c) $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$

However, (b) does not hold for all pairs of \mathbf{x} and \mathbf{y} . For instance, let $\lambda = 2$ and $\mathbf{y} = [3]$, then

$$\|\lambda\mathbf{y}\| = \|[6]\| = 1 \neq 2 = |2| \times \|[3]\|$$

Problem 3

a.

The Young's Inequality states that

$$\gamma\delta \leq \frac{\gamma^p}{p} + \frac{\delta^q}{q}$$

Let $\gamma_i = \frac{|u_i^*|}{\|\mathbf{u}\|_p}$ and $\delta_i = \frac{|v_i|}{\|\mathbf{v}\|_q}$, we have

$$\frac{|u_i^*|}{\|\mathbf{u}\|_p} \frac{|v_i|}{\|\mathbf{v}\|_q} \leq \frac{1}{p} \left(\frac{|u_i^*|}{\|\mathbf{u}\|_p} \right)^p + \frac{1}{q} \left(\frac{|v_i|}{\|\mathbf{v}\|_q} \right)^q$$

summing by i for both sides, we get

$$\sum_i \frac{|u_i^*| |v_i|}{\|\mathbf{u}\|_p \|\mathbf{v}\|_q} \leq \frac{1}{p} \sum_i \left(\frac{|u_i^*|}{\|\mathbf{u}\|_p} \right)^p + \frac{1}{q} \sum_i \left(\frac{|v_i|}{\|\mathbf{v}\|_q} \right)^q = \frac{1}{p} + \frac{1}{q} = 1$$

$$\rightarrow \sum_i |u_i^* v_i| \leq \|\mathbf{u}\|_p \|\mathbf{v}\|_q - (1)$$

Also, by the triangular inequality, $\sum_i |u_i^* v_i| \geq |\sum_i u_i^* v_i| = |\mathbf{u}^H \mathbf{v}| - (2)$

The inequality $|\mathbf{u}^H \mathbf{v}| \leq \|\mathbf{u}\|_p \|\mathbf{v}\|_q$ then holds due to (1) and (2)

b.

$$|\mathbf{u}^H \mathbf{v}| = |\sum_i u_i^* v_i| = |\sum_i \frac{|v_i|^q}{v_i \|\mathbf{v}\|_q^{q-1}} v_i| = |\sum_i \frac{|v_i|^q}{\|\mathbf{v}\|_q^{q-1}}| = |\frac{\sum_i |v_i|^q}{\|\mathbf{v}\|_q^{q-1}}| = \|\mathbf{v}\|_q$$

Also,

$$\begin{aligned} \|\mathbf{u}\|_p &= \sqrt[p]{\sum_i |u_i|^p} = \sqrt[p]{\sum_i \left[\frac{|v_i|^q}{v_i \|\mathbf{v}\|_q^{q-1}} \right]^p} = \sqrt[p]{\sum_i \frac{|v_i|^{pq}}{|v_i|^p \|\mathbf{v}\|_q^{pq-p}}} = \sqrt[p]{\sum_i \frac{|v_i|^{p+q}}{|v_i|^p \|\mathbf{v}\|_q^q}} = \\ &= \sqrt[p]{\sum_i \frac{|v_i|^q}{\|\mathbf{v}\|_q^q}} = \sqrt[p]{\frac{\|\mathbf{v}\|_q^q}{\|\mathbf{v}\|_q^q}} = 1 \end{aligned}$$

Therefore,

$$|\mathbf{u}^H \mathbf{v}| = \|\mathbf{u}\|_p \|\mathbf{v}\|_q \text{ when the condition holds.}$$

Problem 4

a.

Assume that $\mathbf{A} = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_6]$

Since $\mathbf{a}_i \neq c\mathbf{a}_j$ for all pairs of i, j and scaling factor c , the spark is not 2.

Since $\mathbf{a}_3 - 2\mathbf{a}_5 - \mathbf{a}_6 = \mathbf{0}$, $\mathbf{a}_3, \mathbf{a}_5, \mathbf{a}_6$ is linearly dependent.

Therefore, the spark of the matrix is 3.

b.

The kruskal rank is therefore 2 due to **a.**

The linearly dependent sets are:

$$\mathcal{S} = \{\{\mathbf{a}_i, \mathbf{a}_j\} | i \neq j\}$$

Problem 5

For $\|\mathbf{z}\|_0 = 0$, it is obviously impossible.

For $\|\mathbf{z}\|_0 = 1$, it is clear that no column vectors in \mathbf{A} is a scaled from \mathbf{y} .

For $\|\mathbf{z}\|_0 = 2$, we check if all pairs of column vectors in \mathbf{A} are linear independent with \mathbf{y} .

MATLAB code:

```

A = [-2 -4 4 9 8; ...
-7 9 4 -4 5; ...
5 -3 0 6 3];
z = [-22; 17; -15];
for i = 1 : 4
    for j = i+1 : 5
        if rank([A(:,i) A(:,j) z]) < 3
            fprintf("a_%d & a_%d\n",i,j)
        end
    end
end
end
end

```


The results find that only $\mathbf{a}_2, \mathbf{a}_4, \mathbf{y}$ are linearly dependent and this forms a unique solution.

Therefore, the solution set is:

$$\mathcal{S}_z = \{[0, 1, 0, -2, 0]^T\}$$


Problem 6

a.



$$\begin{aligned}
 \text{Var}(f) &: [9.38 \times 10^{-6}, 2.06 \times 10^{-6}, 6.87 \times 10^{-6}, 3.14 \times 10^{-6}, 2.35 \times 10^{-6}, 3.57 \times 10^{-5}] \\
 \text{Mean}(f) &: [-2.50 \times 10^{-1}, -1.50 \times 10^{-1}, 9.96 \times 10^{-3}, 1.00 \times 10^{-1}, 2.00 \times 10^{-1}, 3.01 \times 10^{-1}] \\
 \text{MSE}(f) &: [9.40 \times 10^{-6}, 2.06 \times 10^{-6}, 6.87 \times 10^{-6}, 3.14 \times 10^{-6}, 2.35 \times 10^{-6}, 3.65 \times 10^{-5}]
 \end{aligned}$$

b.



$$\begin{aligned}
 \text{Var}(f) &: [9.38 \times 10^{-6}, 2.06 \times 10^{-6}, 6.87 \times 10^{-6}, 3.14 \times 10^{-6}, 2.36 \times 10^{-6}, 3.56 \times 10^{-5}] \\
 \text{Mean}(f) &: [-2.50 \times 10^{-1}, -1.50 \times 10^{-1}, 9.95 \times 10^{-3}, 1.00 \times 10^{-1}, 2.00 \times 10^{-1}, 3.01 \times 10^{-1}] \\
 \text{MSE}(f) &: [9.40 \times 10^{-6}, 2.06 \times 10^{-6}, 6.87 \times 10^{-6}, 3.14 \times 10^{-6}, 2.36 \times 10^{-6}, 3.64 \times 10^{-5}]
 \end{aligned}$$