


Lecture 2: Example applications

COMP5930M Scientific Computation

Today

1. Navigation by GPS

- ▶ Receiver calculates its position (x, y, z) based on signals sent by four GPS satellites and received at times t_i^R .
- ▶ Each satellite $i = 1, 2, 3, 4$ transmits messages that include the transmission time t_i^S and the satellite position (x_i, y_i, z_i) at the time of transmission.
- ▶ The clock error, $\Delta t = t_{\text{satellite}} - t_{\text{receiver}}$, is also unknown.



1. Navigation by GPS

- ▶ Receiver calculates its position (x, y, z) based on signals sent by four GPS satellites and received at times t_i^R .
- ▶ Each satellite $i = 1, 2, 3, 4$ transmits messages that include the transmission time t_i^S and the satellite position (x_i, y_i, z_i) at the time of transmission.
- ▶ True range from satellite i :

$$d_i = \sqrt{(x - x_i)^2 + (y - y_i)^2 + (z - z_i)^2}$$

Pseudorange from satellite i :

$$p_i = c \left(t_i^R - t_i^S \right)$$

where c is the speed of light (assuming no clock error)

1. Navigation by GPS

- ▶ Receiver calculates its position (x, y, z) based on signals sent by four GPS satellites and received at times t_i^R .
- ▶ Each satellite $i = 1, 2, 3, 4$ transmits messages that include the transmission time t_i^S and the satellite position (x_i, y_i, z_i) at the time of transmission.
- ▶ True range from satellite i :

$$d_i = \sqrt{(x - x_i)^2 + (y - y_i)^2 + (z - z_i)^2}$$

Pseudorange from satellite i :

$$p_i = c \left(t_i^R - t_i^S \right)$$

where c is the speed of light (assuming no clock error)

1. Navigation by GPS

- ▶ True range from satellite i :

$$d_i = \sqrt{(x - x_i)^2 + (y - y_i)^2 + (z - z_i)^2}$$

Pseudorange from satellite i :

$$p_i = c \left(t_i^R - t_i^S \right)$$

where c is the speed of light (assuming no clock error)

- ▶ By incorporating the clock error, we can relate the true range with the pseudorange:

$$d_i = c \left(t_i^R - t_i^S + \Delta t \right)$$

1. Navigation by GPS

- ▶ True range from satellite i :

$$d_i = \sqrt{(x - x_i)^2 + (y - y_i)^2 + (z - z_i)^2}$$

Pseudorange from satellite i :

$$p_i = c \left(t_i^R - t_i^S \right)$$

where c is the speed of light (assuming no clock error)

- ▶ By incorporating the clock error, we can relate the true range with the pseudorange:

$$d_i = c \left(t_i^R - t_i^S + \Delta t \right)$$

Navigation equations (four satellites)

By collecting all four equations into one system, we can in theory solve for the four unknowns $(x, y, z, \Delta t)$:

$$\begin{cases} \sqrt{(x - x_1)^2 + (y - y_1)^2 + (z - z_1)^2} - c(t_1^R - t_1^S + \Delta t) = 0 \\ \sqrt{(x - x_2)^2 + (y - y_2)^2 + (z - z_2)^2} - c(t_2^R - t_2^S + \Delta t) = 0 \\ \sqrt{(x - x_3)^2 + (y - y_3)^2 + (z - z_3)^2} - c(t_3^R - t_3^S + \Delta t) = 0 \\ \sqrt{(x - x_4)^2 + (y - y_4)^2 + (z - z_4)^2} - c(t_4^R - t_4^S + \Delta t) = 0 \end{cases}.$$

\Rightarrow nonlinear equation system $F(\vec{x}) = 0$

Possible issues in solving the navigation equations

- ▶ The equations may have multiple solutions or no solutions¹
⇒ Are the solutions **unique** (up to some conditions)?
- ▶ Receiver motion introduces uncertainty to the times t_i^R
⇒ How **sensitive** are the solutions to the coefficients?

¹Abel J and Chauffe J. Existence and uniqueness of GPS solutions. IEEE Transactions on Aerospace and Electronic Systems 27:952956, 1991.

Possible issues in solving the navigation equations

- ▶ The equations may have multiple solutions or no solutions¹
⇒ Are the solutions **unique** (up to some conditions)?
- ▶ Receiver motion introduces uncertainty to the times t_i^R
⇒ How **sensitive** are the solutions to the coefficients?
- ▶ Nonlinear equation solution algorithms often iterative
⇒ How to generate **initial solutions** for the location?

¹Abel J and Chauffe J. Existence and uniqueness of GPS solutions. IEEE Transactions on Aerospace and Electronic Systems 27:952956, 1991.

Possible issues in solving the navigation equations

- ▶ The equations may have multiple solutions or no solutions¹
⇒ Are the solutions **unique** (up to some conditions)?
- ▶ Receiver motion introduces uncertainty to the times t_i^R
⇒ How **sensitive** are the solutions to the coefficients?
- ▶ Nonlinear equation solution algorithms often iterative
⇒ How to generate **initial solutions** for the location?

¹Abel J and Chauffe J. Existence and uniqueness of GPS solutions. IEEE Transactions on Aerospace and Electronic Systems 27:952956, 1991.

2. Simple differential equation

Predator-prey -model in computational ecology:

"Assume we have an ecological system with two species: prey and predator. The prey reproduce without limits due to large quantities of food. The predator needs to eat the prey to reproduce."

Objective: describe the dynamics of the two populations

- ▶ t is our time variable
- ▶ $x(t)$ is the number of animals in the prey species
- ▶ $y(t)$ is the number of animals in the predator species
- ▶ The initial populations are $x(0) = x_0$ and $y(0) = y_0$
- ▶ The change in time of the populations are $\frac{dx}{dt}$ and $\frac{dy}{dt}$

2. Simple differential equation

Lotka-Volterra model:

$$\begin{aligned}\frac{dx}{dt} &= \alpha x(t) - \beta x(t)y(t) \\ \frac{dy}{dt} &= \delta x(t)y(t) - \gamma y(t)\end{aligned}\tag{1}$$

where $\alpha, \beta, \gamma, \delta$ are positive constants

2. Simple differential equation

Interpretation of terms in the equation:

- ▶ $\alpha x(t)$ is the reproduction rate of new prey species

Since $\frac{dx}{dt} = \alpha x(t)$ has solution $x(t) = C \exp(\alpha t)$, the prey population grows exponentially when there are no predators

2. Simple differential equation

Interpretation of terms in the equation:

- ▶ $\alpha x(t)$ is the reproduction rate of new prey species

Since $\frac{dx}{dt} = \alpha x(t)$ has solution $x(t) = C \exp(\alpha t)$, the prey population grows exponentially when there are no predators

- ▶ $-\beta x(t)y(t)$ is the predation rate, i.e. how many prey are being eaten by the predators. Note this is a non-linear term.

2. Simple differential equation

Interpretation of terms in the equation:

- ▶ $\alpha x(t)$ is the reproduction rate of new prey species

Since $\frac{dx}{dt} = \alpha x(t)$ has solution $x(t) = C \exp(\alpha t)$, the prey population grows exponentially when there are no predators

- ▶ $-\beta x(t)y(t)$ is the predation rate, i.e. how many prey are being eaten by the predators. Note this is a non-linear term.
- ▶ $\delta x(t)y(t)$ is the growth rate of predators, which depends on how much prey is available. Another non-linear term.

2. Simple differential equation

Interpretation of terms in the equation:

- ▶ $\alpha x(t)$ is the reproduction rate of new prey species

Since $\frac{dx}{dt} = \alpha x(t)$ has solution $x(t) = C \exp(\alpha t)$, the prey population grows exponentially when there are no predators

- ▶ $-\beta x(t)y(t)$ is the predation rate, i.e. how many prey are being eaten by the predators. Note this is a non-linear term.
- ▶ $\delta x(t)y(t)$ is the growth rate of predators, which depends on how much prey is available. Another non-linear term.
- ▶ $-\gamma y(t)$ is the the natural death rate of the predator species

Predators will die off exponentially if there is no prey

2. Simple differential equation

Interpretation of terms in the equation:


- ▶ $\alpha x(t)$ is the reproduction rate of new prey species

Since $\frac{dx}{dt} = \alpha x(t)$ has solution $x(t) = C \exp(\alpha t)$, the prey population grows exponentially when there are no predators

- ▶ $-\beta x(t)y(t)$ is the predation rate, i.e. how many prey are being eaten by the predators. Note this is a non-linear term.
- ▶ $\delta x(t)y(t)$ is the growth rate of predators, which depends on how much prey is available. Another non-linear term.
- ▶ $-\gamma y(t)$ is the the natural death rate of the predator species
Predators will die off exponentially if there is no prey

2. Simple differential equation

In many cases, the solution can be shown to approach a periodic orbit where the number of predators/prey behave cyclically:



Phase-space plot in the (x, y) space tracks the solution from a given initial solution (x_0, y_0)

Numerical approximation of differential equations

In practice, differential equations are often **discretised** in time on a numerical grid with discrete time step $\Delta t = t_j - t_{j-1}$:

$$X_j = x(t_j) \text{ for } t_0 < t_1 < \dots < t_N$$

by applying difference approximations for the derivatives, e.g.

$$\frac{\partial x}{\partial t}(t_j) \approx \frac{X_j - X_{j-1}}{\Delta t}.$$

This leads to a system $F(\vec{X}) = 0$ of algebraic nonlinear equations for the X_j to be solved at each time step $t_{j-1} \rightarrow t_j$.

Return to these techniques later in the module...

Numerical approximation of differential equations

In practice, differential equations are often **discretised** in time on a numerical grid with discrete time step $\Delta t = t_j - t_{j-1}$:

$$X_j = x(t_j) \text{ for } t_0 < t_1 < \dots < t_N$$

by applying difference approximations for the derivatives, e.g.

$$\frac{\partial x}{\partial t}(t_j) \approx \frac{X_j - X_{j-1}}{\Delta t}.$$

This leads to a system $F(\vec{X}) = 0$ of algebraic nonlinear equations for the X_j to be solved at each time step $t_{j-1} \rightarrow t_j$.

Return to these techniques later in the module...

2. Simple differential equation

Nonlinear equations for continuous Lotka-Volterra model:

$$\frac{dx}{dt} = \alpha x(t) - \beta x(t)y(t) \tag{2}$$

$$\frac{dy}{dt} = \delta x(t)y(t) - \gamma y(t)$$

2. Simple differential equation

Nonlinear equations for discrete Lotka-Volterra model:

$$\frac{X_j - X_{j-1}}{\Delta t} = \alpha X_j - \beta X_j Y_j \quad (3)$$

$$\frac{Y_j - Y_{j-1}}{\Delta t} = \delta X_j Y_j - \gamma Y_j$$

for $j = 1, 2, \dots, N$

Replace derivatives with discrete difference approximations

2. Simple differential equation

Nonlinear equations for discrete Lotka-Volterra model:

$$X_j - X_{j-1} = \Delta t [\alpha X_j - \beta X_j Y_j] \quad (4)$$

$$Y_j - Y_{j-1} = \Delta t [\delta X_j Y_j - \gamma Y_j]$$

for $j = 1, 2, \dots, N$

Multiply both sides by Δt

2. Simple differential equation

Nonlinear equations for discrete Lotka-Volterra model:

$$X_j - X_{j-1} - \Delta t [\alpha X_j - \beta X_j Y_j] = 0 \quad (5)$$

$$Y_j - Y_{j-1} - \Delta t [\delta X_j Y_j - \gamma Y_j] = 0$$

for $j = 1, 2, \dots, N$

Move all terms to the left-hand side

2. Simple differential equation

Nonlinear equations for discrete Lotka-Volterra model:

$$F_1(X_j, Y_j) = X_j - X_{j-1} - \Delta t [\alpha X_j - \beta X_j Y_j] = 0 \quad (6)$$

$$F_2(X_j, Y_j) = Y_j - Y_{j-1} - \Delta t [\delta X_j Y_j - \gamma Y_j] = 0$$

for $j = 1, 2, \dots, N$

Identify the non-linear equations

Numerical approximation of differential equations

- ▶ Depending on application, number of equations can be large
⇒ Do our algorithms **scale** for large problems?
- ▶ Does the approximate solution U approach u as $N \rightarrow \infty$?
⇒ What are the criteria for **convergence**?

Numerical approximation of differential equations

- ▶ Depending on application, number of equations can be large
⇒ Do our algorithms **scale** for large problems?
- ▶ Does the approximate solution U approach u as $N \rightarrow \infty$?
⇒ What are the criteria for **convergence**?
- ▶ Time-dependent problems require choice of time step Δt
⇒ How to choose time step Δt that is **stable**?

Numerical approximation of differential equations

- ▶ Depending on application, number of equations can be large
⇒ Do our algorithms **scale** for large problems?
- ▶ Does the approximate solution U approach u as $N \rightarrow \infty$?
⇒ What are the criteria for **convergence**?
- ▶ Time-dependent problems require choice of time step Δt
⇒ How to choose time step Δt that is **stable**?