


∞ -Spaces



ND

ND

ND

ND

1. Introduction

Implementation Progress

Writing Progress

1. In this text I would like to develop a functorial construction of the classifying space in homotopy which can be applied indefinitely.
2. This construction is an endofunctor of algebras for operadic groups in operadic groups in ∞ -Grpd.
- 3.

I would like to notate it as B^1 .

This notation distinguishes my attempt at an abelian-classifying-space construction which is an endofunctor of grouplike E^∞ -spaces, here referred to as ∞ -spaces, from the classifying-space construction involving ∞ -groupoids and grouplike A^∞ -spaces. Note that our ∞ -categories always have $r \equiv 1$, i.e. ∞ is short for $(\infty, 1)$.

I will write B^n for the n -fold composition of B^1 , $\text{Pow } B^n$. Note that B is not an endofunctor so that it cannot be iterated. It is more typical to divide up the construction B , for instance constructing the Grassmanians $\text{Gr}^\infty(\mathbb{C}, n) \cong B.\text{obj } \text{GL}_n(\mathbb{C})$, from which it follows that $[-, \text{Gr}^\infty(\mathbb{C}, n)]$ is equivalent to the category of $\text{GL}_n(\mathbb{C})$ -principal bundles and therefore to n -dimensional vector bundles.

In “TheWhiteheadTheoremandTwoVariations”, we developed two models of ∞ -Grpd_(A) and ∞ -Grpd_(B). These will produce two models on which B^1 can be defined:

$$B^1 : \text{OperadicGroup} \bullet \text{OperadicGroup } \infty\text{-Grpd} \longrightarrow \text{OperadicGroup} \bullet \text{OperadicGroup } \infty\text{-Grpd}$$

∞ -Spaces ...

In this section, which makes use of the previous section concerning Haar integral, I intend to cover the ordinary versions of Poincare duality, Pontrjagin duality, and Fourier duality, as well as versions of these theorems using language enabled by the previous repositories. This won't culminate until far into the future, so for now I have jotted down some sketches.

after this we develop chain complexes of these.

The table of contents below reflects the tentative long-term goals of the authors, with the main goal the pursuit of the Whitehead theorem for a point-set model involving Mathlib's predefined homotopy groups.

1. cup and cap product

Ideas for future applications:

1. <https://arxiv.org/pdf/2206.13563.pdf>

1. One of the basic things I wanted out of this was homotopy colimit preserving maps $(E^{\text{inf}} \text{-Alg } A)^{\text{op}} \rightarrow \infty\text{-Grpd}$

Note that this repository does not implement $Q X := \text{colimit}_n \Omega^n \Sigma^n X$ or the stable homotopy groups. It concerns the relationship between O and B and not Ω and Σ .

2. Contents

Section	Description
Unfinished	
Contents	
Unicode	
Introduction	
PART I: ∞ -SPACES	
Chapter 1: Abelian Groups	
abeliangroup	The type of abelian groups
Maps of abelian groups	Constructing homomorphisms of abelian groups
Negation	
The Eckman-Hilton Argument	
AbelianGroup \longrightarrow Group	The forgetful functor from abelian groups to groups
Eilenberg-MacLane Spaces	
Chain Complexes	
Realization of Chain Complexes	
Tensor Product of Chain Complexes	
Chapter 2: ∞ -Spaces	
∞ -space	The type of ∞ -spaces
Maps of ∞ -spaces	Constructing maps of ∞ -spaces
Negation	
The Eckman-Hilton Argument	
OperadicGroup OperadicGroup ∞ -Grpd ₋₁ \longrightarrow OperadicGroup ∞ -Grpd ₋₁	
B^1 and $B^{\mathbb{N}}$	
Chn : ???	
Realization of Chain Complexes	
Tensor Product of Chain Complexes	
Chapter 3: Tensor Product of Abelian Groups	
- \otimes (AbelianGroups) -	Mathlib's tensor product of abelian groups
[\cdot , \cdot] (AbelianGroups)	Mathlib's hom of abelian groups
AbelianGroup	The symmetric monoidal category of abelian groups
Chapter 4: Tensor Product of ∞ -Spaces	
- \otimes (∞ -Space) -	
[\cdot , \cdot] (∞ -Space)	
∞ -Space	The symmetric monoidal category of ∞ -spaces
Chapter 5: Set ₋₁ \rightleftarrows AbelianGroups	
???	The free abelian group functor
???	The forgetful functor from abelian groups to pointed sets
??? : Set ₋₁ \rightleftarrows AbelianGroup : ???	The adjunction between pointed sets and abelian groups
Chapter 6: ∞ -Grpd ₋₁ \rightleftarrows ∞ -Space	
???	The free ∞ -space given a based ∞ -groupoid
???	The forgetful functor from ∞ -spaces to ∞ -groupoids
??? : ∞ -Grpd ₋₁ \rightleftarrows ∞ -Space : ???	The ??? between ∞ -Grpd ₋₁ and ∞ -Spaces
PART II: RINGS, COMMUTATIVE RINGS, A^∞ -RINGS, AND E^∞ -RINGS	
Chapter 7: Rings and Commutative Rings	

ring	The type of rings
Ring	The category of rings
commutative_ring	The type of commutative rings
CommutativeRing	The category of commutative rings
Chapter 8: A^∞ -Rings and E^∞ -Rings	
A^∞ -ring	The type of A^∞ -rings
A^∞ -Ring	The category of A^∞ -Rings
E^∞ -ring	The type of E^∞ -rings
E^∞ -Ring	The category of E^∞ -Rings
Chapter 9: Modules and Modules over Commutative Rings	
InternalMonoidAction (InternalMonoid C) \cong InternalMonoid (InternalMonoidAction C)	The ??? theorem
CommutativeAlgebra : CommutativeRing \rightarrow Cat	The category of commutative algebras
Maps (Algebra A) : Cat	The category of maps of commutative A-algebras
Chapter 10: A^∞ -Modules and E^∞ -Modules	
A^∞ -RingAction (A^∞ -Ring C) \cong A^∞ -Ring (A^∞ -RingAction C)	The ??? theorem
Maps A^∞ -Algebras	
PART III: DERIVATIONS AND CONNECTIONS	
Chapter 11: Lie Algebras	
lie_algebra	The type of Lie-algebras
LieAlgebra	The category of Lie-algebras
Chapter 12: Derivations	
InternalAbelianGroup (Maps (Algebra A)) \cong MonoidActionObject A	???
??? : (Maps (Algebra A)) \rightleftarrows InternalAbelianGroup (Maps (Algebra A)) : ???	The free abelian group functor for (Maps (Algebra A))
Λ : ??? \rightleftarrows ??? : FstDeg	
??? : (Algebra A) \rightleftarrows Chn (Algebra A) : ???	The free DGA functor
derivation	Definition of a derivation
Der : () \rightleftarrows (InternalMonoidAction A) : ???	A derivation is a primitive element
Chapter 13: L^∞ Algebras	
L^{inf} _algebra	The type of L^∞ -algebras
L^∞ Algebra	
Chapter 14: ∞ -Derivations	
OperadicAbelianGroup (Maps (∞ -Algebra A)) \cong E^∞ MonoidAction A	???
??? : A^∞ -Algebras \rightleftarrows ???	The free abelian group
Λ : ??? \rightleftarrows ??? : FstDeg	
??? : (???) \rightleftarrows (???) : ???	The free ???
∞ -derivation	Definition of an ∞ -derivation
∞ -Der : () \rightleftarrows () : ???	A derivation is an ∞ -primitive element
Chapter 15: Tensor Product of Lie Algebras	
- \otimes () -	
LieAlgebra : ???	The monoidal category of Lie algebras
Chapter 16: Tensor Product of L^∞ -Algebras	
- \otimes () -	
: ???	The symmetric monoidal category of L^∞ -algebras
Chapter 17: Lie Algebra Representations	
Chapter 18: Connections	
???	The ??? equivalence
???	The free internal abelian group action functor
???	The first degree of the free E^∞ -DGM on an algebra is $s_()$

???	???
connection	Definition of a connection
???	A connection is a d-action
Chapter 19: L^∞ -Representations	
Chapter 20: ∞ -Connections	
???	The ??? equivalence
???	The free operadic abelian group action functor
???	The first degree of the free E^∞ -DGM on an algebra is s^{inf}_-
???	???
connection	Definition of a connection
???	A connection is a d^{inf} -action
Chapter 21: Tensor Product of Lie-Algebra Representations	
$- \otimes_0 -$	
$[-, -]_0$	
???	The symmetric monoidal closed category of Lie-algebra representations
Chapter 22: Tensor Product of L^∞ -Algebra Representations	
$- \otimes_0 -$	
$[-, -]_0$	
???	The symmetric monoidal closed category of L^∞ -algebra representations

PART 1: ABELIAN GROUPS AND ∞ -SPACES


In this first section I will construct eight structures for monoidal categories. These structures will be constructed so as to be endofunctions of a particular kind of monoidal category (as opposed to a cartesian category).

Eight Structures			
Strict		Lax	
Unital	Actional	Unital	Actional
InternalCommutativeMonoid	InternalCommutativeMonoidAction	OperadicCommutativeMonoid	OperadicCommutativeMonoidAction
InternalAbelianGroup	InternalAbelianGroupAction	OperadicAbelianGroup	OperadicAbelianGroupAction


One particular kind of monoidal category is a *cartesian category*, i.e. a monoidal category in which the monoidal operation is cartesian product. Fox's theorem says that a symmetric monoidal category is cartesian if and only if it is isomorphic to co-commutative comonoids in itself.

We will also be interested to form the tensor product of abelian groups and the tensor product of ∞ -spaces.

Abelian Groups



3. The Category of Commutative Monoids and the Category




4. `AbelianGroup` \longrightarrow `Group`

In this chapter I cover the type of abelian groups as well as the category of abelian groups.

1. `Mathlib.Algebra.Group.Defs`




5. Negation




6. The Eckman-Hilton Argument

The Eckman-Hilton Argument demonstrates that internal groups in the monoidal category of groups with product as monoidal operation is equivalent to the category of abelian groups.




7. Eilenberg-MacLane Spaces

Definition 1. Let A be an abelian group, and let $n \in \mathbb{N}$ be a non-negative integer. An Eilenberg-MacLane space .



8. Chain Complexes

Definition 2. ...




9. Realization of Chain Complexes

Definition 3. ...




10. Tensor Product of Chain Complexes


Definition 4. ...



∞ -Spaces




11. The category of ∞ -Spaces and the category of E^∞ -Spa



(x, y)



ND



12. ∞ -Spaces \longrightarrow OperadicGroup • OperadicGroup ∞ -Grpd.

An ∞ -space is an algebra for the little-squares operad.

The little-squares operad is OperadicGroup 2 ∞ -Grpd.

, which we have here taken to mean an operadic group in operadic groups in based connected ∞ -groupoids, is a particular operadic group. In this approach, we have considered OperadicGroup to have type $\mathbb{N} \longrightarrow \text{Cat}$ rather than $\infty\text{-Cat} \longrightarrow \infty\text{-Cat}$ or $\infty_-(\infty\text{-Cat}) \longrightarrow \infty_-(\infty\text{-Cat})$. Specifically, we can supply a non-negative integer to obtain a particular operad resembling little n-cubes but which features no "empty space".


13. Negation


For an ∞ -space A , negation $\neg : A \rightarrow A \dots$


∞ -spaces




14. The Eckman-Hilton Argument











OperadicGroup 2 ∞ -Grpd₁




15. B^1 and B^n

Definition 5. B^1


Definition 6. B^n is $B^1 \bullet B^{n-1}$.




16. Chain Complexes



17. Realization of Chain Complexes




18. Tensor Product of Chain Complexes




Tensor Product

1.



Tensor Product of ∞ -Spaces

$\infty\text{-Grpd}_1$ with smash product forms an operadic commutative monoid, but not a monoidal category.



$$\mathbf{Set}_{-1} \rightleftarrows \mathbf{AbelianGroup}$$

Construction	Description
$??? : \mathbf{InternalAbelianGroup} \mathbf{Set}_{-1} \cong \mathbf{AbelianGroup} : ???$	The $???$ theorem
$??? : \mathbf{Set}_{-1} \rightleftarrows \mathbf{AbelianGroup} : ???$	The $???$ adjunction

Abelian groups are internal groups in internal groups in sets.

In forming the free group on a set, based sets intermediate the construction.

$$\infty\text{-Grpd}_1 \rightleftarrows \infty\text{-Space}$$

In this section, we construct a

Construction	Description
$??? : \text{InternalAbelianGroup Set}_1 \cong \text{AbelianGroup} : ???$	The ??? theorem
$??? : \infty\text{-Grpd}_1 \rightleftarrows \infty\text{-Space} : ???$	The ??? adjunction

1. π_n of an ∞ -space arising from an ∞ -groupoid

2. H_n of an

3. Dold-Thom theorem

4.

5. B is iterable on this

6. $\text{Chn} \dots$

7. $\mu : \text{Chn} \times \text{Chn} \longrightarrow \text{Chn}$

$S_-(???)$ in general...

PART 2: RINGS, COMMUTATIVE RINGS, A^∞ -RINGS, AND E^∞ -RINGS

In this second section, I consider four constructions for monoidal categories which have occurred less generally elsewhere for cartesian categories: internal monoids, internal commutative monoids, algebras for A^∞ -operads, and algebras for E^∞ -operads. The main difference with the structures featured previously is that these structures concern an operation which is more general than product, but less general than pull-back. Four of the sixteen structures formed in the repository concerning the Whitehead theorem can re-create the rest, are not instances of the structures here. Meanwhile, the four structures defined here will be constructed using Mathlib 4's monoidal categories and symmetric monoidal categories. Tensor product of abelian groups and smash product of ∞ -spaces, coincide with the previous structures in the case where the monoidal operation is product (see Fox's theorem).

To reflect the use of monoidal categories as opposed to categories (in which the cartesian monoidal structure can be recovered from the structure as a seven entry with the addition of a single Lean universe), I use different names for the constructions:

Categories of Internal Objects			
Strict		Lax	
Unital	Actional	Unital	Actional
MonoidObjects : ??? \rightarrow ???	MonoidActionObjects : ??? \rightarrow ???	A^∞ -Monoid ∞ -Space	A^∞ -Monoid
CommutativeMonoidObjects : ??? \rightarrow ???	CommutativeMonoidActionObjects : ??? \rightarrow ???	E^∞ -Monoid ∞ -Space	E^∞ -Monoid

Rings and Commutative Rings

Rings and DGAs

Commutative rings and CDGAs


1. A thread on creating the six-entry category of commutative algebras.
2. In this section we use slightly different internal structures than the internal monoid in the last section; these internal monoids are defined in a monoidal category and the others are defined for product only. As such we may like to re-examine those structures, or alternatively to keep separate definitions.
3. What's more clear is that the ∞ -analogues are more difficult to reconcile with the choices made for the first sixteen structures and the four doubled structures (see the section on the Eckman-Hilton argument)

A^∞ -Rings and E^∞ -Rings


make sure to include Alg
make sure to include ∞ -Alg...

1. What's more clear is that the ∞ -analogues are more difficult to reconcile with the choices made for the first sixteen structures and the four doubled structures (see the section on the Eckman-Hilton argument).

Modules over Rings and
Commutative Rings



A^∞ -Modules and E^∞ -Modules



PART 3: DERIVATIONS AND CONNECTIONS

In this section, I define four more structures:

Four Definitions		
	Strict	Lax
Unital	Derivation	∞ -Derivation
Actional	Connection	∞ -Connection

I also construct four adjunctions which feature the internal abelian group structure:

Four Free Constructions in Algebra		
	Strict	Lax
Unital	$??? : \text{Map } ??? \rightleftarrows \text{InternalAbelianGroup}(\text{Map } ???) : ???$	$??? : \text{Map } ??? \rightleftarrows (\text{OperadicAbelianGroup}(\text{Map } ???)) : ???$
Actional	$??? : ??? \rightleftarrows \text{InternalAbelianGroupAction } ??? : ???$	$??? : ??? \rightleftarrows \text{OperadicAbelianGroupAction } ??? : ???$

In this second part, I define eight adjunctions associated to the algebraic structures defined in the last section.

Lie Algebras

Definition 7 (Lie Algebra). Let A be a (commutative unital) ring. A Lie-algebra is an A -module M such that...

Definition 8 (Lie Algebra Map). Let A be a (commutative unital) ring.

1. A thread on the classification of Lie algebras.

Derivations

In a blog post here,

Let A be a ring and suppose that $B : \text{Alg } A \rightarrow \text{dom.}$

$$S_*(\text{Mon}(\text{Act } R)) : (\text{ComMon } R) \rightleftarrows S_*(\text{Mon}(\text{Act } R))$$


Theorem 1. the category of internal abelian groups in $\text{Mon}(\text{Act } R)$ is equivalent to $\text{Act } R$

$$S_*(\text{Mon}(\text{Ch}(\text{Act } R))) : \text{Mon}(\text{Ch}(\text{Act } R)) \rightleftarrows \text{Mod } R : S_*(\text{Mon}(\text{Ch}(\text{Act } R)))$$

$$(\text{Alg } A)/A \rightleftarrows (\text{Alg } A)$$

1. I would like to first construct the lie-algebra of derivations using the spectrum $\Omega^{\text{inf}}. \text{obj } X$. It seems related to coalgebra endomorphisms from $\Omega^{\text{inf}}. \text{obj } X$ to itself.
2. Lie algebras and $\text{Der}^? (A, A)$

L^∞ Algebras



∞ -Derivations

$$\Omega_{-}(). : (E^{\text{inf}}\text{-Alg } A)/A \rightleftarrows E^{\text{inf}}\text{-Mod } A : \Omega_{-}()$$

$$\Lambda_{-}(). : \text{Ch } (E^{\text{inf}}\text{-Alg } A) \rightleftarrows E^{\text{inf}}\text{-Mod } A : \Lambda_{-}()$$

In this section, I construct:

$$\Omega^{\text{inf}}_{-}(A) : (E^{\infty}\text{-Alg } A)/A \rightleftarrows E^{\infty}\text{-Mod } A : \Lambda^{\text{inf}}_{-}(A)$$

And I also define the concept of an ∞ -derivation.


This adjunction factors like so:

$$\Omega^{\text{inf}}_{-}(A) : (E^{\infty}\text{-Alg } A)/A \rightleftarrows ??? \rightleftarrows E^{\infty}\text{-Mod } A : \Lambda^{\text{inf}}_{-}(A)$$


The more typical concept of a free abelian group .

Given an E^{∞} -algebra...., we can form an

Tensor Product of Lie-Algebras



Tensor Product of L^∞ -Algebras




Connections

s instead of ω and λ

The set of connections forms an affine space. One can then define:

1. Free abelian group action.



∞ -Connections

Connections are elements of the free abelian group action on an A -algebra over A , and ∞ -connections are ...

1. I would like to first construct the lie-algebra representation of flat connections using the spectrum $\omega^{\text{inf}} \cdot \text{obj } X) \cdot \text{obj } V$.
- 2.
3. Lie algebra representations and ...
- 4.
- 5.
- 6.
7. Somehow connections are the dual of the free abelian group action

Some goals:

- 1.
1. smooth (etale locally ???)
2. analytic (etale locally ???)

$$\mathbb{CP}^1, \mathbb{CP}^1 \cong \mathbb{C}(x)$$


3. Our \mathbb{R} is $\partial \cdot \text{obj } \mathbb{Z}$ regarded as an object in $[\vec{\gamma}, \infty_-(\infty\text{-Grpd})]$
4. $B \cdot \text{obj } \det : B \cdot \text{obj } U(n) \longrightarrow B \cdot \text{obj } U(1)$
5. Which E_∞ Space have a Chern class?

Cohomology with coefficients in $[-, B\mathbb{C}^x]$ plays a .


- 1.
2. d_2 is wedge with (representable...)

Given an E^∞ -algebra....., we can form an

Tensor Product of Lie-Algebra Representations



L^∞ -Algebra Representations



Bibliography

1. Samuel Eilenberg and Saunders Mac Lane, "On the Groups $H(\pi, n)$. I", Annals of Mathematics, Second Series, Vol. 58, No. 1 (Jul., 1953), pp. 55-106.
2. Samuel Eilenberg and Saunders Mac Lane, "On the Groups $H(\pi, n)$. II", Annals of Mathematics, Second Series, Vol. 60, No. 1 (Jul., 1954), pp. 49-139.
3. Saunders Mac Lane, "On the Homology Theory of Eilenberg-Mac Lane", Proceedings of the National Academy of Sciences of the United States of America, Vol. 35, No. 11 (Nov. 15, 1949), pp. 657-663.
4. Eilenberg, S., & MacLane, S. (1945). Relations Between Homology and Homotopy Groups of Spaces. Proceedings of the National Academy of Sciences of the United States of America, 31(2), 83-87.

Further reading:

1. The nlab article on ∞ -spaces
2. A blog post of Akhil Mathew explaining how $B^n X \cong \Omega B^{n+1} X$ for an ∞ -space X and $n \geq 2$
3. The n-lab article on the Eckman-Hilton argument
4. Operads, Algebras, and Modules, an exposition of J. P. May.

ND