

Coding Theory Homework 1

Duolei Wang
224010091

June 29, 2026 ~

1 Problem 1.

(a) Let $a = (a_1, a_2, \dots, a_k) \in \mathbb{F}_q^k$ be a vector, $f : \mathbb{F}_q^k \rightarrow \mathbb{F}_q$ be a function defined as:

$$f_a(x_1, \dots, x_k) = \sum_{i=1}^k a_i x_i$$

Show that each element of $\mathbb{F}_q$ is the image of exactly q^{k-1} vectors in $\mathbb{F}_q^k$ under the mapping of f .

Solution. Denote the preimage of an arbitrary vector $b \in \mathbb{F}_q^k$ as:

$$f^{-1}(v) := \left\{ x_i \in \mathbb{F}_q \mid \sum_{i=1}^k a_i x_i = v \right\}$$

Assume the number of solution with fixing the first k coordinates is S_k , the total number of solution is S_0 , we have the iteration equation:

$$S_0 = qS_1$$

As the same, we have:

$$S_i = qS_{i+1}$$

until $S_k = 1$. As a result, the number of solution is exactly:

$$S_0 = \underbrace{q \cdot q \cdot \dots \cdot q}_{k+1} \cdot 1 = q^{k-1}$$

□

(b) Let C be a (n, k, d) code over $\mathbb{F}_q$ and let T be a $q^k \times n$ matrix whose rows are codewords of C . Show that each nonzero elements of $\mathbb{F}_q$ appears in every nonzero column in T exactly q^{k-1} times.

Proof. Assume $x \in \mathbb{F}_q$ be an element, we prove it by contradiction. If there exist some column who has the number of element x less than q^{k-1} , as there are totally q^k rows, there must exist some element $x' \in \mathbb{F}_q$ such that it occurs at more than q^{k-1} rows. As the rows are codewords of C , we know that such rows containing x' should be different. As there are $k-1$ coordinates and q elements for each position, there are at most q^{k-1} different vectors. As a result, we know that there must be some repeating codewords, which is contradict to the hypothesis. By the contradiction, we get the result that there number cannot be strict less or greater than q^{k-1} . $\square$

2 Uniformly Random Parity-Check Matrix

Let $R \in (0, 1)$, $k = \lfloor Rn \rfloor$, and $r = n - k$. Choose a matrix $H \in \mathbb{F}_q^{r \times n}$ uniformly at random, and define C to be the code having H as its parity-check matrix, i.e.:

$$C := \{x \in \mathbb{F}_q^n \mid Hx^T = 0\}$$

Prove that for every fixed δ satisfying that $H(\delta) < 1 - R$, with high probability H has rank r and:

$$d(C) > \delta n$$

Proof. We have:

1. If H is full rank, we have the facts that the first row is nonzero, the $k+1$ row is not the linear combination of the first k rows. The probability of the first row to be nonzero is exactly:

$$1 - \left(\frac{1}{q}\right)^n$$

The probability of the second row is not a linear combination (multiplier) of the first row is:

$$1 - \left(\frac{q}{q^n}\right)$$

As a result, the probability should be:

$$P[H \text{ is full row rank}] = \left(1 - \left(\frac{1}{q}\right)^n\right) \cdot \left(1 - \left(\frac{q}{q^n}\right)\right) \cdot \dots \cdot \left(1 - \left(\frac{q^{r-1}}{q}\right)\right)$$

It's:

$$P[H \text{ is full row rank}] = \prod_{i=0}^{r-1} \left(1 - \frac{1}{q^{n-i}}\right)$$

which is really high if q is large.

2. Firstly, we know that:

$$d(C) = \min_{x \in C} w(x)$$

Thus, we have:

$$P[d(C) > \delta n] = 1 - P[d(C) \leq \delta n]$$

We try to check the bound of the right hand side. Considering that:

$$P[d(C) \leq \delta n] = P[x \in \mathbb{F}_q^n, x \in C, w(x) \leq \delta n]$$

As we know that every vector x in C satisfy that $Hx^T = 0$, we get the probability to be:

$$P[x \in \mathbb{F}_q^n, w(x) \leq \delta n, Hx^T = 0]$$

For a given row, we have:

$$h_i x^T = 0 \iff \sum_{j=1}^n h_{ij} x_j = 0$$

According to the question 1, we know there are q^{n-1} solutions. Thus the probability is

$$\frac{q^{n-1}}{q^n} = \frac{1}{q}$$

As the rows are independent, the probability of $Hx^T = 0$ is exactly $\frac{1}{q^r}$. Thus, we have:

$$P[x \in \mathbb{F}_q^n, w(x) \leq \delta n, Hx^T = 0] \leq \sum_{x, w(x) \leq \delta n} P[Hx^T = 0] = |\{x \mid w(x) \leq \delta n\}| \cdot \frac{1}{q^r}$$

According to the Hamming bound, we know that:

$$|B(0, \lfloor \delta n \rfloor)| \leq q^{H_q(\delta)n}$$

and $H_q(\delta) < 1 - R$; thus, we have $r = n - k = n - \lfloor nR \rfloor \geq n(1 - R)$:

$$P[d(C) \leq \delta n] \leq q^{H_q(\delta)n - n(1-R)} \leq q^{n(1-R+\varepsilon) - n(1-R)} = q^{n\varepsilon} \rightarrow 0$$

As a result, we have:

$$\lim_{n \rightarrow \infty} P[d(C) > \delta n] = 1 - \lim_{n \rightarrow \infty} q^{n\varepsilon} = 1$$

□

3 Minimum distance of a code

Let $R \in (0, 1/2)$ and $M = \lceil q^{Rn} \rceil$, a random q -ary (n, M) code over Σ is obtained by choosing codewords $X_1, X_2, \dots, X_M \in \Sigma^n$ independently, uniformly, and randomly. Let $\delta(C)$ be the relative minimum distance of the random code, we write H_q^{-1} for the inverse of H_q on the interval $[0, 1 - 1/q]$.

(a) Show that for any fixed $\varepsilon > 0$ with $1 - 2R - \varepsilon > 0$, with high probability we have:

$$\delta(C) \geq H_q^{-1}(1 - 2R - \varepsilon) - o(1)$$

Proof. It's equivalent to prove, with high probability:

$$d(C) \geq nH_q^{-1}(1 - 2R - \varepsilon) - o(n)$$

□

(b) Show that for any fixed $\varepsilon > 0$, with high probability we have:

$$\delta(C) \leq H_q^{-1}(1 - 2R) + \varepsilon$$

4 Hamming Ball

Denote the volume of a Hamming Ball of radius in $\mathbb{F}_q^n$ by:

$$V_q(n, t) := \sum_{i=0}^t \binom{n}{i} \cdot (q-1)^i$$

Let $r = n - k$, and let D be an integer satisfying:

$$V_q(n-1, D-2) < q^r$$

Prove that one can construct a $r \times n$ parity-check matrix H of an (n, k, d) linear code over $\mathbb{F}_q$ with $d \geq D$.

5 Q5

Let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{F}_q^n$ be a vector with distinct entries, and $v \in (F_q^*)^n$, prove that for any $1 \leq k \leq n-1$, the dual code of $\text{GRS}_q(\alpha, k, v)$ is $\text{GRS}_q(\alpha, n-k, v')$ for some $v' = (v'_1, v'_2, \dots, v'_n) \in (F_q^*)^n$

Index

hamming distance , 3

t-error channel , 3

block length, 2

bound

Generalized Hamming Bound, 6

code, 2

code

binary code, 2

dimension, 2

perfect, 16

rate, 2

Reed-Solomon Code, 15

encoding function, 3

parity code, 2

repeating code, 3

sphere-packing bound, 16