

Multilinear Regression: CarSeats dataset

Table of contents

Import notebook funcs	1
Import ISLP objects	2
Import User Funactions	2
Load dataset	2
Display dataset stats	2
Set categorical types	3
Standardize variables	3
(a) Fit a multiple regression model to predict Sales using Price, Urban, and US.	3
(b) Provide an interpretation of each coefficient in the model. Be careful—some of the variables in the model are qualitative!	4
(c) Write out the model in equation form, being careful to handle the qualitative vari- ables properly.	4
(d) For which of the predictors can you reject the null hypothesis $H_0 : \beta_j = 0$?	4
(e) On the basis of your response to the previous question, fit a smaller model that only uses the predictors for which there is evidence of association with the outcome.	5
(f) How well do the models in (a) and (e) fit the data?	5
(g) Using the model from (e), obtain 95 % confidence intervals for the coefficient(s).	6
(h) Is there evidence of outliers or high leverage observations in the model from (e)?	6
For a more conservative cutoff values for hat_diag, we have the following influence point(s):	14
Using DFFITS cutoff, we have the following influential points	14
Using Cooks Distance, we have the following influential points	14
Using Cooks Distance p-values, we have the following influential points	14
Using DFBeta for intercept, we have the following influential points	14
Using DFBeta for US, we have the following influential points	14
Using DFBeta for Price, we have the following influential points	15

Import notebook funcs

```
from notebookfuncs import *
```

Import ISLP objects

```
from ISLP import load_data
```

Import User Functions

```
from userfuncs import *
```

Load dataset

```
Carseats = load_data("Carseats")  
Carseats.head()
```

	Sales	CompPrice	Income	Advertising	Population	Price	ShelveLoc	Age	Education	Urban	US
0	9.50	138	73	11	276	120	Bad	42	17	Yes	Yes
1	11.22	111	48	16	260	83	Good	65	10	Yes	Yes
2	10.06	113	35	10	269	80	Medium	59	12	Yes	Yes
3	7.40	117	100	4	466	97	Medium	55	14	Yes	Yes
4	4.15	141	64	3	340	128	Bad	38	13	Yes	No

```
Carseats.shape
```

```
(400, 11)
```

```
Carseats = Carseats.dropna()  
Carseats.shape
```

```
(400, 11)
```

Display dataset stats

```
Carseats.describe()
```

	Sales	CompPrice	Income	Advertising	Population	Price	Age	Education
count	400.000000	400.000000	400.000000	400.000000	400.000000	400.000000	400.000000	400.000000
mean	7.496325	124.975000	68.657500	6.635000	264.840000	115.795000	53.322500	13.900000
std	2.824115	15.334512	27.986037	6.650364	147.376436	23.676664	16.200297	2.620528
min	0.000000	77.000000	21.000000	0.000000	10.000000	24.000000	25.000000	10.000000
25%	5.390000	115.000000	42.750000	0.000000	139.000000	100.000000	39.750000	12.000000

	Sales	CompPrice	Income	Advertising	Population	Price	Age	Education
50%	7.490000	125.000000	69.000000	5.000000	272.000000	117.000000	54.500000	14.000000
75%	9.320000	135.000000	91.000000	12.000000	398.500000	131.000000	66.000000	16.000000
max	16.270000	175.000000	120.000000	29.000000	509.000000	191.000000	80.000000	18.000000

Set categorical types

```
Carseats["US"] = Carseats["US"].astype("category")
Carseats["Urban"] = Carseats["Urban"].astype("category")
```

Standardize variables

```
Carseats["Sales"] = standardize(Carseats["Sales"])
Carseats["Price"] = standardize(Carseats["Price"])
```

(a) Fit a multiple regression model to predict Sales using Price, Urban, and US.

```
formula = "Price + Urban + US"
perform_analysis("Sales", formula, Carseats)
```

```

=====
                        OLS Regression Results
=====
Dep. Variable:          Sales    R-squared:                0.239
Model:                  OLS      Adj. R-squared:           0.234
Method:                 Least Squares    F-statistic:         41.52
Date:                  Tue, 25 Feb 2025    Prob (F-statistic):    2.39e-23
Time:                  14:40:07      Log-Likelihood:       -512.88
No. Observations:      400          AIC:                  1034.
Df Residuals:          396          BIC:                  1050.
Df Model:               3
Covariance Type:       nonrobust
=====
                        coef    std err          t      P>|t|      [0.025    0.975]
-----
Intercept             -0.2691      0.098      -2.734      0.007     -0.463    -0.076
Urban[T.Yes]          -0.0078      0.096     -0.081      0.936     -0.197     0.182
US[T.Yes]              0.4256      0.092      4.635      0.000      0.245     0.606
Price                 -0.4566      0.044     -10.389      0.000     -0.543    -0.370
=====
Omnibus:               0.676    Durbin-Watson:           1.912

```

```

Prob(Omnibus):          0.713   Jarque-Bera (JB):          0.758
Skew:                  0.093   Prob(JB):          0.684
Kurtosis:              2.897   Cond. No.          4.27
=====

```

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

	df	sum_sq	mean_sq	F	PR(>F)
Urban	1.0	0.095104	0.095104	0.123767	7.251713e-01
US	1.0	12.675983	12.675983	16.496407	5.877444e-05
Price	1.0	82.939070	82.939070	107.936143	1.609917e-22
Residual	396.0	304.289843	0.768409	NaN	NaN

<statsmodels.regression.linear_model.RegressionResultsWrapper at 0x7fe79017d160>

(b) Provide an interpretation of each coefficient in the model. Be careful—some of the variables in the model are qualitative!

- The coefficient of -0.0078 for Urban (True) indicates that -0.0078 of the SD of Sales can be explained by the level urban store as compared to a rural store. However, the p-value of 0.936 indicates that this difference is not significant and can be discounted or discarded.
- The coefficient of 0.4256 for US (True) indicates that 0.4256 of the typical deviation of Sales are explained by a US store as compared to a non-US store.
- The coefficient of -0.4566 for Price indicates that -0.4566 of the typical deviation of Sales is explained by one SD of change in the Price variable.
- We can also conclude that Price has the highest effect on Sales, the response variable, since the absolute value of its coefficient 0.4566 is the highest amongst all the coefficients.
- <https://blogs.sas.com/content/iml/2023/07/17/standardize-reg-coeff-class.html>
- <https://www.statlect.com/fundamentals-of-statistics/linear-regression-with-standardized-variables>

(c) Write out the model in equation form, being careful to handle the qualitative variables properly.

- The equation can be written out as follows:
- Sales = -0.0078 * Urban + 0.4256 * US -0.4566 * Price (Standardized) - 0.2691

(d) For which of the predictors can you reject the null hypothesis $H_0 : \beta_j = 0$?

- The p-value for the Urban predictor is 0.936 which is much higher than our chosen level of significance 0.01. So we cannot reject the null Hypothesis in this case that its coefficient is zero.
- The p-values for US, Price and Intercept are zero. Hence, we reject the null hypothesis for them that their coefficients are zero.

(e) On the basis of your response to the previous question, fit a smaller model that only uses the predictors for which there is evidence of association with the outcome.

```
formula = "Price + US"
results = perform_analysis("Sales", formula, Carseats)
```

```

=====
                        OLS Regression Results
=====
Dep. Variable:          Sales    R-squared:                0.239
Model:                  OLS      Adj. R-squared:           0.235
Method:                 Least Squares    F-statistic:         62.43
Date:                  Tue, 25 Feb 2025    Prob (F-statistic):    2.66e-24
Time:                  14:40:07    Log-Likelihood:       -512.88
No. Observations:      400      AIC:                  1032.
Df Residuals:          397      BIC:                  1044.
Df Model:               2
Covariance Type:       nonrobust
=====

```

	coef	std err	t	P> t	[0.025	0.975]
Intercept	-0.2743	0.074	-3.730	0.000	-0.419	-0.130
US[T.Yes]	0.4253	0.092	4.641	0.000	0.245	0.605
Price	-0.4567	0.044	-10.416	0.000	-0.543	-0.371

```

=====
Omnibus:                0.666    Durbin-Watson:           1.912
Prob(Omnibus):          0.717    Jarque-Bera (JB):        0.749
Skew:                   0.092    Prob(JB):                0.688
Kurtosis:               2.895    Cond. No.:               3.12
=====

```

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

	df	sum_sq	mean_sq	F	PR(>F)
US	1.0	12.544810	12.544810	16.366658	6.273751e-05
Price	1.0	83.160345	83.160345	108.495617	1.272157e-22
Residual	397.0	304.294845	0.766486	NaN	NaN

<statsmodels.regression.linear_model.RegressionResultsWrapper at 0x7fe74d1bf8c0>

(f) How well do the models in (a) and (e) fit the data?

- Model(a) has an explanatory value R² adjusted value of 0.234
- Model(e) has an explanatory value R² adjusted value of 0.235


(g) Using the model from (e), obtain 95 % confidence intervals for the coefficient(s).

From the summary analysis, it can be seen that the 95% confidence limits for the three terms are as follows: + Intercept (-0.419, -0.130) + US[T.Yes] (0.245, 0.605) + Price (-0.543, -0.371) + None of them include zero in their range unlike that for Urban[T.Yes] in Model(a) which is another indicator that the coefficient is not significant.

(h) Is there evidence of outliers or high leverage observations in the model from (e)?

We can check for presence of outliers by plotting the residuals plot and seeing if there are any outliers.


```
display_residuals_plot(results)
```



- From the plot above, there doesn't appear to be any obvious outliers.


We can plot studentized residuals to see whether there are any visible there.

```
display_studentized_residuals(results)
```

- From the above plot, no observation lies outside the $(-3,3)$ range. Hence, we can safely conclude that there are no evident outliers in the dataset.


```
display_hat_leverage_cutoffs(results)
```




- We can see from the above graph that we have a few leverage points that exceed the cutoff of 3 * average leverage value. These are plotted in red.
- The ones in yellow exceed the less conservative estimate of 2 * average leverage value
- We could also use more conservative estimates for the cutoff of either 4 * average leverage value or 5 * average cutoff value


References: - <https://online.stat.psu.edu/stat501/lesson/11/11.2>


```
display_cooks_distance_plot(results)
```





```
display_DFFITS_plot(results)
```





```
display_hat_leverage_plot(results)
```

Unable to display output for mime type(s): text/html

Unable to display output for mime type(s): text/html

- Looking at all three studentized plots for leverage, it can be concluded that even if there are a few outliers, none wield a significant influence on the regression since the points with high leverage values have low studentized residual values.

```
inf_df, _ = get_influence_points(results)
inf_df
print(_)
```

```
n = 400.0, p = 3
```

```
Average Hat Leverage: 0.0075000000000000015
```

```
Hat Leverage Cutoff = 2 * Average Hat Leverage = 0.015000000000000003
```

```
DFBetas Cutoff = 3 / sqrt(n) = 0.15
```

```
DFFITS Cutoff = 2 * sqrt(p/n) = 0.17320508075688773
```

```
Cooks Distance Cutoff = 1.0
```

```
Cooks Distance p-value Cutoff = 0.05
```

```
Studentized Residuals Cutoff = 3.0
```

```
Studentized Residuals p-value Cutoff = 0.01
```

```
{'n': 400.0, 'p': 3, 'average_hat': 0.0075000000000000015, 'hat_leverage_cutoff': 0.015000000000000003}
```

For a more conservative cutoff values for `hat_diag`, we have the following influence point(s):

```
inf_df[inf_df["hat_diag"] > (3 * np.mean(inf_df["hat_diag"]))]
```

	dfb_Intercept	dfb_US[T.Yes]	dfb_Price	cooks_d	hat_diag	student_resid	dffits	student_resid_pvalue
--	---------------	---------------	-----------	---------	----------	---------------	--------	----------------------

Using DFFITS cutoff, we have the following influential points

```
inf_df[inf_df["dffits"] > 2 * np.sqrt(len(results.params) / results.nobs)]
```

	dfb_Intercept	dfb_US[T.Yes]	dfb_Price	cooks_d	hat_diag	student_resid	dffits	student_resid_pvalue
25	0.210972	-0.165700	-0.176952	0.026109	0.011622	2.599652	0.281894	0.004840
68	0.004485	0.092526	0.096454	0.011988	0.005202	2.642364	0.191069	0.004280
376	-0.007088	0.116257	-0.152448	0.018282	0.006637	2.891521	0.236355	0.002023

Using Cooks Distance, we have the following influential points

```
inf_df[inf_df["cooks_d"] > 1.0]
```

	dfb_Intercept	dfb_US[T.Yes]	dfb_Price	cooks_d	hat_diag	student_resid	dffits	student_resid_pvalue
--	---------------	---------------	-----------	---------	----------	---------------	--------	----------------------

Using Cooks Distance p-values, we have the following influential points

```
inf_df[inf_df["cooks_d_pvalue"] < 0.05]
```

	dfb_Intercept	dfb_US[T.Yes]	dfb_Price	cooks_d	hat_diag	student_resid	dffits	student_resid_pvalue
--	---------------	---------------	-----------	---------	----------	---------------	--------	----------------------

Using DFBeta for intercept, we have the following influential points

```
inf_df[inf_df["dfb_Intercept"] > (3 / np.sqrt(results.nobs))]
```

	dfb_Intercept	dfb_US[T.Yes]	dfb_Price	cooks_d	hat_diag	student_resid	dffits	student_resid_pvalue
25	0.210972	-0.1657	-0.176952	0.026109	0.011622	2.599652	0.281894	0.00484

Using DFBeta for US, we have the following influential points

```
inf_df[inf_df["dfb_US[T.Yes]"] > (3 / np.sqrt(results.nobs))]
```

dfb_Intercept	dfb_US[T.Yes]	dfb_Price	cooks_d	hat_diag	student_resid	dffits	student_resid_pvalue
---------------	---------------	-----------	---------	----------	---------------	--------	----------------------

Using DFBeta for Price, we have the following influential points

```
inf_df[inf_df["dfb_Price"] > (3 / np.sqrt(results.nobs))]
```

dfb_Intercept	dfb_US[T.Yes]	dfb_Price	cooks_d	hat_diag	student_resid	dffits	student_resid_pvalue
---------------	---------------	-----------	---------	----------	---------------	--------	----------------------

```
allDone()
```

```
<IPython.lib.display.Audio object>
```