

Equilibrium measures on curves

Damian Dąbrowski

Institute of Mathematics of the Polish Academy of Sciences (IMPAN)



Funded by
the European Union



European Research Council
Established by the European Commission



Suomen Akatemia
Finlands Akademi
Research Council of Finland

What is an equilibrium measure?

Logarithmic energy

Given a compactly supported Radon measure μ on \mathbb{R}^n its **logarithmic energy** is

$$\mathcal{I}(\mu) := \iint \log \frac{1}{|x - y|} d\mu(x) d\mu(y) \in (-\infty, +\infty].$$

Given a compactly supported Radon measure μ on \mathbb{R}^n its **logarithmic energy** is

$$\mathcal{I}(\mu) := \iint \log \frac{1}{|x-y|} d\mu(x)d\mu(y) \in (-\infty, +\infty].$$

Examples:

- $\mathcal{I}(\delta_0) = +\infty$
- $\mathcal{I}(\mu) < +\infty$ whenever μ is α -Frostman for some $\alpha > 0$, i.e.

$$\mu(B(x, r)) < Cr^\alpha \quad x \in \mathbb{R}^n, r > 0.$$

Equilibrium measures

Suppose that $E \subset \mathbb{R}^n$ is compact, and let $\mathcal{P}(E) := \{\mu : \text{supp } \mu \subset E, \mu(E) = 1\}$.

Problem

Minimize $\mathcal{I}(\mu)$ among $\mu \in \mathcal{P}(E)$.

Equilibrium measures

Suppose that $E \subset \mathbb{R}^n$ is compact, and let $\mathcal{P}(E) := \{\mu : \text{supp } \mu \subset E, \mu(E) = 1\}$.

Problem

Minimize $\mathcal{I}(\mu)$ among $\mu \in \mathcal{P}(E)$.

If for all $\mu \in \mathcal{P}(E)$ we have $\mathcal{I}(\mu) = +\infty$, nothing to do.

- Example: $E = \{0\}$.

Equilibrium measures

Suppose that $E \subset \mathbb{R}^n$ is compact, and let $\mathcal{P}(E) := \{\mu : \text{supp } \mu \subset E, \mu(E) = 1\}$.

Problem

Minimize $\mathcal{I}(\mu)$ among $\mu \in \mathcal{P}(E)$.

If for all $\mu \in \mathcal{P}(E)$ we have $\mathcal{I}(\mu) = +\infty$, nothing to do.

- Example: $E = \{0\}$.

Theorem (Frostman, Fuglede)

If there is at least one $\nu \in \mathcal{P}(E)$ with $\mathcal{I}(\nu) < \infty$, then there exists a unique minimizer of $\mathcal{I}(\mu)$ among $\mu \in \mathcal{P}(E)$.

This unique minimizer is called the **(logarithmic) equilibrium measure for E** .

Examples

- $E = [-1, 1]$




Examples

• $E = [-1, 1]$



\rightsquigarrow

$$\mu = \frac{c}{\sqrt{1-x^2}} \mathcal{H}^1|_{[-1,1]}$$





Examples

- $E = [-1, 1]$ \rightsquigarrow $\mu = \frac{c}{\sqrt{1-x^2}} \mathcal{H}^1|_{[-1,1]}$
- $E = \mathbb{S}^1$ or $E = B(0, 1) \subset \mathbb{R}^2$





Examples

- $E = [-1, 1]$ \rightsquigarrow $\mu = \frac{c}{\sqrt{1-x^2}} \mathcal{H}^1|_{[-1,1]}$
- $E = \mathbb{S}^1$ or $E = B(0, 1) \subset \mathbb{R}^2$ \rightsquigarrow $\mu = \mathcal{H}^1|_{\mathbb{S}^1}$



Examples

- $E = [-1, 1]$ \rightsquigarrow $\mu = \frac{c}{\sqrt{1-x^2}} \mathcal{H}^1|_{[-1,1]}$
- $E = \mathbb{S}^1$ or $E = B(0, 1) \subset \mathbb{R}^2$ \rightsquigarrow $\mu = \mathcal{H}^1|_{\mathbb{S}^1}$
- $E = \mathbb{S}^{n-1}$ or $E = B(0, 1) \subset \mathbb{R}^n$ \rightsquigarrow $\mu = \mathcal{H}^{n-1}|_{\mathbb{S}^{n-1}}$



Why is this interesting?

- If $n = 2$, then μ models the distribution of electric charges inside a conducting material.

Why is this interesting?

- If $n = 2$, then μ models the distribution of electric charges inside a conducting material.
- If $n = 2$ and $E = \partial\Omega$ for a Wiener regular unbounded open set, then μ is the harmonic measure with pole at ∞ for Ω .

Why is this interesting?

- If $n = 2$, then μ models the distribution of electric charges inside a conducting material.
- If $n = 2$ and $E = \partial\Omega$ for a Wiener regular unbounded open set, then μ is the harmonic measure with pole at ∞ for Ω .
- Identifying the equilibrium measure is a limit version of an important discrete problem: for a fixed $E \subset \mathbb{R}^n$ and $N \in \mathbb{N}$, what's the configuration $x_1, \dots, x_N \subset E$ minimizing

$$\sum_{i=1}^N \sum_{j=1}^N \log \frac{1}{|x_i - x_j|}?$$

Why is this interesting?

- If $n = 2$, then μ models the distribution of electric charges inside a conducting material.
- If $n = 2$ and $E = \partial\Omega$ for a Wiener regular unbounded open set, then μ is the harmonic measure with pole at ∞ for Ω .
- Identifying the equilibrium measure is a limit version of an important discrete problem: for a fixed $E \subset \mathbb{R}^n$ and $N \in \mathbb{N}$, what's the configuration $x_1, \dots, x_N \subset E$ minimizing

$$\sum_{i=1}^N \sum_{j=1}^N \log \frac{1}{|x_i - x_j|}?$$

- For $n \geq 3$ a connection to higher order PDEs?

Main result

- If $n = 2$, a lot of results on the structure of equilibrium measures. For example:

Theorem (F. and M. Riesz, 1916)

If $\Gamma \subset \mathbb{R}^2$ is a Jordan curve of finite length, then

$$\mathcal{H}^1|_{\Gamma} \ll \mu \ll \mathcal{H}^1|_{\Gamma}.$$

- If $n \geq 3$: very little is known.

Main result

- If $n = 2$, a lot of results on the structure of equilibrium measures. For example:

Theorem (F. and M. Riesz, 1916)

If $\Gamma \subset \mathbb{R}^2$ is a Jordan curve of finite length, then

$$\mathcal{H}^1|_{\Gamma} \ll \mu \ll \mathcal{H}^1|_{\Gamma}.$$

- If $n \geq 3$: very little is known.

Theorem (D.-Orponen '25)

If $\Gamma \subset \mathbb{R}^n$ is a $C^{1,\alpha}$ curve with $\alpha > 0$, then $\mu \ll \mathcal{H}^1|_{\Gamma}$.

Before nothing was known even for C^∞ curves.

Logarithmic potentials

Logarithmic potentials

Let ν be a compactly supported Radon measure on \mathbb{R}^n . Its **logarithmic potential** is

$$\mathcal{U}\nu(x) := \int \log \frac{1}{|x - y|} d\nu(y).$$

Logarithmic potentials

Let ν be a compactly supported Radon measure on \mathbb{R}^n . Its **logarithmic potential** is

$$\mathcal{U}\nu(x) := \int \log \frac{1}{|x-y|} d\nu(y).$$

Remarks:

- $\mathcal{U}\nu \in C^\infty(\mathbb{R}^n \setminus \text{supp } \nu)$

Logarithmic potentials

Let ν be a compactly supported Radon measure on \mathbb{R}^n . Its **logarithmic potential** is

$$\mathcal{U}\nu(x) := \int \log \frac{1}{|x-y|} d\nu(y).$$

Remarks:

- $\mathcal{U}\nu \in C^\infty(\mathbb{R}^n \setminus \text{supp } \nu)$
- $\mathcal{U}\nu(x) \sim -\log |x|$ as $|x| \rightarrow \infty$

Logarithmic potentials

Let ν be a compactly supported Radon measure on \mathbb{R}^n . Its **logarithmic potential** is

$$\mathcal{U}\nu(x) := \int \log \frac{1}{|x-y|} d\nu(y).$$

Remarks:

- $\mathcal{U}\nu \in C^\infty(\mathbb{R}^n \setminus \text{supp } \nu)$
- $\mathcal{U}\nu(x) \sim -\log |x|$ as $|x| \rightarrow \infty$
- $\mathcal{I}(\nu) = \int \mathcal{U}\nu(x) d\nu(x)$

Logarithmic potentials

Let ν be a compactly supported Radon measure on \mathbb{R}^n . Its **logarithmic potential** is

$$\mathcal{U}\nu(x) := \int \log \frac{1}{|x-y|} d\nu(y).$$

Remarks:

- $\mathcal{U}\nu \in C^\infty(\mathbb{R}^n \setminus \text{supp } \nu)$
- $\mathcal{U}\nu(x) \sim -\log |x|$ as $|x| \rightarrow \infty$
- $\mathcal{I}(\nu) = \int \mathcal{U}\nu(x) d\nu(x)$
- $\Delta^{n/2}(\mathcal{U}\nu) = \nu$ in a weak sense
 - \rightsquigarrow if $n = 2$ then $\mathcal{U}\nu$ is harmonic on $\mathbb{R}^2 \setminus \text{supp } \nu$
 - \rightsquigarrow if $n = 4$ then $\mathcal{U}\nu$ is biharmonic on $\mathbb{R}^4 \setminus \text{supp } \nu$

Why are they useful?

We can use potentials to

- solve PDEs,

Why are they useful?

We can use potentials to

- solve PDEs,
- **study measures.**

regularity of potentials \Leftrightarrow regularity of measures

Why are they useful?

We can use potentials to

- solve PDEs,
- **study measures.**

regularity of potentials \Leftrightarrow regularity of measures

Theorem (Carleson, Wallin '60s)

For $0 < \alpha < 1$

$$\mathcal{U}\nu \in C^\alpha(\mathbb{R}^n) \Leftrightarrow \nu(B(x, r)) \leq Cr^\alpha.$$

Why are they useful?

We can use potentials to

- solve PDEs,
- **study measures.**

regularity of potentials \Leftrightarrow regularity of measures

Theorem (Carleson, Wallin '60s)

For $0 < \alpha < 1$

$$\mathcal{U}\nu \in C^\alpha(\mathbb{R}^n) \quad \Leftrightarrow \quad \nu(B(x, r)) \leq Cr^\alpha.$$

If we show that our equilibrium measure satisfies $\mathcal{U}\mu \in \text{Lip}(\mathbb{R}^n)$, we'll get $\mu \in L^\infty(\mathcal{H}^1|_\Gamma)$!

Potentials of equilibrium measures

Theorem (Frostman, Fuglede)

If μ is the equilibrium measure on E , then its potential $\mathcal{U}\mu$ satisfies




E

Potentials of equilibrium measures

Theorem (Frostman, Fuglede)

If μ is the equilibrium measure on E , then its potential \mathcal{U}_μ satisfies

- $\mathcal{U}_\mu(x) \leq \mathcal{I}(\mu)$ for all $x \in \text{supp } \mu$




Potentials of equilibrium measures

Theorem (Frostman, Fuglede)

If μ is the equilibrium measure on E , then its potential $\mathcal{U}\mu$ satisfies

- $\mathcal{U}\mu(x) \leq \mathcal{I}(\mu)$ for all $x \in \text{supp } \mu$
- $\mathcal{U}\mu(x) \geq \mathcal{I}(\mu)$ for “approximately all” $x \in E$




Potentials of equilibrium measures

Theorem (Frostman, Fuglede)

If μ is the equilibrium measure on E , then its potential $\mathcal{U}\mu$ satisfies

- $\mathcal{U}\mu(x) \leq \mathcal{I}(\mu)$ for all $x \in \text{supp } \mu$
- $\mathcal{U}\mu(x) \geq \mathcal{I}(\mu)$ for “approximately all” $x \in E$
- $\mathcal{U}\mu$ is continuous on \mathbb{R}^n .




Potentials of equilibrium measures

Theorem (Frostman, Fuglede)

If μ is the equilibrium measure on E , then its potential \mathcal{U}_μ satisfies

- $\mathcal{U}_\mu(x) \leq \mathcal{I}(\mu)$ for all $x \in \text{supp } \mu$
- $\mathcal{U}_\mu(x) \geq \mathcal{I}(\mu)$ for “approximately all” $x \in E$
- \mathcal{U}_μ is continuous on \mathbb{R}^n .



Potentials of equilibrium measures

Theorem (Frostman, Fuglede)

If μ is the equilibrium measure on E , then its potential $\mathcal{U}\mu$ satisfies

- $\mathcal{U}\mu(x) \leq \mathcal{I}(\mu)$ for all $x \in \text{supp } \mu$
- $\mathcal{U}\mu(x) \geq \mathcal{I}(\mu)$ for “approximately all” $x \in E$
- $\mathcal{U}\mu$ is continuous on \mathbb{R}^n .

Problem

Given the equilibrium measure μ on a $C^{1,\alpha}$ -graph Γ , how do we use the information above to conclude $\mathcal{U}\mu \in \text{Lip}(\mathbb{R}^n)$?

Potentials of equilibrium measures

Theorem (Frostman, Fuglede)

If μ is the equilibrium measure on E , then its potential $\mathcal{U}\mu$ satisfies

- $\mathcal{U}\mu(x) \leq \mathcal{I}(\mu)$ for all $x \in \text{supp } \mu$
- $\mathcal{U}\mu(x) \geq \mathcal{I}(\mu)$ for “approximately all” $x \in E$
- $\mathcal{U}\mu$ is continuous on \mathbb{R}^n .

Problem

Given the equilibrium measure μ on a $C^{1,\alpha}$ -graph Γ , how do we use the information above to conclude $\mathcal{U}\mu \in \text{Lip}(\mathbb{R}^n)$?

Idea: perhaps it's enough to show that $\mathcal{U}\mu \in \text{Lip}(\Gamma)$?

Goal

If μ is the equilibrium measure on a $C^{1,\alpha}$ -graph Γ , then $\mu \ll \mathcal{H}^1|_{\Gamma}$.

Plan of the proof

Goal

If μ is the equilibrium measure on a $C^{1,\alpha}$ -graph Γ , then $\mu \ll \mathcal{H}^1|_{\Gamma}$.

Step 1. Show that

$$\mathcal{U}\mu \in \text{Lip}(\Gamma) \quad \Rightarrow \quad \mu(B(x, r)) \leq Cr, \text{ i.e. } \mu \in L^\infty(\Gamma)$$

Goal

If μ is the equilibrium measure on a $C^{1,\alpha}$ -graph Γ , then $\mu \ll \mathcal{H}^1|_{\Gamma}$.

Step 1. Show that

$$\mathcal{U}\mu \in \dot{H}^{1,p}(\Gamma) \quad \Rightarrow \quad \mu \in L^p(\Gamma)$$

Plan of the proof

Goal

If μ is the equilibrium measure on a $C^{1,\alpha}$ -graph Γ , then $\mu \ll \mathcal{H}^1|_{\Gamma}$.

Step 1. Show that

$$\mathcal{U}\mu \in \dot{H}^{1,p}(\Gamma) \quad \Rightarrow \quad \mu \in L^p(\Gamma)$$

Step 2. Prove Sobolev regularity of $\mathcal{U}\mu$ on Γ .

Step 1: Potentials seen “intrinsically”

Theorem (Frostman, Fuglede)

Given a compact set E and measures $\mu, \nu \in \mathcal{P}(E)$ we have

$$\mathcal{U}\mu|_E = \mathcal{U}\nu|_E \quad \Leftrightarrow \quad \mu = \nu.$$

Thus, the operator $\mathcal{U} : \mathcal{P}(E) \rightarrow \{\text{functions on } E\}$ is injective.

- What are its mapping properties on subspaces of $\mathcal{P}(E)$?
- When does it have a bounded inverse?

Theorem (Carleson, Wallin '60s)

For $0 < \beta < 1$

$$\mathcal{U}\nu \in C^\beta(\mathbb{R}^n) \quad \Leftrightarrow \quad \nu(B(x, r)) \leq Cr^\beta.$$

Theorem (Carleson, Wallin '60s)

For $0 < \beta < 1$

$$\mathcal{U}\nu \in C^\beta(\mathbb{R}^n) \iff \nu(B(x, r)) \leq Cr^\beta.$$

Proposition (D.-Orponen '25)

[almost true]

If Γ is a Lipschitz graph with $\text{Lip}(\Gamma) \ll 1$, then for $\nu \in \mathcal{P}(\Gamma)$

$$\mathcal{U}\nu \in C^\beta(\Gamma) \iff \nu(B(x, r)) \leq Cr^\beta.$$

Theorem (Carleson, Wallin '60s)

For $0 < \beta < 1$

$$\mathcal{U}\nu \in C^\beta(\mathbb{R}^n) \iff \nu(B(x, r)) \leq Cr^\beta.$$

Proposition (D.-Orponen '25)

[almost true]

If Γ is a Lipschitz graph with $\text{Lip}(\Gamma) \ll 1$, then for $\nu \in \mathcal{P}(\Gamma)$

$$\mathcal{U}\nu \in C^\beta(\Gamma) \iff \nu(B(x, r)) \leq Cr^\beta.$$

Almost-almost true: for $\beta \in [0, 1]$ and $p \in (1, \infty)$ we show that

$$\mathcal{U} : \dot{H}^{\beta-1,p}(\Gamma) \rightarrow \dot{H}^{\beta,p}(\Gamma)$$


is bounded with bounded inverse, provided $\text{Lip}(\Gamma) \ll_p 1$.

Graph potentials

Given $A : \mathbb{R} \rightarrow \mathbb{R}^{n-1}$ denote by $\Gamma(x) = (x, A(x)) \in \mathbb{R}^n$ the graph map.

Given a compactly supported measure ν on \mathbb{R} we define its **graph potential** $\mathcal{U}^\Gamma : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\mathcal{U}^\Gamma \nu(x) = \int_{\mathbb{R}} \log \frac{1}{|\Gamma(x) - \Gamma(y)|} d\nu(y).$$



Graph potentials

Given $A : \mathbb{R} \rightarrow \mathbb{R}^{n-1}$ denote by $\Gamma(x) = (x, A(x)) \in \mathbb{R}^n$ the graph map.

Given a compactly supported measure ν on \mathbb{R} we define its **graph potential** $\mathcal{U}^\Gamma : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\mathcal{U}^\Gamma \nu(x) = \int_{\mathbb{R}} \log \frac{1}{|\Gamma(x) - \Gamma(y)|} d\nu(y).$$

Note that

measure ν on Γ and $\mathcal{U}\nu : \Gamma \rightarrow \mathbb{R} \iff$ measure $\tilde{\nu}$ on \mathbb{R} and $\mathcal{U}^\Gamma \tilde{\nu} : \mathbb{R} \rightarrow \mathbb{R}$.

$$\tilde{\nu} = (\pi_1)_* \nu$$

$$\mathcal{U}^\Gamma \tilde{\nu}(x) = \mathcal{U}\nu(\Gamma(x))$$

Graph potentials

Given $A : \mathbb{R} \rightarrow \mathbb{R}^{n-1}$ denote by $\Gamma(x) = (x, A(x)) \in \mathbb{R}^n$ the graph map.

Given a compactly supported measure ν on \mathbb{R} we define its **graph potential** $\mathcal{U}^\Gamma : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\mathcal{U}^\Gamma \nu(x) = \int_{\mathbb{R}} \log \frac{1}{|\Gamma(x) - \Gamma(y)|} d\nu(y).$$

Proposition (D.-Orponen '25)

For $\beta \in [0, 1]$ and $p \in (1, \infty)$ the operators

$$T_\beta^\Gamma := \Delta^{(1-\beta)/2} \mathcal{U}^\Gamma \Delta^{\beta/2}$$

are bounded $L^p \rightarrow L^p$, and invertible provided $\text{Lip}(\Gamma) \ll_p 1$.

Proposition (D.-Orponen '25)

For $\beta \in [0, 1]$ and $p \in (1, \infty)$ the operators

$$T_{\beta}^{\Gamma} := \Delta^{(1-\beta)/2} \mathcal{U}^{\Gamma} \Delta^{\beta/2}$$

are bounded $L^p \rightarrow L^p$, and invertible provided $\text{Lip}(\Gamma) \ll_p 1$.

- In the flat case $A \equiv 0$ we have $T_{\beta}^{\Gamma} = \text{id}$.

Proposition (D.-Orponen '25)

For $\beta \in [0, 1]$ and $p \in (1, \infty)$ the operators

$$\mathcal{T}_\beta^\Gamma := \Delta^{(1-\beta)/2} \mathcal{U}^\Gamma \Delta^{\beta/2}$$

are bounded $L^p \rightarrow L^p$, and invertible provided $\text{Lip}(\Gamma) \ll_p 1$.

- In the flat case $A \equiv 0$ we have $\mathcal{T}_\beta^\Gamma = \text{id}$.
- We employ the method of continuity to show that

$$\|\mathcal{T}_\beta^\Gamma - \text{id}\|_{L^p \rightarrow L^p} \lesssim_p \text{Lip}(A).$$

Proposition (D.-Orponen '25)

For $\beta \in [0, 1]$ and $p \in (1, \infty)$ the operators

$$\mathcal{T}_\beta^\Gamma := \Delta^{(1-\beta)/2} \mathcal{U}^\Gamma \Delta^{\beta/2}$$

are bounded $L^p \rightarrow L^p$, and invertible provided $\text{Lip}(\Gamma) \ll_p 1$.

- In the flat case $A \equiv 0$ we have $\mathcal{T}_\beta^\Gamma = \text{id}$.
- We employ the method of continuity to show that

$$\|\mathcal{T}_\beta^\Gamma - \text{id}\|_{L^p \rightarrow L^p} \lesssim_p \text{Lip}(A).$$

- The case $\beta \in \{0, 1\}$ relies on the L^p -boundedness of nice enough SIOs on Lipschitz graphs.

Proposition (D.-Orponen '25)

For $\beta \in [0, 1]$ and $p \in (1, \infty)$ the operators

$$T_\beta^\Gamma := \Delta^{(1-\beta)/2} \mathcal{U}^\Gamma \Delta^{\beta/2}$$

are bounded $L^p \rightarrow L^p$, and invertible provided $\text{Lip}(\Gamma) \ll_p 1$.

- In the flat case $A \equiv 0$ we have $T_\beta^\Gamma = \text{id}$.
- We employ the method of continuity to show that

$$\|T_\beta^\Gamma - \text{id}\|_{L^p \rightarrow L^p} \lesssim_p \text{Lip}(A).$$

- The case $\beta \in \{0, 1\}$ relies on the L^p -boundedness of nice enough SIOs on Lipschitz graphs.
- For $\beta \in (0, 1)$ we use complex interpolation.

In what follows, we'll pretend we have the "almost true" result:

Proposition

If Γ is a Lipschitz graph with $\text{Lip}(\Gamma) \ll 1$, then for $\nu \in \mathcal{P}([0, 1])$

$$\mathcal{U}^\Gamma \nu \in C^\beta([0, 1]) \quad \Leftrightarrow \quad \nu(B(x, r)) \leq Cr^\beta.$$

We move on to **Step 2**:

Goal


If μ is the (projection of the) equilibrium measure on a $C^{1,\alpha}$ -graph Γ , then $\mathcal{U}^\Gamma \mu \in \text{Lip}([0, 1])$.

This will imply $\mu \ll \mathcal{H}^1$.

Step 2: Lipschitz regularity of \mathcal{U}^Γ_μ


Planar case

- Recall: for the equilibrium measure we have $\mathcal{U}^\Gamma_\mu = \text{const}$ on $\text{supp } \mu$.



Planar case

- Recall: for the equilibrium measure we have $\mathcal{U}^\Gamma \mu = \text{const}$ on $\text{supp } \mu$.
- If we knew $\text{supp } \mu = [0, 1]$, then we'd have $\mathcal{U}^\Gamma \mu = \text{const}$ on $[0, 1]$ and in particular $\mathcal{U}^\Gamma \in \text{Lip}([0, 1])$.



Planar case

- Recall: for the equilibrium measure we have $\mathcal{U}^\Gamma \mu = \text{const}$ on $\text{supp } \mu$.
- If we knew $\text{supp } \mu = [0, 1]$, then we'd have $\mathcal{U}^\Gamma \mu = \text{const}$ on $[0, 1]$ and in particular $\mathcal{U}^\Gamma \in \text{Lip}([0, 1])$.
- This is the case in \mathbb{R}^2 !

Theorem (Frostman)

If $E \subset \mathbb{R}^2$ is compact and ν is the equilibrium measure on E , then the potential $\mathcal{U}\nu$ is constant on E .


- Reason: $\mathcal{U}\nu$ is harmonic on $\mathbb{R}^2 \setminus \text{supp } \nu$, and we have the maximum principle.
- This does not work in higher dimensions:

$$\Delta^{n/2}(\mathcal{U}\nu) = \nu$$

Kernel decomposition

Our graph potential has kernel

$$k^\Gamma(x, y) = -\log |\Gamma(x) - \Gamma(y)| = -\log |(x - y, A(x) - A(y))|.$$




Kernel decomposition

Our graph potential has kernel

$$K^\Gamma(x, y) = -\log |\Gamma(x) - \Gamma(y)| = -\log |(x - y, A(x) - A(y))|.$$

We decompose the kernel $K^\Gamma(x, y) = P(x, y) + R(x, y)$, where

$$\begin{aligned} P(x, y) &= -\log |(x - y, \nabla A * \psi_{|x-y|}(x)(x - y))| \\ &= -\log |x - y| - \log |(1, \nabla A * \psi_{|x-y|}(x))| \end{aligned}$$



Potential decomposition

We use the kernel decomposition to decompose the potential $\mathcal{U}^\Gamma_\mu = \mathcal{P}_\mu + \mathcal{R}_\mu$:

$$\mathcal{P}_\mu(x) := \int P(x, y) d\mu(y),$$

$$\mathcal{R}_\mu(x) := \int R(x, y) d\mu(y).$$

Potential decomposition

We use the kernel decomposition to decompose the potential $\mathcal{U}^\Gamma_\mu = \mathcal{P}_\mu + \mathcal{R}_\mu$:

$$\mathcal{P}_\mu(x) := \int P(x, y) d\mu(y),$$

$$\mathcal{R}_\mu(x) := \int R(x, y) d\mu(y).$$

\mathcal{P} -Lemma

If $\text{Lip}(A) \ll 1$, then \mathcal{P}_μ is convex on $\mathbb{R} \setminus \text{supp } \mu$.

Potential decomposition

We use the kernel decomposition to decompose the potential $\mathcal{U}^\Gamma \mu = \mathcal{P}\mu + \mathcal{R}\mu$:

$$\mathcal{P}\mu(x) := \int P(x, y) d\mu(y),$$

$$\mathcal{R}\mu(x) := \int R(x, y) d\mu(y).$$

\mathcal{P} -Lemma

If $\text{Lip}(A) \ll 1$, then $\mathcal{P}\mu$ is convex on $\mathbb{R} \setminus \text{supp } \mu$.


\mathcal{R} -Lemma

If $A \in C^{1,\alpha}$ and $\beta \in [0, 1 - \alpha]$, then


$$\mu(B(x, r)) \leq Cr^\beta \quad \Rightarrow \quad \mathcal{R}\mu \in C^{\beta+\alpha}(\mathbb{R}).$$

In particular, $\mathcal{R}\mu \in C^\alpha(\mathbb{R})$.


Potential decomposition



Potential decomposition



Potential decomposition



Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P} \mu + \mathcal{R} \mu$
- $\mathcal{P} \mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R} \mu \in C^\alpha(\mathbb{R})$

Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P} \mu + \mathcal{R} \mu$
- $\mathcal{P} \mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R} \mu \in C^\alpha(\mathbb{R})$

$$\xRightarrow{\text{Missing Lemma}} \mathcal{U}^\Gamma \mu \in C^\alpha([0, 1])$$

Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P}\mu + \mathcal{R}\mu$
- $\mathcal{P}\mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R}\mu \in C^\alpha(\mathbb{R})$

Missing Lemma
 \implies

$$\mathcal{U}^\Gamma \mu \in C^\alpha([0, 1])$$

Step 1
 \nwarrow

$$\mu(B(x, r)) \leq Cr^\alpha$$

Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P}\mu + \mathcal{R}\mu$
- $\mathcal{P}\mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R}\mu \in C^{2\alpha}(\mathbb{R})$

Missing Lemma
 \Longrightarrow

$$\mathcal{U}^\Gamma \mu \in C^\alpha([0, 1])$$

\nwarrow \mathcal{R} -Lemma

\nwarrow Step 1

$$\mu(B(x, r)) \leq Cr^\alpha$$

Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P} \mu + \mathcal{R} \mu$
- $\mathcal{P} \mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R} \mu \in C^{2\alpha}(\mathbb{R})$

Missing Lemma
 \Longrightarrow

$$\mathcal{U}^\Gamma \mu \in C^{2\alpha}([0, 1])$$

\nwarrow \mathcal{R} -Lemma

\nwarrow Step 1

$$\mu(B(x, r)) \leq Cr^\alpha$$

Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P} \mu + \mathcal{R} \mu$
- $\mathcal{P} \mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R} \mu \in C^{2\alpha}(\mathbb{R})$

Missing Lemma
 \Longrightarrow

$$\mathcal{U}^\Gamma \mu \in C^{2\alpha}([0, 1])$$

\nwarrow \mathcal{R} -Lemma

\nwarrow Step 1

$$\mu(B(x, r)) \leq Cr^{2\alpha}$$

Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P} \mu + \mathcal{R} \mu$
- $\mathcal{P} \mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R} \mu \in C^{3\alpha}(\mathbb{R})$

Missing Lemma
 \Longrightarrow

$$\mathcal{U}^\Gamma \mu \in C^{3\alpha}([0, 1])$$

\nwarrow \mathcal{R} -Lemma

\nwarrow Step 1

$$\mu(B(x, r)) \leq Cr^{3\alpha}$$

Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P} \mu + \mathcal{R} \mu$
- $\mathcal{P} \mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R} \mu \in C^{4\alpha}(\mathbb{R})$

Missing Lemma
 \Longrightarrow

$$\mathcal{U}^\Gamma \mu \in C^{4\alpha}([0, 1])$$

\nwarrow \mathcal{R} -Lemma
 \nwarrow

\nwarrow Step 1
 \nwarrow

$$\mu(B(x, r)) \leq Cr^{4\alpha}$$

Bootstrapping scheme

- $\mathcal{U}^\Gamma \mu = \mathcal{P}\mu + \mathcal{R}\mu$
- $\mathcal{P}\mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$
- $\mathcal{R}\mu \in \text{Lip}(\mathbb{R})$

Missing Lemma
 \Longrightarrow

$$\mathcal{U}^\Gamma \mu \in \text{Lip}([0, 1])$$

\nwarrow \mathcal{R} -Lemma

\nwarrow Step 1

$$\mu(B(x, r)) \leq Cr$$


Sums of Hölder and convex functions

An elementary lemma

Suppose that $U : [0, 1] \rightarrow \mathbb{R}$ satisfies $U = P + R$ where

- P is convex on $[0, 1]$
- $R \in C^\beta([0, 1])$
- $U(0) = U(1) = \inf_{t \in [0, 1]} U(t)$.

Then $U \in C^\beta([0, 1])$.




Corollary: the Missing Lemma

Missing Lemma

If μ is the equilibrium measure on Γ , and $\mathcal{U}^\Gamma \mu = \mathcal{P}\mu + \mathcal{R}\mu$ with

- $\mathcal{P}\mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$,
- $\mathcal{R}\mu \in C^\beta(\mathbb{R})$,

then $\mathcal{U}^\Gamma \mu \in C^\beta(\mathbb{R})$.




Corollary: the Missing Lemma

Missing Lemma

If μ is the equilibrium measure on Γ , and $\mathcal{U}^\Gamma \mu = \mathcal{P}\mu + \mathcal{R}\mu$ with

- $\mathcal{P}\mu$ convex on $\mathbb{R} \setminus \text{supp } \mu$,
- $\mathcal{R}\mu \in C^\beta(\mathbb{R})$,

then $\mathcal{U}^\Gamma \mu \in C^\beta(\mathbb{R})$.



Open questions

1-dimensional Lipschitz graphs

Question 1

Suppose that $\Gamma \subset \mathbb{R}^n$ is a 1-dimensional Lipschitz graph. Is the equilibrium measure on Γ absolutely continuous with respect to \mathcal{H}^1 ? What about $\mathcal{H}^1 \ll \mu$?

1-dimensional Lipschitz graphs

Question 1

Suppose that $\Gamma \subset \mathbb{R}^n$ is a 1-dimensional Lipschitz graph. Is the equilibrium measure on Γ absolutely continuous with respect to \mathcal{H}^1 ? What about $\mathcal{H}^1 \ll \mu$?

In our proof:

Step 1. Regularity of $\mathcal{U}^\Gamma \mu \Rightarrow$ regularity of μ ✓

1-dimensional Lipschitz graphs

Question 1

Suppose that $\Gamma \subset \mathbb{R}^n$ is a 1-dimensional Lipschitz graph. Is the equilibrium measure on Γ absolutely continuous with respect to \mathcal{H}^1 ? What about $\mathcal{H}^1 \ll \mu$?

In our proof:

Step 1. Regularity of $\mathcal{U}^\Gamma \mu \Rightarrow$ regularity of μ ✓

Step 2. Prove regularity of $\mathcal{U}^\Gamma \mu$. ✗

Bootstrapping falls apart ☹

Riesz equilibrium measures

Equilibrium measures can be considered for many different energies. One of the classical is the s -Riesz energy:

$$\mathcal{I}_s(\mu) = \iint \frac{1}{|x - y|^s} d\mu(x) d\mu(y).$$

Question 2

Suppose that $\Gamma \subset \mathbb{R}^n$ is a 1-dimensional $C^{1,\alpha}$ graph. If $0 < s < 1$, is the s -equilibrium measure on Γ absolutely continuous with respect to \mathcal{H}^1 ?

Riesz equilibrium measures

Equilibrium measures can be considered for many different energies. One of the classical is the s -Riesz energy:

$$\mathcal{I}_s(\mu) = \iint \frac{1}{|x - y|^s} d\mu(x) d\mu(y).$$

Question 2

Suppose that $\Gamma \subset \mathbb{R}^n$ is a 1-dimensional $C^{1,\alpha}$ graph. If $0 < s < 1$, is the s -equilibrium measure on Γ absolutely continuous with respect to \mathcal{H}^1 ?

In our proof:

Step 1. Regularity of $\mathcal{U}^\Gamma \mu \Rightarrow$ regularity of μ ✗

Unclear how to treat corresponding operators T_β^Γ ☹

Step 2. Prove regularity of $\mathcal{U}^\Gamma \mu$. ✓ (probably)

Higher dimensional surfaces

Just as log-equilibrium measures on curves in \mathbb{R}^2 are classical, $(n - 2)$ -equilibrium measures on $(n - 1)$ -dimensional surfaces in \mathbb{R}^n are classical.

Question 3

Suppose that $\Sigma \subset \mathbb{R}^n$ is a k -dimensional $C^{1,\alpha}$ graph. Is the $(k - 1)$ -equilibrium measure on Σ absolutely continuous with respect to \mathcal{H}^k ? What about s -equilibrium measures for $0 < s < k$?

Higher dimensional surfaces

Just as log-equilibrium measures on curves in \mathbb{R}^2 are classical, $(n - 2)$ -equilibrium measures on $(n - 1)$ -dimensional surfaces in \mathbb{R}^n are classical.

Question 3

Suppose that $\Sigma \subset \mathbb{R}^n$ is a k -dimensional $C^{1,\alpha}$ graph. Is the $(k - 1)$ -equilibrium measure on Σ absolutely continuous with respect to \mathcal{H}^k ? What about s -equilibrium measures for $0 < s < k$?

At least for $s = k - 1$:

Step 1. Regularity of $\mathcal{U}^\Gamma \mu \Rightarrow$ regularity of μ ✓ (probably)

Higher dimensional surfaces

Just as log-equilibrium measures on curves in \mathbb{R}^2 are classical, $(n - 2)$ -equilibrium measures on $(n - 1)$ -dimensional surfaces in \mathbb{R}^n are classical.

Question 3

Suppose that $\Sigma \subset \mathbb{R}^n$ is a k -dimensional $C^{1,\alpha}$ graph. Is the $(k - 1)$ -equilibrium measure on Σ absolutely continuous with respect to \mathcal{H}^k ? What about s -equilibrium measures for $0 < s < k$?

At least for $s = k - 1$:

Step 1. Regularity of $\mathcal{U}^\Gamma \mu \Rightarrow$ regularity of μ ✓ (probably)

Step 2. Prove regularity of $\mathcal{U}^\Gamma \mu$. ✗

Unclear how to get the corresponding elementary lemma on subharmonic functions 😞

Thank you!