



Neurobiology of the major psychoses: a translational perspective on brain structure and function—the FOR2107 consortium

Tilo Kircher^{1,2} · Markus Wöhr^{2,3} · Igor Nenadic^{1,2} · Rainer Schwarting^{2,3} · Gerhard Schratt⁴ · Judith Alferink⁵ · Carsten Culmsee^{2,6} · Holger Garn⁷ · Tim Hahn⁵ · Bertram Müller-Myhsok⁸ · Astrid Dempfle⁹ · Maik Hahmann¹⁰ · Andreas Jansen^{1,2,11} · Petra Pfefferle¹² · Harald Renz⁷ · Marcella Rietschel¹³ · Stephanie H. Witt¹³ · Markus Nöthen¹⁴ · Axel Krug^{1,2} · Udo Dannlowski⁵

Received: 27 June 2018 / Accepted: 19 September 2018 / Published online: 28 September 2018
 © Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Genetic (G) and environmental (E) factors are involved in the etiology and course of the major psychoses (MP), i.e. major depressive disorder (MDD), bipolar disorder (BD), schizoaffective disorder (SZA) and schizophrenia (SZ). The neurobiological correlates by which these predispositions exert their influence on brain structure, function and course of illness are poorly understood. In the FOR2107 consortium, animal models and humans are investigated. A human cohort of MP patients, healthy subjects at genetic and/or environmental risk, and control subjects ($N=2500$) has been established. Participants are followed up after 2 years and twice underwent extensive deep phenotyping (MR imaging, clinical course, neuropsychology, personality, risk/protective factors, biomaterials: blood, stool, urine, hair, saliva). Methods for data reduction, quality assurance for longitudinal MRI data, and (deep) machine learning techniques are employed. In the parallelised animal cluster, genetic risk was introduced by a rodent model (*Cacna1c* deficiency) and its interactions with environmental risk and protective factors are studied. The animals are deeply phenotyped regarding cognition, emotion, and social function, paralleling the variables assessed in humans. A set of innovative experimental projects connect and integrate data from the human and animal parts, investigating the role of microRNA, neuroplasticity, immune signatures, (epi-)genetics and gene expression. Biomaterial from humans and animals are analyzed in parallel. The FOR2107 consortium will delineate pathophysiological entities with common neurobiological underpinnings (“biotypes”) and pave the way for an etiologic understanding of the MP, potentially leading to their prevention, the prediction of individual disease courses, and novel therapies in the future.

Keywords Cohort study · Animal model · Mental disorder · Etiology · Course of illness

Introduction

The major psychoses (MP), i.e. major depressive disorder (MDD), bipolar disorder (BD), schizoaffective disorder (SZA) and schizophrenia (SZ) are common, chronic, costly and debilitating disorders. Their etiology and pathophysiology remain largely elusive. They are caused by a complex interplay of genetic susceptibility and environmental factors. The MP share a familial risk, with genetic correlations

estimated around 0.4–0.7 using molecular genetic methods [1, 2]. Similarly, brain structural studies have reported convergence of effects across the MP [3–6]. Beneficial environmental factors (such as social support) may interact with risk genes in the development of resilience [7, 8]. Little is known about the neurobiological correlates (humans) and causal processes (animal models) of these protective as compared to risk factors.

The mechanisms by which epidemiologically validated risk factors exert their influence on neurobiology, the onset and course of illness remain largely elusive. Understanding these pathways is inevitable for improving prognosis and developing new treatment approaches for modifying or even reversing the detrimental effects of these risks. Hence, there is a pressing need for large-scale, deeply phenotyped, human, longitudinal studies that reach the size of

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00406-018-0943-x>) contains supplementary material, which is available to authorized users.

✉ Tilo Kircher
 kircher2@staff.uni-marburg.de

Extended author information available on the last page of the article

epidemiological cohorts [9, 10]. Functional and structural neuroimaging provides a particularly useful tool for bridging the gap between risk factors and clinical presentations by providing a layer beyond symptoms and behaviour [11].

Animal models must parallel human correlational cohorts for establishing causal mechanisms [8]. Ideally, human and animal studies will mutually inform each other regarding novel mechanisms and targets, which can then be validated further in a human sample or mechanistically explored in animal models.

A number of molecular mechanisms have been discussed in the etiology of the MP [12–14]. Altered immune mechanisms [15–17], dysfunctions in neuroplasticity [18, 19] such as abnormal dendritic arborisation, dendritic spine morphology, and reduced altered neurogenesis [20, 21] among others have been implicated in the MP. In rodents, miRNAs were found to regulate cellular phenotypes associated with MP, including the plasticity of dendrites and spines [22–26], expression of neurotransmitter receptors [27–29], and synaptic plasticity [30–33].

Genome-wide association studies (GWAS) are a powerful approach to identify the genetic underpinnings of disorders. In several independent GWAS the *CACNA1C* gene has been implicated as a major shared risk gene across the MP. *CACNA1C* codes for the $\alpha 1C$ subunit of the voltage-gated L-type calcium channel (LTCC) Cav1.2. It plays a pivotal role in regulating neuronal excitability, synaptic plasticity, and gene expression. Cav1.2 thus represents a primary target for both drugs and second messengers acting on LTCCs [13, 14]. However, the associated variants in *CACNA1C* are found in its intronic region, and the neurobiological mechanisms whereby such variants modify brain structure and function are not well understood [34–38].

In our research consortium briefly outlined here, we have integrated a large, multi-omics, longitudinal cohort study with deeply phenotyped humans and a parallelised animal model to constitute a neurobiologically informed, pathophysiological model for the etiology and the course of major psychoses.

Research programme

The research consortium FOR2107 comprises a human cohort and animal models. All human subjects ($n=2500$) have been deeply phenotyped (see Table 2) and are followed up after 2 years. In complementary animal and cell models, gene by environment (GxE) risk and protective factors on brain structure and function are being mechanistically explored. The clinical (human) and experimental (animal) cluster are tailored to reflect each other regarding selected environmental and genetic risk/protective factors and phenotyping procedures (Fig. 1). They include, among others,

the characterization of brain structure and function, miRNA biogenesis and profiles, immune signatures, (epi-)genetic analyses and gene expression.

The main aims of the FOR2107 are

1. Establishing consequences (humans, animals), respectively, neurobiological mechanisms (animals) of genetic and environmental risk factors for the major psychoses, particularly on the level of brain structure and function in humans and animal models.
2. Uncovering neurobiological determinants for the course of illness (onset, relapse, diagnostic shifts e.g., from MDD to BD, alterations in brain structure and function over time).
3. De novo subgrouping of homogeneous patient groups based on biological, cognitive, and psychopathological longitudinal data (“biotypes”).

Objectives

- (a) Baseline deep phenotyping of $N=2500$, subjects with genetic and/or environmental risk factors, and controls (see Table 1).
- (b) Second deep phenotyping after follow-up of 2 years of all participants (see Table 1). Data are stored in professional database and biobank.
- (c) Provide reliable neurobiological markers for diagnostic switches (e.g., MDD to BD or SZ). Ultimately, a standardized diagnostic toolkit using clinical, neuroimaging and molecular markers could be provided for early detection and specific treatment of potential individual “switchers”.
- (d) Implementation and development of MR imaging standards for large-scale clinical MRI studies in a longitudinal design [39].
- (e) Development and refinement of statistical analysis tools for clustering multi-dimensional and longitudinal data (including machine learning techniques), facilitating the identification of characteristic genotype–phenotype signatures of MP and their courses.
- (f) Acquire information regarding the role of three epigenetic processes (DNA methylation, histone modification, miRNA) in the etiology and clinical course of MP.
- (g) Deep phenotyping of GxE rat models for MP, including risk and protection factors, with correspondence to those in the human cohort.
- (h) Establishment of miRNAs for the longitudinal course of MP, cross-correlation with functional studies in GxE rodent models.
- (i) Analysis of novel immunological markers as biomarkers for the longitudinal course of MP, cross-correlation with functional studies in GxE rodent models.




Fig. 1 Parallel structure of the human and animal clusters. A common genetic risk factors in the human and animal part is (blue boxes) CACNA1C. Common environmental risk factor is maltreatment, protective factor is a beneficial environment [such as social sup-

port (humans)/enrichment (animals)]. Common phenotypes are (red boxes): bonding; social behavior; emotion and cognition. WP work project, CP central project

- (j) Analysis of selected causal effects of GxE risk on brain structure and function in animals.
- (k) Cooperation and biobanking with the national DGPPN cohort, DFG KFO 241; further collaboration with other large-scale imaging studies for validation (BMBF BiDirect, BMBF BIPOLIFE, BMBF PROTECT-AD, DGBS) resting state initiative, SHIP/ SHIP-TREND, and others. Regarding data sharing policy, collaborators submit their suggestions to one of the authors, i.e. a short outline of their project or data analysis strategy. This will be evaluated by the data sharing committee (elected PIs), and if positive, data are transferred to the collaborators.

Work programme

Below we will describe the human (Work Package 1; WP1) and the animal backbone studies (WP2), the microRNA (WP3), the immunological (WP4), (Epi-)genetic (WP5), machine learning, statistics and quality control (WP6) and biobank (CP1) projects. The study and all procedures have been approved by the respective local Ethics committees and by the data protection committees, and are in accordance with the Declaration of Helsinki.

Table 1 Inclusion and exclusion criteria of the FOR2107 study population ($n = 2500$)

Condition	Affective disorders (MDD $n = 700$; BD $n = 300$)	Psychotic disorders (SZ $n = 200$; SZA $n = 100$)	Group			
			Healthy subjects without risks ($n = 1000$)	Healthy subjects, genetic risk ($n = 200$) ^e	Healthy subjects, environmental risk ($n = 200$) ^f	Healthy subjects, environmental and genetic risk ($n = 100$) ^{e,f}
Verbal IQ > 80	+	+	+	+	+	+
West-European ancestry	+	+	+	+	+	+
MRI compatibility ^a	+	+	+	+	+	+
Substance-related disorders	–	– ^c	–	–	–	–
History of severe neurological b or medical disorders ^d	–	–	–	–	–	–
At least one 1st degree relative with life time diagnosis MDD or BD	+	+	– ^e	+	– ^e	+
At least one CTQ- subscale reaching the threshold for maltreatment	+	+	– ^f	– ^f	+	+
Family history of any psychiatric disorder	+	+	–	– ^g	–	– ^g
Current psycho- tropic medication	+	+	–	–	–	–
Current benzodiaz- epine use	– ⁱ	– ⁱ	–	–	–	–
Past or present SCID-diagnosis ^h	+	+	–	–	–	–

^aExclusion due to, e.g., metal implants, pacemakers, claustrophobia, pregnancy, etc.^bE.g., seizures, stroke, multiple sclerosis, dementia, traumatic brain injury^cIn case of SZA and SZ cannabis abuse is tolerated, these patients are included, other comorbid substance-related diagnoses are excluded^dE.g., cancer, auto-immune disorders, chronic inflammatory diseases, and cardio-vascular diseases^eSubjects with at least one relative with the diagnosis of MDD or BD^fAt least one subscale of the CTQ reaching the threshold for maltreatment^gInclude in case of family history (first degree relative) of MDD or BD is inclusion criterion, otherwise exclude^hSCID-confirmed diagnosis of MDD and BD (any polarity); according to DSM-IV criteria (296.xx) or SZ (295.1-3/295.9); and SZA (295.7). Dysthymia and cyclothymia are not inclusion criteria, but are allowed as comorbid disorders within any patient groupⁱWash-out of > 3 half-lives required to be included (within patients only)

Deeply phenotyped human longitudinal cohort (WP1)

Subjects

We have characterized $n = 2500$ individuals across 6 samples with the T1 battery at baseline (first subject in 9/11/2014). We re-evaluate these subjects with the T2 battery after 2 years of follow-up at Marburg and Münster

Universities, Germany. See Table 1 for inclusion and exclusion criteria and Table 2 for acquired data:

1. $n = 700$ patients suffering from MDD.
2. $n = 300$ patients suffering from BD.
3. $n = 200$ patients suffering from SZ.
4. $n = 100$ patients suffering from SZA.
5. $n = 500$ healthy subjects at genetic risk ($n = 200$), environmental risk ($n = 200$) or both ($n = 100$).

Table 2 Data acquired at baseline T1 and after 2 years of follow-up T2

Psychopathology		Baseline	Follow-up
Diagnosis (DSM-IV)	SCID-I, II [1]	X	X
Psychiatric symptoms	OPCRIT 4 [2]	X	X
Personality	SPQ-B [3], NEO-FFI [4]	X	
	TAS-20 [5], ADP-IV [6]		X
Attachment style	RSQ [7]	X	
Subclinical affective symptoms	TEMPS-M [8]		X
Subclinical psychotic symptoms	CAPE [9]		X
Anhedonia	SHAPS-D [10]	X	
Depression level	BDI-II [11], HAM-D [12]	X	X
Suicidality	SBQ-R [13]		X
Anxiety	STAI-T [14]	X	
	STAI-S [14], HAMA [15]	X	X
	PROMIS I-VI/ CROSS-D [16], ASI-3 [17]		X
Positive symptoms	SAPS [18]	X	X
Negative symptoms	SANS [19]	X	X
Mania symptoms	YMRS [20]	X	X
Clinical course	Life chart [21]	X	X
Global/social functioning	GAF [1]	X	X
	SOFAS [22], FAST [23], WHODAS 2.0 [24]		X
Physical activity	IPAQ [25]		X
General symptoms	SCL-90-R [26]	X	X
Lithium response	Alda-Skala [27]		X
MRI			
fMRI	Emotion processing [28]	X	X
	Memory [29]	X	X
	Subliminal face processing [30]	X	X
	Resting state	X	X
sMRI	T1	X	X
	DTI	X	X
General data			
Family history		X	X
Sociodemographics		X	X
Ethnic background		X	X
Employment/salary		X	X
General health	SF-36 [31]	X	X
Smoking	FTQ [32]	X	X
Alcohol consumption	Audit	X	X
Drugs		X	X
Diet	FFQ2 [33]		X
Past and current medication		X	X
Handedness	HQ [34]	X	
Neuropsychology			
Memory	VLMT [35]	X	X
Verbal intelligence (IQ)	MWT-B [36]	X	X
Executive function	TMT-A and -B [37], digit-coding	X	X
Semantic processing	Verbal fluency [38]	X	X
Attention	d2 [39]	X	X
Verbal working memory	Letter–number span [40]	X	X
Visuospatial WM	Block span	X	X

Table 2 (continued)

Psychopathology		Baseline	Follow-up
Risk factors			
Maltreatment	CTQ [41]	X	X
Maltreatment in childhood and youth	ACE [42]	X	
Life events	LEQ [43]	X	X
	THQ [44]	X	
Perceived stress	PSS [45]	X	X
Obstetric complications		X	
Protective factors			
Social support	F-SozU [46]	X	X
Positive life events	Life event-Q [47]	X	
Parental bonding	PBI [48], FEB	X	
Resilience	RS-25 [49]	X	X
Biomaterial			
EDTA blood	Sarstedt tubes EDTA	X	X
PAX gene	PAX-gene tube	X	X
Cell separation	BD Vacutainer CPT 1&2	X	X
Urine	Sterile and DNA-free PE tubes white	X	X
Feces	Sterile and DNA-free stool sample tubes (PE)	X	X
Hair	Sterile and DNA-free Eppendorf tubes	X	X
Heparin blood	Sarstedt tube heparinized	X	X
Serum blood	Sarstedt tube serum	X	X

For references, please see supplementary material

6. $n = 1000$ healthy subjects without any of these risk factors.

MRI scanning

The MRI scanning consists of T1-weighted high-resolution anatomical images, a DTI sequence and four fMRI paradigms: (1) face matching. This paradigm is widely published and robustly elicits amygdala responses to fearful and angry faces (e.g., [40–42]). (2) Episodic memory encoding and retrieval. This face-encoding task induces robust activation of the hippocampal formation and cortical areas during episodic memory processes [43–45]. A retrieval task is presented outside the scanner. (3) Subliminal affective priming. This task uses an event-related subliminal priming design presenting affectively neutral faces from a standardized battery [46–49]. (4) Resting state sequence. All MRI sequences were kept as similar as possible in Marburg and Münster, which was possible due to the same scanner platform (Siemens 3T) [39]. The MRI battery takes less than 60 min scanning time in one session.

Measures for data quality assurance

We have introduced several measures for quality assurance regarding MRI data, human phenotype data, as well as biomaterial processing, storage, and distribution.

MRI data We have implemented a comprehensive quality assurance (QA) protocol for assessing the general quality of the MRI data, to detect potential malfunctions in the scanning equipment, and to evaluate inter-site differences that need to be accounted for in subsequent analyses. The QA protocol is based on the regular measurement of an MRI phantom and an extensive variety of published QA statistics [39]. Human MRI data were assessed including the following procedures: All structural images are visually inspected by trained staff and processed with the SPM CAT12-Toolbox providing ratings for image noise and bias. All gray matter segments undergo the “check data quality using covariance” procedure, which additionally identifies segmentation outliers due to anatomical abnormalities and/or artifacts. All image abnormalities are categorized as (A) exclusion of all participant data (e.g., due to pathologies meeting exclusion criteria a posteriori), (B) exclusion of all MRI data only due to anatomical abnormalities without pathological significance (e.g., abnormally large ventricles; these participants will be followed up but without MRI), (C) exclusion of structural MRI data only due to image artifacts (but will be invited for follow-up MRI). Finally, the “check data quality using covariance” is able to robustly detect dual participation of subjects.

For all fMRI data, movement parameters are extracted, allowing for exclusion of participants due to excessive head

movement for each paradigm. Furthermore, individual activation maps (1st level data) are generated for each subject and paradigm in an automated pipeline showing the “effects of interest” contrast at an exploratory threshold. These images are inspected for “abnormal” activation, e.g., non-activation of the visual cortex, artifacts at tissue borders, etc. All individual MRI quality measures are entered into the Coordination Centre for Clinical Trials (KKS) Marburg database.

Phenotype data Interview ratings are conducted by trained PhD students at both sites according to detailed SOPs. All raters undergo standardized training, and interviews are regularly supervised and videotaped. Videotaped interviews are regularly rated by all other raters, reliability measures are generated and potential mismatches of ratings are discussed and fine-tuned. Problematic cases are discussed at a weekly conference. Currently, the overall intra-class correlations (ICC) across all videotaped interviews are HAMD (ICC = 0.918), HAMA (ICC = 0.959), YMRS (ICC = 0.829), SANS (ICC = 0.845), and SAPS (ICC = 0.923).

Rating and questionnaire sets from the probands are generated as PDF files with an individualized barcode for each subject and are handed out in paper form. These forms are then scanned (using Remark Office OMR software) under quality-controlled conditions based on an annotated CFR (aCRF) (1st quality control) and converted to a database-compatible file structure. Through the data import pipeline, all datasets are checked for plausibility (e.g., range of values, missing data) according to comprehensive aCRFs. For each data import, quality control protocols are generated which gives the data quality manager an immediate and detailed feedback and enables to correct data directly (2nd quality control). By implementing a mirrored test server, this process is repeated until the error rate is zero percent (i.e. error reduction loop). Once this value is reached, data sets are released for export and can be downloaded. Pseudonymization is used for data storage for all subjects. Data storage and supervision of data management according to GCP Guidelines lie with the KKS Marburg.

Biomaterial All biomaterial acquisition is performed by thoroughly trained technical staff, according to SOPs. The quality is constantly monitored by trained staff; the biobank has been certified (<http://www.cbbmr.de>).

Animal backbone study (WP2)

In WP2, we experimentally address the question how *Cacna1c* affects brain structure and function by studying the effects of *Cacna1c* haploinsufficiency in rats. To this aim, we use a newly developed genetic *Cacna1c* rat model and compare wild-type (*Cacna1c*^{+/+}) and constitutive heterozygous

(*Cacna1c*^{+/-}) males and females. Initially, we perform deep and longitudinal behavioral phenotyping in *Cacna1c*^{+/-} rats as compared to *Cacna1c*^{+/+} littermate controls, including both sexes and spanning the period from early postnatal days to adulthood. For behavioral phenotyping of rat models for neuropsychiatric disorders, several readouts with face validity to human core symptoms are available [50]. This includes psychomotor activity (e.g., open field behavior), behavioral despair/passive coping to stress (e.g., immobility in the forced swim test), anxiety (e.g., elevated plus-maze), anhedonia (e.g., sucrose consumption), cognition (e.g., object recognition, attention, risk-taking), and social behavior. Regarding affect and social behavior, rodent ultrasonic vocalizations (USV) have become an important tool, which cannot only be used as readout of positive and negative affect, but also as a means to study a subject's responsiveness to social stimuli, which is of major importance in models of neuropsychiatric disorders, typically characterized by social deficits [51].

Rodents are highly social animals, displaying a rich repertoire of social behaviors and emitting distinct types of USV. Such USV serve as situation-dependent socio-affective signals [51]. In rats, three types of USV are known. Pups emit 40-kHz USV (“distress calls”) when socially isolated from mother and littermates, probably reflecting a negative affective state akin to anxiety. Adult rats emit 50-kHz USV (“rat laughter”) in appetitive situations such as social investigation and play or when exposed to psychostimulants, such as amphetamine, while 22-kHz USV (“alarm calls”) occur in aversive situations such as predator exposure, fear conditioning or social defeat. Measuring USV emission does not only provide a unique tool to gauge affective states in rodents, but they also help to detect changes in affective information processing that are not detectable by conventional behavioral approaches. For instance, we found that maternal immune activation during pregnancy via poly-I:C affects the production of 22-kHz USV emitted during fear conditioning in adult offspring [52]. Poly-I:C administration to pregnant rats during specific gestational periods mimics a viral infection and leads to behavioral impairments in the offspring with relevance to neuropsychiatric disorders in humans. In our fear-conditioning paradigm, poly-I:C exposure during pregnancy caused an increase of 22-kHz USV emission to 300% of saline controls. However, despite this strong effect, a detailed analysis of visible behavior, including freezing, did not reveal any group differences, highlighting the importance of assessing 22-kHz USV as a measure of affective state. Most recently, we have introduced a novel technique to model mania-like elevated mood together with elevated drive. Specifically, we showed that amphetamine treatment leads not only to increased levels of locomotor activity but also to strongly enhanced levels of 50-kHz USV [53] and social responsivity [54], reflecting mania-like elevated mood

and heightened levels of social interest. Notably, amphetamine-induced 50-kHz calling can be blocked by lithium treatment [53]. This allows us to conduct pharmacological studies on the effects of potential anti-manic drugs in *Cacna1c* haploinsufficient rats, as compared to the gold standard lithium. All types of USV serve important communicative functions and induce call-specific behavioral responses in the receiver. While aversive 22-kHz USV induce freezing, indicating an alarming function, appetitive 50-kHz USV induce social approach behavior, suggesting that they serve as social contact calls [55]. These opposite behavioral responses are paralleled by distinct brain activation patterns (22-kHz USV: “fear circuit”, e.g., amygdala; 50-kHz USV: “pleasure circuit”, e.g., nucleus accumbens, where they evoke phasic release of dopamine [56]). Importantly, social approach evoked by playback of pro-social 50-kHz USV can be used as readout for detecting social deficits and is sensitive for aversive experiences during early development, such as post-weaning social isolation [57].

In a second step, we will apply a GxE interaction approach. Specifically, we will analyze whether an established maltreatment, namely post-weaning social isolation, can induce or exaggerate deficits in *Cacna1c* haploinsufficient rats, and whether beneficial environment, namely social and physical enrichment can prevent a deficit. Control subjects are kept in normal group housing. Besides deep and longitudinal behavioral phenotyping, we analyze brain neurochemistry (biogenic amines), hippocampal morphology and plasticity (neurogenesis), as well as neuronal activation patterns (immediate early genes). Together with the other WPs, we further study the role of the immune system and how microRNAs affect brain structure and function in a genotype-dependent manner, comparing *Cacna1c*^{+/-} rats and *Cacna1c*^{+/+} littermate controls exposed to different environmental conditions. In a first study, we showed that post-weaning social isolation leads to a strong increase in hippocampal miRNA-134 expression levels [58]. This effect was paralleled by an increase of an alternative transcript encoding for the ubiquitin ligases Ube3a and Ube3a1. In line with the behavioral alterations evident following post-weaning SI [57, 58], UBE3A duplications are among the most frequent copy number variations associated with social deficits and intellectual disabilities in humans [59]. It is planned to extend our current experimental approach by including additional rat models, e.g., with genetic modifications targeting *ANK3*, *NCAN*, or *TCF4*, in future projects.

miRNA (WP3)

microRNAs (miRNAs), a large family of small non-coding RNAs, have been demonstrated to control synaptic plasticity in hippocampal neurons in vitro and in vivo [60]. Studies suggest an important pro-resilient function of

miRNA-dependent gene regulation in various stress-related diseases, including affective disorders (AD) [61]. However, it is not known how stressful events (e.g., maltreatment) impair miRNA function to promote depressive behaviour.

We pursue four different aims to test the hypothesis that impaired Ca²⁺-dependent miRNA biogenesis at the level of the Dicer1 pre-miRNA processing complex is causally involved in the etiology of endophenotypes of MP in humans. First, we will attempt to characterize modifications of the Dicer1 complex in the rat hippocampus in response to maltreatment. Second, we will try to decipher the signaling pathways that are engaged in neurons of maltreated rats to control miRNA processing activity, with a focus on Ca²⁺-dependent signaling cascades. The biomaterial for these biochemical and molecular biology experiments will be obtained from the rat gene–environment (GxE) cohort run in WP2. Third, in collaboration with WP2, we plan to restore activity of the Dicer1 complex in the hippocampus of maltreated rats by recombinant viral transgene expression and pharmacological treatment. Fourth, in collaboration with WP5, we will perform an integrative analysis of transcriptome and epigenome signatures of miRNA biogenesis factors based on large-scale longitudinal datasets from AD patients with childhood trauma and available data on course of illness. This extensive expression analysis will be complemented by the identification of rare genetic variants in miRNA biogenesis genes using genome-wide genotyping data of AD cohorts that are available through WP5.

In conclusion, our project promises to obtain novel insight regarding the function of altered miRNA activity in AD etiology and course of illness. Hence, miRNA biogenesis genes might emerge as novel candidates for biomarkers and treatment strategies.

Immunology (WP4)

Immune dysregulations represent a longstanding enigma in the pathophysiology of the MP [62]. For example, individuals diagnosed with AD exhibited elevated peripheral blood levels of pro-inflammatory cytokines, e.g., IL-1β, TNF, and IL-6 [63]. Besides that, early life stress such as childhood maltreatment was associated with long-term alterations in immune responses and concomitantly increased vulnerability to AD [63]. Studies in rodents and humans suggested a link between peripheral inflammatory cytokines and perturbed microglial function in the brain that may significantly contribute to depression-associated behaviour [64]. Since the underlying mechanisms of GxE interactions in MP and associated immune alterations are largely unknown, longitudinal studies are needed in humans exposed to GxE interactions and in related animal models to gain novel insights into immune signatures.

In WP4, we perform a longitudinal study in MP patients and in healthy participants of the FOR 2107 cohort, including those with familial genetic risk and exposure to early life stress (childhood maltreatment). We established a cellular large-scale multi-parameter flow cytometry screen for cell-type-specific characterisations of inflammatory responses in MP and immune signatures of specific cell subsets which shall harbor prognostic potential. Further, we investigate whether the altered immune signatures identified in patients are also associated with GxE risk interaction in the *Cacna1c*^{+/-} rat model in conditions of social isolation or enriched environment in the presence or absence of an additional immune stimulus. The rat model and derived microglial cultures allow for mechanistic analyses of interactions between the altered peripheral immune capacity and neuro-inflammatory responses in defined GxE risk settings. Together, this translational approach will provide new insights into immune signatures in subsets of peripheral immune cells in animals and humans exposed to GxE risk factors for MP.

Integrative analyses of genetic, epigenetic, transcriptomic, and environmental vulnerability factors of affective disorders (WP5)

The aim of WP5 is to identify how genetic and environmental factors impact the etiology and course of MP, using genetic, epigenetic and transcriptomic methods. To achieve this aim, systematic genome-wide genotyping of FOR2107 probands is performed. Epigenome-wide methylation and expression profiles are assessed in a subsample at baseline and follow-up. Longitudinal analyses of the methylation and expression profiles are performed in relation to the course of illness or the occurrence of life events. This dataset allows us to further analyze the impact of the cumulative effect of genetic variation on subphenotypes. In addition, bisulfite sequencing analysis of *CACNA1C* is performed longitudinally in part of the sample.

Using data from different platforms, we perform integrative analyses of the genetic, epigenetic and expression data in collaboration with WP1, WP3, WP4 and WP6. This includes multimarker, polygenic and pathway analyses to identify yet undetected genotype–phenotype and GxE effects and the current large GWAS datasets of the PGC. The data are used particularly for in-depth analysis of genes which are interaction partners or in pathways of the risk gene *CACNA1C*. The results of our analyses will elucidate disease-specific factors, as well as factors which are of relevance across diagnostic boundaries.

Analyses of data generated so far have already led to several important findings. For example, besides *CACNA1C*, we also investigated further candidate genes for MP, such as *NCAN*, for which we showed an association with limbic

gray matter deficits in healthy subjects and MDD patients [65]. Using genome-wide genetic data, it was demonstrated that polygenic risk for schizophrenia affects working memory and its neural correlates in healthy subjects [66]. With respect to common mechanisms in affective disorders, we identified shared risk loci and pathways for bipolar disorder and schizophrenia [67]. Results from a genome-wide analysis of microRNA-coding genes in BD suggest that BD-associated microRNAs might be enriched within known microRNA loci [68]. We expect that the wealth of forthcoming longitudinal data will further lead to a large number of important new insights.

Multimodal cross-disorder biotype identification and multi-method data analytic support (WP6)

Diagnosis and treatment in psychiatry still relies almost exclusively on a phenotype-based approach. Although this renders the validity of psychiatric classification questionable and severely hampers biomarker discovery, establishing a system of biologically meaningful groups—so-called biotypes—has remained elusive for decades. Building on the large-scale dataset acquired in the FOR2107, it is now finally possible to apply state-of-the-art tools from the fields of machine learning and multivariate statistics to a rich, multimodal database, thereby bringing robust biotype identification across disorders within reach.

In WP6, we identify and validate biologically plausible, homogeneous biotypes across disorders within and across modalities, enable a de novo subgrouping of homogeneous patient groups based on biological, cognitive, and psychopathological longitudinal data. Drawing on all other WPs, from molecular genetics to whole-brain neuroimaging, we will (1) employ domain-knowledge-based and deep learning-based automatic feature engineering to address the curse of dimensionality, (2) develop a principled approach to confounder removal in linear and non-linear multivariate models, and (3) ensure reliability and internal validity by optimizing cluster solution stability within and across modalities. Crucially, the longitudinal design of this FOR2107 allows us to assess the validity of our cluster solutions based solely on predictive utility. We will thereby directly use clinical relevance as the core criterion for evaluating clustering results. Specifically, predictive biomarker models, such as those designed to predict disease trajectory and outcome, will be trained using our cluster solutions. We expect them to substantially outperform the same models trained on classic disorder groups.

Within this framework, we are uniquely positioned to uncover biologically plausible patient clusters across disorders, which will strengthen the validity of psychiatric classification and crucially simplify biomarker discovery in the future.

Biobanking (CP1)

Structured biomaterial biobanks contribute to translational biomedical research by facilitating quality-controlled acquisition, processing, documentation and dissemination of human bio-specimens. A tailor-made biobank with specimens collected from FOR2107 probands (WP1) was implemented (<http://www.cbbmr.de/da>). Core of the repository is a large collection of blood compounds to allow deep phenotyping of probands by assessment of genetic, epigenetic, immunologic and physiologic parameters. In addition, stool and buccal swabs were collected for further microbiome analyses. We employed quality-controlled workflows which includes core processes such as recruitment, transport, processing, storage and dissemination for 12 different matrices collected within the biobank. Storage of sample data was organized in a secured and for the FOR2107 biobank parametrized section of biobank-information-management system CentraXX (Fig. 2).

Discussion and limitations

The FOR 2107 is an exemplary programme for translational research in mental disorder. Basic neuroscience research findings are applied in a large-scale, deeply phenotyped human longitudinal cohort. Experimental projects allow a multimodal evaluation of pathophysiological and causal etiological processes. The collected data range from behavioral, psychophysiological, neural and (epi)genetic to clinical self-report measures, thus linking various levels of information [see current R-DoC matrix, National Institute of Mental Health (NIMH), 2016] [69].

A particularly important feature is the longitudinal approach of the FOR2107 with a 2-year follow-up of all data domains and an in-depth characterization of clinical course and environmental factors during the follow-up

interval. Large, longitudinal cohorts are largely missing in the field, particularly in neuroimaging research [70], although this is the only way to draw conclusions regarding temporal associations and causal mechanisms of clinical presentations and neurobiological underpinnings, and also for the prediction of clinical trajectories from baseline data. From smaller studies, there is evidence that neuroimaging markers can be related to specific aspects of disease course in major depression [71], and that differences in disease course severity are associated with brain volume changes over time [72].

A further advantage of longitudinal designs is the possibility to investigate to which extent baseline data can be employed to predict clinical outcomes regarding treatment response, relapse, and other aspects of longitudinal clinical presentations. It appears that single markers derived from univariate statistics are inferior compared to multivariate approaches, particularly machine learning techniques as implemented in WP6, e.g., using machine learning on still limited sample sizes, prediction of treatment response to psychotherapy [73] or electroconvulsive treatment [74] has been successfully shown. Furthermore, diagnostic features such as comorbidity [75], or the differentiation of unipolar and bipolar depression [5] have been predicted by machine learning on brain imaging data.

The data acquisition programme is time consuming and demanding for the human participants. This might have led to a sampling bias. As yet, dropout rates during baseline and after the 2-year follow-up are in line with comparable studies. The inclusion of comorbid diagnoses such as anxiety, somatoform disorders, or obsessive–compulsive disorder as a secondary diagnosis is useful to generalize findings. Multiple quality control procedures on all level of data acquisition and analyses minimize heterogeneity of primary data. Diagnoses are established via SCID interviews, done by trained psychologists. This is a high, international research standard and gives reliable results,




Fig. 2 Samples and processes of the biobank

which, however, might not be as valid as if the interviews were done by certified, experienced psychiatrists or psychotherapists.

In sum, the FOR2107 will make a large contribution to the understanding of molecular pathways and neurobiological mechanisms of genetic and environmental risk markers for affective disorders and their course. These results will serve future experiments in disentangling the mechanisms and pathways leading from individual biological risk factors to the clinical presentation of symptoms.

Acknowledgements The FOR 2107 consortium is supported by the German Research Council (Deutsche Forschungsgemeinschaft, DFG, Grant nos. KI 588/14-1, KI 588/14-2, KR 3822/7-1, KR 3822/7-2, NE 2254/1-2, DA 1151/5-1, DA 1151/5-2, SCHW 559/14-1, SCHW 559/14-2, WO 1732/4-1, WO 1732/4-2, AL 1145/5-2, CU 43/9-2, GA 545/7-2, RI 908/11-2, WI 3439/3-2, NO 246/10-2, DE 1614/3-2, HA 7070/2-2, JA 1890/7-1, JA 1890/7-2, MU 1315/8-2, RE 737/20-2, PF 784/1-2, KI 588/17-1, CU 43/9-1). We are deeply indebted to all participants of this study, the recruitment sites and their staff.

Compliance with ethical standards

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.


References

- Bulik-Sullivan BK, Finucane HK, Anttila V, Gusev A, Day FR, Loh P-R, Duncan LE, Perry JRB, Patterson N, Robinson EB, Daly MJ, Price AL, Neale BM (2015) An atlas of genetic correlations across human diseases and traits. *Nat Genet* 47:1236–1241
- Cross-Disorder Group of the Psychiatric Genomics Consortium (2013) Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet* 381:1371–1379
- Chang M, Womer FY, Edmiston EK, Bai C, Zhou Q, Jiang X, Wei S, Wei Y, Ye Y, Huang H, He Y, Xu K, Tang Y, Wang F (2018) Neurobiological commonalities and distinctions among three major psychiatric diagnostic categories: a structural MRI study. *Schizophr Bull* 44:65–74
- Goodkind M, Eickhoff SB, Oathes DJ, Jiang Y, Chang A, Jones-hagata LB, Ortega BN, Zaiko YV, Roach EL, Korgaonkar MS, Grieve SM, Galatzer-Levy IR, Fox PT, Etkin A (2015) Identification of a common neurobiological substrate for mental illness. *JAMA Psychiatry* 5797:305–315
- Redlich R, Almeida JRC, Grotegerd D, Opel N, Kugel H, Heindel W, Arolt V, Phillips ML, Dannlowski U (2014) Brain morphometric biomarkers distinguishing unipolar and bipolar depression. A voxel-based morphometry-pattern classification approach. *JAMA Psychiatry* 71:1222–1230
- Nenadic I, Maitra R, Langbein K, Dietzek M, Lorenz C, Smesny S, Reichenbach JR, Sauer H, Gaser C (2015) Brain structure in schizophrenia vs. psychotic bipolar I disorder: a VBM study. *Schizophr Res* 165:212–219
- Kaufman J, Yang B-Z, Douglas-Palumberi H, Grasso D, Lipschitz D, Houshyar S, Krystal JH, Gelernter J (2006) Brain-derived neurotrophic factor-5-HTTLPR gene interactions and environmental modifiers of depression in children. *Biol Psychiatry* 59:673–680
- Berton O, Hahn C-G, Thase ME (2012) Are we getting closer to valid translational models for major depression? *Science* 338:75–79
- Hyde LW, Bogdan R, Hariri AR (2011) Understanding risk for psychopathology through imaging gene-environment interactions. *Trends Cogn Sci* 15:417–427
- Caspi A, Moffitt TE (2006) Gene-environment interactions in psychiatry: joining forces with neuroscience. *Nat Rev Neurosci* 7:583–590
- Yüksel D, Dietsche B, Forstner AJ, Witt SH, Maier RM, Rietschel M, Konrad C, Nöthen MM, Dannlowski U, Baune BT, Kircher T, Krug A (2017) Polygenic risk for depression and the neural correlates of working memory in healthy subjects. *Prog Neuropsychopharmacol Biol Psychiatry* 79:67–76
- Kupfer DJ, Frank E, Phillips ML (2012) Major depressive disorder: new clinical, neurobiological, and treatment perspectives. *Lancet* 379:1045–1055
- Moylan S, Maes M, Wray NR, Berk M (2013) The neuroprogressive nature of major depressive disorder: pathways to disease evolution and resistance, and therapeutic implications. *Mol Psychiatry* 18:595–606
- Anglin RES, Mazurek MF, Tarnopolsky MA, Rosebush PI (2012) The mitochondrial genome and psychiatric illness. *Am J Med Genet B Neuropsychiatr Genet* 159B:749–759
- Messay B, Lim A, Marsland AL (2012) Current understanding of the bi-directional relationship of major depression with inflammation. *Biol Mood Anxiety Disord* 2:4
- Raison CL, Miller AH (2011) Is depression an inflammatory disorder? *Curr Psychiatry Rep* 13:467–475
- Raedler TJ (2011) Inflammatory mechanisms in major depressive disorder. *Curr Opin Psychiatry* 24:519–525
- Manji HK, Quiroz JA, Sporn J, Payne JL, Denicoff K, Gray NA, Zarate CA, Charney DS (2003) Enhancing neuronal plasticity and cellular resilience to develop novel, improved therapeutics for difficult-to-treat depression. *Biol Psychiatry* 53:707–742
- Zarate CA, Singh J, Manji HK (2006) Cellular plasticity cascades: targets for the development of novel therapeutics for bipolar disorder. *Biol Psychiatry* 59:1006–1020
- Kang HJ, Voleti B, Hajszan T, Rajkowska G, Stockmeier CA, Licznarski P, Lepack A, Majik MS, Jeong LS, Banasr M, Son H, Duman RS (2012) Decreased expression of synapse-related genes and loss of synapses in major depressive disorder. *Nat Med* 18:1413–1417
- Samuels BA, Hen R (2011) Neurogenesis and affective disorders. *Eur J Neurosci* 33:1152–1159
- Schratt GM, Tuebing F, Nigh EA, Kane CG, Sabatini ME, Kiebler M, Greenberg ME (2006) A brain-specific microRNA regulates dendritic spine development. *Nature* 439:283–289
- Impey S, Davare M, Lesiak A, Lasiek A, Fortin D, Ando H, Varlamova O, Obrietan K, Soderling TR, Goodman RH, Wayman GA (2010) An activity-induced microRNA controls dendritic spine formation by regulating Rac1-PAK signaling. *Mol Cell Neurosci* 43:146–156
- Hansen KF, Sakamoto K, Wayman GA, Impey S, Obrietan K (2010) Transgenic miR132 alters neuronal spine density and impairs novel object recognition memory. *PLoS One* 5:e15497
- Lippi G, Steinert JR, Marczylo EL, D'Oro S, Fiore R, Forsythe ID, Schratt GM, Zoli M, Nicotera P, Young KW (2011) Targeting of the Arp3 actin nucleation factor by miR-29a/b regulates dendritic spine morphology. *J Cell Biol* 194:889–904
- Fiore R, Khudayberdiev S, Christensen M, Siegel G, Flavell SW, Kim T-K, Greenberg ME, Schratt GM (2009) Mef2-mediated transcription of the miR379-410 cluster regulates activity-dependent dendritogenesis by fine-tuning Pumilio2 protein levels. *EMBO J* 28:697–710

27. Saba R, Störchel PH, Aksoy-Aksel A, Kepura F, Lippi G, Plant TD, Schratt GM (2012) Dopamine-regulated microRNA MiR-181a controls GluA2 surface expression in hippocampal neurons. *Mol Cell Biol* 32:619–632
28. Edbauer D, Neilson JR, Foster KA, Wang C-F, Seeburg DP, Batterton MN, Tada T, Dolan BM, Sharp PA, Sheng M (2010) Regulation of synaptic structure and function by FMRP-associated microRNAs miR-125b and miR-132. *Neuron* 65:373–384
29. Harraz MM, Eacker SM, Wang X, Dawson TM, Dawson VL (2012) MicroRNA-223 is neuroprotective by targeting glutamate receptors. *Proc Natl Acad Sci USA* 109:18962–18967
30. Gao J, Wang W-Y, Mao Y-W, Gräff J, Guan J-S, Pan L, Mak G, Kim D, Su SC, Tsai L-H (2010) A novel pathway regulates memory and plasticity via SIRT1 and miR-134. *Nature* 466:1105–1109
31. Mellios N, Sugihara H, Castro J, Banerjee A, Le C, Kumar A, Crawford B, Strathmann J, Tropea D, Levine SS, Edbauer D, Sur M (2011) miR-132, an experience-dependent microRNA, is essential for visual cortex plasticity. *Nat Neurosci* 14:1240–1242
32. Cohen JE, Lee PR, Chen S, Li W, Fields RD (2011) MicroRNA regulation of homeostatic synaptic plasticity. *Proc Natl Acad Sci* 108:11650–11655
33. Rajasethupathy P, Fiumara F, Sheridan R, Betel D, Puthanveetil SV, Russo JJ, Sander C, Tuschl T, Kandel E (2009) Characterization of small RNAs in aplysia reveals a role for miR-124 in constraining synaptic plasticity through CREB. *Neuron* 63:803–817
34. Krauthelm JT, Straube B, Dannlowski U, Pyka M, Schneider-Hassloff H, Drexler R, Krug A, Sommer J, Rietschel M, Witt SH, Kircher T (2018) Outgroup emotion processing in the vACC is modulated by childhood trauma and CACNA1C risk variant. *Soc Cogn Affect Neurosci* 13:341–348
35. Dietsche B, Backes H, Laneri D, Weikert T, Witt SH, Rietschel M, Sommer J, Kircher T, Krug A (2014) The impact of a CACNA1C gene polymorphism on learning and hippocampal formation in healthy individuals: a diffusion tensor imaging study. *Neuroimage* 89:256–261
36. Thimm M, Kircher T, Kellermann T, Markov V, Krach S, Jansen A, Zervas K, Eggermann T, Stöcker T, Shah NJ, Nöthen MM, Rietschel M, Witt SH, Mathiak K, Krug A (2011) Effects of a CACNA1C genotype on attention networks in healthy individuals. *Psychol Med* 41:1551–1561
37. Krug A, Nieratschker V, Markov V, Krach S, Jansen A, Zervas K, Eggermann T, Stöcker T, Shah NJ, Treutlein J, Mühleisen TW, Kircher T (2010) Effect of CACNA1C rs1006737 on neural correlates of verbal fluency in healthy individuals. *Neuroimage* 49:1831–1836
38. Paulus FM, Bedenbender J, Krach S, Pyka M, Krug A, Sommer J, Mette M, Nöthen MM, Witt SH, Rietschel M, Kircher T, Jansen A (2014) Association of rs1006737 in CACNA1C with alterations in prefrontal activation and fronto-hippocampal connectivity. *Hum Brain Mapp* 35:1190–1200
39. Vogelbacher C, Möbius TWD, Sommer J, Schuster V, Dannlowski U, Kircher T, Dempfle A, Jansen A, Bopp M (2018) The Marburg-Münster affective disorders cohort study (MACS): a quality assurance protocol for MR neuroimaging data. *Neuroimage* 172:450–460
40. Hariri AR, Mattay VS, Tessitore A, Kolachana BS, Fera F, Goldman D, Egan MF, Weinberger DR (2002) Serotonin transporter genetic variation and the response of the human amygdala. *Science* 297:400–403
41. Pezawas L, Meyer-Lindenberg A, Drabant EM, Verchinski BA, Muñoz KE, Kolachana BS, Egan MF, Mattay VS, Hariri AR, Weinberger DR (2005) 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic susceptibility mechanism for depression. *Nat Neurosci* 8:828–834
42. Dannlowski U, Kugel H, Franke F, Stuhrmann A, Hohoff C, Zwanzger P, Lenzen T, Grotegerd D, Suslow T, Arolt V, Heindel W, Domschke K (2011) Neuropeptide-S (NPS) receptor genotype modulates basolateral amygdala responsiveness to aversive stimuli. *Neuropsychopharmacology* 36:1879–1885
43. Krug A, Krach S, Jansen A, Nieratschker V, Witt SH, Shah NJ, Nöthen MM, Rietschel M, Kircher T (2013) The effect of neurogranin on neural correlates of episodic memory encoding and retrieval. *Schizophr Bull* 39:141–150
44. Krug A, Markov V, Eggermann T, Krach S, Zervas K, Stöcker T, Shah NJ, Schneider F, Nöthen MM, Treutlein J, Rietschel M, Kircher T (2008) Genetic variation in the schizophrenia-risk gene neuregulin1 correlates with differences in frontal brain activation in a working memory task in healthy individuals. *Neuroimage* 42:1569–1576
45. Leube DT, Erb M, Grodd W, Bartels M, Kircher T (2001) Differential activation in parahippocampal and prefrontal cortex during word and face encoding tasks. *Neuroreport* 12:2773–2777
46. Stuhrmann A, Dohm K, Kugel H, Zwanzger P, Redlich R, Grotegerd D, Rauch AV, Arolt V, Heindel W, Suslow T, Zwitterlood P, Dannlowski U (2013) Mood-congruent amygdala responses to subliminally presented facial expressions in major depression: associations with anhedonia. *J Psychiatry Neurosci* 38:249–258
47. Suslow T, Konrad C, Kugel H, Rumstadt D, Zwitterlood P, Schöning S, Ohrmann P, Bauer J, Pyka M, Kersting A, Arolt V, Heindel W, Dannlowski U (2010) Automatic mood-congruent amygdala responses to masked facial expressions in major depression. *Biol Psychiatry* 67:155–160
48. Suslow T, Kugel H, Rauch AV, Dannlowski U, Bauer J, Konrad C, Arolt V, Heindel W, Ohrmann P (2009) Attachment avoidance modulates neural response to masked facial emotion. *Hum Brain Mapp* 30:3553–3562
49. Dannlowski U, Kugel H, Huber F, Stuhrmann A, Redlich R, Grotegerd D, Dohm K, Sehlmeier C, Konrad C, Baune BT, Arolt V, Heindel W, Zwitterlood P, Suslow T (2013) Childhood maltreatment is associated with an automatic negative emotion processing bias in the amygdala. *Hum Brain Mapp* 34:2899–2909
50. Nestler EJ, Hyman SE (2010) Animal models of neuropsychiatric disorders. *Nat Neurosci* 13:1161–1169
51. Wöhr M, Schwarting RKW (2013) Affective communication in rodents: ultrasonic vocalizations as a tool for research on emotion and motivation. *Cell Tissue Res* 354:81–97
52. Yee N, Schwarting RKW, Fuchs E, Wöhr M (2012) Increased affective ultrasonic communication during fear learning in adult male rats exposed to maternal immune activation. *J Psychiatr Res* 46:1199–1205
53. Pereira M, Andreolini R, Schwarting RKW, Brenes JC (2014) Amphetamine-induced appetitive 50-kHz calls in rats: a marker of affect in mania? *Psychopharmacology* 231:2567–2577
54. Engelhardt KA, Fuchs E, Schwarting RKW, Wöhr M (2017) Effects of amphetamine on pro-social ultrasonic communication in juvenile rats: implications for mania models. *Eur Neuropsychopharmacol* 27:261–273
55. Wöhr M, Schwarting RKW (2007) Ultrasonic communication in rats: can playback of 50-kHz calls induce approach behavior? *PLoS One* 2:e1365
56. Willuhn I, Tose A, Wanat MJ, Hart AS, Hollon NG, Phillips PEM, Schwarting RKW, Wöhr M (2014) Phasic dopamine release in the nucleus accumbens in response to pro-social 50 kHz ultrasonic vocalizations in rats. *J Neurosci* 34:10616–10623
57. Seffer D, Rippberger H, Schwarting RKW, Wöhr M (2015) Pro-social 50-kHz ultrasonic communication in rats: post-weaning but not post-adolescent social isolation leads to social impairments-phenotypic rescue by re-socialization. *Front Behav Neurosci* 9:102
58. Valluy J, Bicker S, Aksoy-Aksel A, Lackinger M, Sumer S, Fiore R, Wüst T, Seffer D, Metge F, Dieterich C, Wöhr M, Schwarting RKW, Schratt GM (2015) A coding-independent function of an

- alternative Ube3a transcript during neuronal development. *Nat Neurosci* 18:666–673
59. Glessner JT, Wang K, Cai G, Korvatska O, Kim CE, Wood S, Zhang H, Estes A, Brune CW, Bradfield JP, Imielinski M, Frackelton EC, Reichert J, Crawford EL, Munson J, Sleiman PMA, Chiavacci R, Annaiah K, Thomas K, Hou C, Glaberson W, Flory J, Otieno F, Garriss M, Soorya L, Klei L, Piven J, Meyer KJ, Anagnostou E, Sakurai T, Game RM, Rudd DS, Zurawiecki D, McDougle CJ, Davis LK, Miller J, Posey DJ, Michaels S, Kolevzon A, Silverman JM, Bernier R, Levy SE, Schultz RT, Dawson G, Owley T, McMahon WM, Wassink TH, Sweeney JA, Nurnberger JI, Coon H, Sutcliffe JS, Minshew NJ, Grant SFA, Bucan M, Cook EH, Buxbaum JD, Devlin B, Schellenberg GD, Hakonarson H (2009) Autism genome-wide copy number variation reveals ubiquitin and neuronal genes. *Nature* 459:569–573
60. Schratz GM (2009) microRNAs at the synapse. *Nat Rev Neurosci* 10:842–849
61. Geaghan M, Cairns MJ (2015) MicroRNA and posttranscriptional dysregulation in psychiatry. *Biol Psychiatry* 78:231–239
62. Gibney SM, Drexhage HA (2013) Evidence for a dysregulated immune system in the etiology of psychiatric disorders. *J Neuroimmune Pharmacol* 8:900–920
63. Köhler CA, Freitas TH, Maes M, de Andrade NQ, Liu CS, Fernandes BS, Stubbs B, Solmi M, Veronese N, Herrmann N, Raison CL, Miller BJ, Lancôt KL, Carvalho AF (2017) Peripheral cytokine and chemokine alterations in depression: a meta-analysis of 82 studies. *Acta Psychiatr Scand* 135:373–387
64. Hodes GE, Kana V, Menard C, Merad M, Russo SJ (2015) Neuroimmune mechanisms of depression. *Nat Neurosci* 18:1386–1393
65. Dannlowski U, Kugel H, Grotegerd D, Redlich R, Suchy J, Opel N, Suslow T, Konrad C, Ohrmann P, Bauer J, Kircher T, Krug A, Jansen A, Baune BT, Heindel W, Domschke K, Forstner AJ, Nöthen MM, Treutlein J, Arolt V, Hohoff C, Rietschel M, Witt SH (2015) NCAN cross-disorder risk variant is associated with limbic gray matter deficits in healthy subjects and major depression. *Neuropsychopharmacology* 40:2510–2516
66. Krug A, Dietsche B, Zöllner R, Yüksel D, Nöthen MM, Forstner AJ, Rietschel M, Dannlowski U, Baune BT, Maier RM, Witt SH, Kircher T (2018) Polygenic risk for schizophrenia affects working memory and its neural correlates in healthy subjects. *Schizophr Res*. <https://doi.org/10.1016/j.schres.2018.01.013>
67. Forstner AJ, Hecker J, Hofmann A, Maaser A, Reinbold CS, Mühleisen TW, Leber M, Strohmaier J, Degenhardt FA, Treutlein J, Mattheisen M, Schumacher J, Streit F, Meier S, Herms SL, Hoffmann P, Lacour A, Witt SH, Reif A, Müller-Myhsok B, Lucae S, Maier W, Schwarz MJ, Vedder H, Kammerer-Ciernioch J, Pfennig A, Bauer M, Hautzinger M, Moebus S, Schenk LM, Fischer SB, Sivalingam S, Czerski PM, Hauser J, Lissowska J, Szeszenia-Dabrowska N, Brennan P, McKay JD, Wright A, Mitchell PB, Fullerton JM, Schofield PR, Montgomery GW, Medland SE, Gordon SD, Martin NG, Krasnov V, Chuchalin A, Babadjanova G, Pantelejeva G, Abramova LI, Tiganov AS, Polonikov A, Khusnutdinova E, Alda M, Cruceanu C, Rouleau GA, Turecki G, Laprise C, Rivas F, Mayoral F, Kogevinas M, Grigoriu-Serbanescu M, Becker T, Schulze TG, Rietschel M, Cichon S, Fier H, Nöthen MM (2017) Identification of shared risk loci and pathways for bipolar disorder and schizophrenia. *PLoS One* 12:e0171595
68. Forstner AJ, Hofmann A, Maaser A, Sumer S, Khudayberdiev S, Mühleisen TW, Leber M, Schulze TG, Strohmaier J, Degenhardt FA, Treutlein J, Mattheisen M, Schumacher J, Breuer R, Meier S, Herms SL, Hoffmann P, Lacour A, Witt SH, Reif A, Müller-Myhsok B, Lucae S, Maier W, Schwarz MJ, Vedder H, Kammerer-Ciernioch J, Pfennig A, Bauer M, Hautzinger M, Moebus S, Priebe L, Sivalingam S, Verhaert A, Schulz H, Czerski PM, Hauser J, Lissowska J, Szeszenia-Dabrowska N, Brennan P, McKay JD, Wright A, Mitchell PB, Fullerton JM, Schofield PR, Montgomery GW, Medland SE, Gordon SD, Martin NG, Krasnov V, Chuchalin A, Babadjanova G, Pantelejeva G, Abramova LI, Tiganov AS, Polonikov A, Khusnutdinova E, Alda M, Cruceanu C, Rouleau GA, Turecki G, Laprise C, Rivas F, Mayoral F, Kogevinas M, Grigoriu-Serbanescu M, Propping P, Becker T, Rietschel M, Cichon S, Schratz GM, Nöthen MM (2015) Genome-wide analysis implicates microRNAs and their target genes in the development of bipolar disorder. *Transl Psychiatry* 5:e678
69. Straube B, Reif A, Richter J, Lueken U, Weber H, Arolt V, Jansen A, Zwanzger P, Domschke K, Pauli P, Konrad C, Gerlach AL, Lang T, Fydrich T, Alpers GW, Ströhle A, Wittmann A, Pfeleiderer B, Wittchen H-U, Hamm A, Deckert J, Kircher T (2014) The functional -1019C/G HTR1A polymorphism and mechanisms of fear. *Transl Psychiatry* 4:e490
70. Dohm K, Redlich R, Zwitterlood P, Dannlowski U (2017) Trajectories of major depression disorders: a systematic review of longitudinal neuroimaging findings. *Aust N Z J Psychiatry* 51:441–454
71. Zaremba D, Enneking V, Meinert S, Förster K, Bürger C, Dohm K, Grotegerd D, Redlich R, Dietsche B, Krug A, Kircher T, Kugel H, Heindel W, Arolt V, Dannlowski U (2018) Effects of cumulative illness severity on hippocampal gray matter volume in major depression—a voxel-based morphometry study. *Psychol Med*. <https://doi.org/10.1017/S0033291718000016>
72. Zaremba D, Dohm K, Redlich R, Grotegerd D, Strojny R, Meinert S, Bürger C, Enneking V, Förster K, Repple J, Opel N, Baune BT, Zwitterlood P, Heindel W, Arolt V, Kugel H, Dannlowski U (2018) Association of relapse and cortical changes in patients with major depressive disorder. *JAMA Psychiatry* 75:484–492
73. Hahn T, Kircher T, Straube B, Wittchen H-U, Konrad C, Ströhle A, Wittmann A, Pfeleiderer B, Reif A, Arolt V, Lueken U (2015) Predicting treatment response to cognitive behavioral therapy in panic disorder with agoraphobia by integrating local neural information. *JAMA Psychiatry* 72:68–74
74. Redlich R, Opel N, Grotegerd D, Dohm K, Zaremba D, Bürger C, Munker S, Mühlmann L, Wahl P, Heindel W, Arolt V, Alferink J, Zwanzger P, Zavorotnyy M, Kugel H, Dannlowski U (2016) Prediction of individual response to electroconvulsive therapy via machine learning on structural magnetic resonance imaging data. *JAMA Psychiatry* 73:557–564
75. Lueken U, Straube B, Yang Y, Hahn T, Beesdo-Baum K, Wittchen H-U, Konrad C, Ströhle A, Wittmann A, Gerlach AL, Pfeleiderer B, Arolt V, Kircher T (2015) Separating depressive comorbidity from panic disorder: a combined functional magnetic resonance imaging and machine learning approach. *J Affect Disord* 184:182–192

Affiliations

Tilo Kircher^{1,2}  · **Markus Wöhr^{2,3}** · **Igor Nenadic^{1,2}** · **Rainer Schwarting^{2,3}** · **Gerhard Schratt⁴** · **Judith Alferink⁵** · **Carsten Culmsee^{2,6}** · **Holger Garn⁷** · **Tim Hahn⁵** · **Bertram Müller-Myhsok⁸** · **Astrid Dempfle⁹** · **Maik Hahmann¹⁰** · **Andreas Jansen^{1,2,11}** · **Petra Pfefferle¹²** · **Harald Renz⁷** · **Marcella Rietschel¹³** · **Stephanie H. Witt¹³** · **Markus Nöthen¹⁴** · **Axel Krug^{1,2}** · **Udo Dannlowski⁵**

¹ Department of Psychiatry and Psychotherapy, University of Marburg, Rudolf-Bultmann-Str. 8, 35039 Marburg, Germany

² Centre for Mind, Brain and Behaviour, University of Marburg, Marburg, Germany

³ Department of Experimental and Biological Psychology, University of Marburg, Marburg, Germany

⁴ Department of Neuroscience, ETH Zürich, Zurich, Germany

⁵ Department of Psychiatry and Psychotherapy, University of Münster, Munich, Germany

⁶ Institute for Pharmacology and Clinical Pharmacy, University of Marburg, Marburg, Germany

⁷ Institute of Laboratory Medicine and Pathobiochemistry, University of Marburg, Marburg, Germany

⁸ Max Planck Institute of Psychiatry, Munich, Germany

⁹ Institute of Medical Informatics and Statistics, University of Kiel, Kiel, Germany

¹⁰ Coordinating Centre for Clinical Trials, University of Marburg, Marburg, Germany

¹¹ Core-Facility Brainimaging, Faculty of Medicine, University of Marburg, Marburg, Germany

¹² Comprehensive Biomaterial Bank Marburg, University of Marburg, Marburg, Germany

¹³ Department of Genetic Epidemiology in Psychiatry, Medical Faculty Mannheim, Central Institute of Mental Health, University of Heidelberg, Mannheim, Germany

¹⁴ Department of Genomics, Life and Brain Centre, University of Bonn, Bonn, Germany

Supplementary References for Table 2

- 1 Wittchen H-U, Wunderlich U, Gruschwitz S, Zaudig M. *SKID-I. Strukturiertes Klinisches Interview für DSM-IV*. Hogrefe: Göttingen, 1997.
- 2 McGuffin P, Farmer AE, Harvey I. A polydiagnostic application of operational criteria in studies of psychotic illness. Development and reliability of the OPCRIT system. *Arch Gen Psychiatry* 1991; **48**: 764–70.
- 3 Raine A, Benishay D. The SPQ-B: A Brief Screening Instrument for Schizotypal Personality Disorder. *J Pers Disord* 1995; **9**: 346–355.
- 4 Costa PT, McCrae RR. *Revised NEO Personality Inventory (NEO-PI-RTM) and NEO Five Factor Inventory (NEO-FFI): Professional manual*. Psychological Assessment Resources: Odessa, FL, 1992.
- 5 Bagby RM, Parker JDA, Taylor GJ. The twenty-item Toronto Alexithymia Scale—I. Item selection and cross-validation of the factor structure. *J Psychosom Res* 1994.<http://www.sciencedirect.com/science/article/pii/0022399994900051> (accessed 5 Jun2016).
- 6 Schotte CKW, Doncker D De, Vankerckhoven C, Vertommen H, Cosyns P. Self-report assessment of the DSM-IV personality disorders. Measurement of trait and distress characteristics: the ADP-IV. *Psychol Med* 1998.
- 7 Steffanowski A, Oppl M, Meyerberg J, Schmidt J, Wittmann WW, Nübling R. Psychometrische Überprüfung einer deutschsprachigen Version des Relationship Scales Questionnaire (RSQ). In: M. Bassler (Ed.), *Störungsspezifische Therapieansätze*. In: *Störungsspezifische Therapieansätze in der stationären Psychotherapie*. Psychosozial Verlag: Gießen, 2001, pp 320–342.
- 8 Erfurth A, Gerlach AL, Hellweg I, Boenigk I, Michael N, Akiskal HS. Studies on a German (Münster) version of the temperament auto-questionnaire TEMPS-A: construction and validation of the briefTEMPS-M. *J Affect Disord* 2005; **85**: 53–69.
- 9 Konings M, Bak M, Hanssen M, Os J Van, Krabbendam L. Validity and reliability of the CAPE: a self-report instrument for the measurement of psychotic experiences in the general population. *Acta Psychiatr Scand* 2006.
- 10 Franz M, Lemke MR, Meyer T, Ulferts J, Puhl P, Snaith RP. [German version of the Snaith-Hamilton-Pleasure Scale (SHAPS-D). Anhedonia in schizophrenic and depressive patients]. *Fortschr Neurol Psychiatr* 1998; **66**: 407–13.
- 11 Hautzinger M, Keller F, Kühner C. *BDI-II. Beck Depressions-Inventar. Manual*. Frankfurt,. Harcourt Test Services, 2006.
- 12 Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960; **23**: 56–62.
- 13 Osman A, Bagge CL, Gutierrez PM, Konick LC, Kopper BA, Barrios FX. The Suicidal Behaviors Questionnaire-Revised (SBQ-R): validation with clinical and nonclinical samples. *Assessment* 2001.
- 14 Laux L, Glanzmann P, Schaffner P, Spielberger CD. *Das State-Trait Angstinventar*. Beltz: Weinheim, Germany, 1981.
- 15 Hamilton M. HAMA. Hamilton Anxiety Scale. In: Guy, W. (ed.), *Rockville*. Rockville, 1976.
- 16 Lebeau RT, Glenn DE, Hanover LN, Beesdo-Baum K, Wittchen H-U, Craske MG. A dimensional approach to measuring anxiety for DSM-5. *Int J Methods Psychiatr Res* 2012; **21**: 258–72.
- 17 Taylor S, Zvolensky MJ, Cox BJ, Deacon B, Heimberg RG, Ledley DR *et al*. Robust dimensions of anxiety sensitivity: Development and initial validation of the Anxiety Sensitivity Index-3. *Psychol Assess* 2007; **19**: 176–188.
- 18 Andreasen NC. The Scale for the Assessment of Positive Symptoms (SAPS). *Univ Iowa* 1984.
- 19 Andreasen NC. Scale for the assessment of negative symptoms. *Iowa City Univ Iowa* 1984.
- 20 Young RC, Biggs JT, Ziegler VE, Meyer A. A rating scale for mania: reliability, validity and sensitivity. *Br J Psychiatry* 1978; **133**: 429–35.
- 21 Post RM, Roy-Byrne P, Uhde T. Graphic representation of the life course of illness in patients with affective disorder. *Am J Psychiatry* 1988; **145**: 844–848.
- 22 Morosini P, Magliano L, Brambilla L, Ugolini S, Pioli R. Development, reliability and acceptability of a new version of the DSM-IV Social and Occupational Functioning Assessment Scale (SOFAS) to assess routine social. *Acta Psychiatr Scand* 2000.
- 23 Rosa A, Sánchez-Moreno J, Martínez-Aran A, Salamero M, Torrent C, Reinares M *et al*. Validity and reliability of the Functioning Assessment Short Test (FAST) in bipolar disorder. *Clin Pract Epidemiol Ment Heal* 2007.
- 24 Üstün TB, Chatterji S, Kostanjsek N, Rehm J, Kennedy C, Epping-Jordan J *et al*. Developing the World Health Organization disability assessment schedule 2.0. *Bull World Health Organ* 2010.
- 25 Hagströmer M, Oja P, Sjöström M. The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutr* 2006.
- 26 Derogatis LR, Unger R. Symptom Checklist-90-Revised. In: *The Corsini Encyclopedia of*

Psychology. John Wiley & Sons, Inc.: Hoboken, NJ, USA, 2010
doi:10.1002/9780470479216.corpsy0970.

- 27 Grof P, Duffy A, Cavazzoni P, Grof E, Granham J, MacDougall M *et al.* Is response to prophylactic lithium a familial trait? *J Clin Psychiatry* 2002.
- 28 Dannlowski U, Stuhrmann A, Beutelmann V, Zwanzger P, Lenzen T, Grotegerd D *et al.* Limbic scars: Long-term consequences of childhood maltreatment revealed by functional and structural MRI. *Biol Psychiatry* 2012; **71**: 286–93.
- 29 Krug A, Krach S, Jansen A, Nieratschker V, Witt SH, Shah NJ *et al.* The Effect of Neurogranin on Neural Correlates of Episodic Memory Encoding and Retrieval. *Schizophr Bull* 2013; **39**: 141–50.
- 30 Stuhrmann A, Dohm K, Kugel H, Zwanzger P, Redlich R, Grotegerd D *et al.* Mood-congruent amygdala responses to subliminally presented facial expressions in Major Depression: Associations with anhedonia. *J Psychiatry Neurosci* 2013; **38**: 249–58.
- 31 Bullinger M. Erfassung der gesundheitsbezogenen Lebensqualität mit dem SF-36-Health Survey. *Bundesgesundheitsblatt - Gesundheitsforsch - Gesundheitsschutz* 2000; **43**: 190–197.
- 32 Heatherton TF, Kozlowski LT, Frecker RC, Fagerström KO. The Fagerström Test for Nicotine Dependence: a revision of the Fagerström Tolerance Questionnaire. *Br J Addict* 1991; **86**: 1119–27.
- 33 von Ruesten A, Feller S, Bergmann MM, Boeing H. Diet and risk of chronic diseases: results from the first 8 years of follow-up in the EPIC-Potsdam study. *Eur J Clin Nutr* 2013; **67**: 412–9.
- 34 Raczkowski D, Kalat JW, Nebes R. Reliability and validity of some handedness questionnaire items. *Neuropsychologia* 1974; **12**: 43–7.
- 35 Heubrock D. Der Auditiv-Verbale Lerntest (AVLT) in der klinischen und experimentellen Neuropsychologie. Durchführung, Auswertung und Forschungsergebnisse. *Z Diff Diagn Psychol* 1992; **3**: 161–174.
- 36 Lehrl S. *Mehrfachwahl-Wortschatz-Intelligenztest MWT-B*. Hogrefe: Göttingen, 1995.
- 37 Reitan RM. *Trail Making Test. Manual for administration and scoring*. Reitan Neuropsychology Laboratory: Tuscon, USA, 1979.
- 38 Daum I, Gräber S, Schugens MM, Mayes AR. Memory dysfunction of the frontal type in normal ageing. *Neuroreport* 1996; **7**: 2625–8.
- 39 Brickenkamp R. *Der Aufmerksamkeits-Belastungstest d2*. Hogrefe: Göttingen, 2002.
- 40 Gold JM, Carpenter C, Randolph C, Goldberg TE, Weinberger DR. Auditory working memory and Wisconsin Card Sorting Test performance in schizophrenia. *Arch Gen Psychiatry* 1997; **54**: 159–65.
- 41 Wingenfeld K, Spitzer C, Mensebach C, Grabe HJ, Hill AB, Gast U *et al.* Die deutsche Version des Childhood Trauma Questionnaire (CTQ): Erste Befunde zu den psychometrischen Kennwerten The German Version of the Childhood Trauma Questionnaire (CTQ): Preliminary Psychometric Properties. *thieme-connect.com* 2010; : 442–450.
- 42 Wingenfeld K, Schäfer I, Terfehr K, Grabski H, Driessen M, Grabe HJ *et al.* [The reliable, valid and economic assessment of early traumatization: first psychometric characteristics of the German version of the Adverse Childhood Experiences Questionnaire (ACE)]. *Psychother Psychosom Med Psychol* 2011; **61**: e10-4.
- 43 Norbeck JS. Modification of life event questionnaires for use with female respondents. *Res Nurs Health* 1984; **7**: 61–71.
- 44 Maercker A. *Deutsche Übersetzung des Trauma History Questionnaire*. Unveröffentlichtes Manuskript, Universität Zürich, 2002.
- 45 Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983; **24**: 385–96.
- 46 Fydrich T, Sommer G, Brähler E. *F-SozU. Fragebogen zur Sozialen Unterstützung*. Hogrefe: Göttingen, 2007.
- 47 Kuepper Y, Wielpuetz C, Alexander N, Mueller E, Grant P, Hennig J. 5-HTTLPR S-allele: a genetic plasticity factor regarding the effects of life events on personality? *Genes Brain Behav* 2012. doi:10.1111/j.1601-183X.2012.00783.x.
- 48 Burbach DJ, Kashani JH, Rosenberg TK. Parental bonding and depressive disorders in adolescents. *J Child Psychol Psychiatry* 1989; **30**: 417–29.
- 49 Leppert K, Gunzelmann T, Schumacher J, Strauss B, Brähler E. [Resilience as a protective personality characteristic in the elderly]. *Psychother Psychosom Med Psychol* 2005; **55**: 365–9.