

The multiplicities of zeros of $\zeta(s)$ and its values over short intervals

Aleksandar Ivić

Serbian Academy of Arts and Sciences, Belgrade

A. A. Karatsuba's 80th Birthday Conference in Number Theory
and Applications

22-27.05.2015, Moscow, Russia

Professor A.A. Karatsuba (1937 – 2008), photo from Oberwolfach (1993), taken by Yoichi Motohashi



Let $r = m(\rho)$ (≥ 1) denote the **multiplicity** of the complex zero $\rho = \beta + i\gamma$ of the Riemann zeta-function $\zeta(s)$. A zero ρ is **simple** if $m(\rho) = 1$. One may assume $\frac{1}{2} \leq \beta < 1, \gamma > 0$ since $\zeta(s) \neq 0$ for $\Re s \geq 1$, $\overline{\zeta(s)} = \zeta(\bar{s})$ and $\zeta(s) = \chi(s)\zeta(1-s)$, $\chi(s) = \frac{\Gamma(\frac{1}{2}(1-s))}{\Gamma(\frac{1}{2}s)}\pi^{s-1/2}$ (the **functional equation**).

Let $r = m(\rho)$ (≥ 1) denote the **multiplicity** of the complex zero $\rho = \beta + i\gamma$ of the Riemann zeta-function $\zeta(s)$. A zero ρ is **simple** if $m(\rho) = 1$. One may assume $\frac{1}{2} \leq \beta < 1, \gamma > 0$ since $\zeta(s) \neq 0$ for $\Re s \geq 1$, $\overline{\zeta(s)} = \zeta(\bar{s})$ and $\zeta(s) = \chi(s)\zeta(1-s)$, $\chi(s) = \frac{\Gamma(\frac{1}{2}(1-s))}{\Gamma(\frac{1}{2}s)}\pi^{s-1/2}$ (the **functional equation**).

This means that $\zeta(\rho) = \zeta'(\rho) = \dots = \zeta^{(r-1)}(\rho) = 0$, but $\zeta^{(r)}(\rho) \neq 0$. It implies that $\zeta(s + \rho)s^{-r}$ is **regular** in a neighborhood of the point $s = 0$.

Let $r = m(\rho)$ (≥ 1) denote the **multiplicity** of the complex zero $\rho = \beta + i\gamma$ of the Riemann zeta-function $\zeta(s)$. A zero ρ is **simple** if $m(\rho) = 1$. One may assume $\frac{1}{2} \leq \beta < 1, \gamma > 0$ since $\zeta(s) \neq 0$ for $\Re s \geq 1$, $\zeta(\bar{s}) = \overline{\zeta(s)}$ and $\zeta(s) = \chi(s)\zeta(1-s)$, $\chi(s) = \frac{\Gamma(\frac{1}{2}(1-s))}{\Gamma(\frac{1}{2}s)}\pi^{s-1/2}$ (the **functional equation**).

This means that $\zeta(\rho) = \zeta'(\rho) = \dots = \zeta^{(r-1)}(\rho) = 0$, but $\zeta^{(r)}(\rho) \neq 0$. It implies that $\zeta(s + \rho)s^{-r}$ is **regular** in a neighborhood of the point $s = 0$.

A very strong conjecture is **that all the zeros ρ are simple**, and this is true for **all known zeros**. The conjecture seems to be independent of the famous, **yet unproved Riemann Hypothesis (RH)**, that $\Re \rho = 1/2$ ($\forall \rho$).

Let $r = m(\rho)$ (≥ 1) denote the **multiplicity** of the complex zero $\rho = \beta + i\gamma$ of the Riemann zeta-function $\zeta(s)$. A zero ρ is **simple** if $m(\rho) = 1$. One may assume $\frac{1}{2} \leq \beta < 1, \gamma > 0$ since $\zeta(s) \neq 0$ for $\Re s \geq 1$, $\zeta(s) = \zeta(\bar{s})$ and $\zeta(s) = \chi(s)\zeta(1-s)$, $\chi(s) = \frac{\Gamma(\frac{1}{2}(1-s))}{\Gamma(\frac{1}{2}s)}\pi^{s-1/2}$ (the **functional equation**).

This means that $\zeta(\rho) = \zeta'(\rho) = \dots = \zeta^{(r-1)}(\rho) = 0$, but $\zeta^{(r)}(\rho) \neq 0$. It implies that $\zeta(s + \rho)s^{-r}$ is **regular** in a neighborhood of the point $s = 0$.

A very strong conjecture is **that all the zeros ρ are simple**, and this is true for **all known zeros**. The conjecture seems to be independent of the famous, **yet unproved Riemann Hypothesis (RH)**, that $\Re \rho = 1/2$ ($\forall \rho$).

Namely the simplicity of zeros and the RH seem to be two statements **independent of one another**. Both could be **true**, or **false**, or **one true** and the **other one false**.

Besides the **strongest possible conjecture** that all complex zeros $\rho = \beta + i\gamma$ are simple, **A.A. Karatsuba** (2006) mentions **two somewhat weaker conjectures**:

Besides the **strongest possible conjecture** that all complex zeros $\rho = \beta + i\gamma$ are simple, **A.A. Karatsuba** (2006) mentions **two somewhat weaker conjectures**:

$$m(\rho) \ll 1 \quad (\forall \rho),$$

meaning that all multiplicities are **bounded** by some constant, and

Besides the **strongest possible conjecture** that all complex zeros $\rho = \beta + i\gamma$ are simple, **A.A. Karatsuba** (2006) mentions **two somewhat weaker conjectures**:

$$m(\rho) \ll 1 \quad (\forall \rho),$$

meaning that all multiplicities are **bounded** by some constant, and

$$m(\rho) \text{ is } \mathbf{unbounded} \text{ as } |\gamma| \rightarrow \infty.$$

Besides the **strongest possible conjecture** that all complex zeros $\rho = \beta + i\gamma$ are simple, **A.A. Karatsuba** (2006) mentions **two somewhat weaker conjectures**:

$$m(\rho) \ll 1 \quad (\forall \rho),$$

meaning that all multiplicities are **bounded** by some constant, and

$$m(\rho) \text{ is } \mathbf{unbounded} \text{ as } |\gamma| \rightarrow \infty.$$

He also says that **the universality of $\zeta(s)$** (**S.M. Voronin**, 1975) **should include the last conjecture**, but that all these **“are merely surmises”**.

Zeta zeros with **large multiplicities**, statistically speaking, are **rare**. **A. Fujii** (1981) proved that

$$N_j(T) \leq C_1 N(T) e^{-C_j} \quad (j \geq 1; C, C_1 > 0; T \geq T_0 > 0).$$

Zeta zeros with **large multiplicities**, statistically speaking, are **rare**. **A. Fujii** (1981) proved that

$$N_j(T) \leq C_1 N(T) e^{-C_j} \quad (j \geq 1; C, C_1 > 0; T \geq T_0 > 0).$$

$N(T)$ denotes the number of complex zeros ρ of $\zeta(s)$ with $0 < \Im \rho \leq T$ (multiplicities counted), while $N_j(T)$ denotes those zeros counted by $N(T)$ whose multiplicities are j , where j is **not necessarily fixed**.

Zeta zeros with **large multiplicities**, statistically speaking, are **rare**. **A. Fujii** (1981) proved that

$$N_j(T) \leq C_1 N(T) e^{-C_j} \quad (j \geq 1; C, C_1 > 0; T \geq T_0 > 0).$$

$N(T)$ denotes the number of complex zeros ρ of $\zeta(s)$ with $0 < \Im \rho \leq T$ (multiplicities counted), while $N_j(T)$ denotes those zeros counted by $N(T)$ whose multiplicities are j , where j is **not necessarily fixed**.

M. Korolev (2006) obtained **explicit numerical values** for the constants C, C_1 .

Zeta zeros with **large multiplicities**, statistically speaking, are **rare**. **A. Fujii** (1981) proved that

$$N_j(T) \leq C_1 N(T) e^{-C_j} \quad (j \geq 1; C, C_1 > 0; T \geq T_0 > 0).$$

$N(T)$ denotes the number of complex zeros ρ of $\zeta(s)$ with $0 < \Im \rho \leq T$ (multiplicities counted), while $N_j(T)$ denotes those zeros counted by $N(T)$ whose multiplicities are j , where j is **not necessarily fixed**.

M. Korolev (2006) obtained **explicit numerical values** for the constants C, C_1 .

It seems plausible that, for any given $j \geq 2$, **almost all zeros are simple**, namely

$$N_j(T) = o(N(T)) \quad (T \rightarrow \infty).$$

It follows when $j \rightarrow \infty$, but, in general, this is **not known yet**.

This relation certainly **cannot hold** for $j = 1$.

This relation certainly **cannot hold** for $j = 1$.

Namely **D.R. Heath-Brown** (1979) showed unconditionally, by modifying **A. Selberg**'s classical zero-detection method (1942), that

This relation certainly **cannot hold** for $j = 1$.

Namely **D.R. Heath-Brown** (1979) showed unconditionally, by modifying **A. Selberg**'s classical zero-detection method (1942), that

$$N_1(T) \gg N(T).$$

This relation certainly **cannot hold** for $j = 1$.

Namely **D.R. Heath-Brown** (1979) showed unconditionally, by modifying **A. Selberg**'s classical zero-detection method (1942), that

$$N_1(T) \gg N(T).$$

The value of the \gg constant is at least 0.34. In fact, his proof shows that

$$N_s(T) \gg N(T).$$

This relation certainly **cannot hold** for $j = 1$.

Namely **D.R. Heath-Brown** (1979) showed unconditionally, by modifying **A. Selberg**'s classical zero-detection method (1942), that

$$N_1(T) \gg N(T).$$

The value of the \gg constant is at least 0.34. In fact, his proof shows that

$$N_s(T) \gg N(T).$$

Selberg was the first to show that **a positive proportion of zeros** lies on the critical line $\Re s = \frac{1}{2}$.

This relation certainly **cannot hold** for $j = 1$.

Namely **D.R. Heath-Brown** (1979) showed unconditionally, by modifying **A. Selberg**'s classical zero-detection method (1942), that

$$N_1(T) \gg N(T).$$

The value of the \gg constant is at least 0.34. In fact, his proof shows that

$$N_s(T) \gg N(T).$$

Selberg was the first to show that **a positive proportion of zeros** lies on the critical line $\Re s = \frac{1}{2}$.

Here $N_s(T)$ denotes the number of **simple zeta-zeros** of the form $\rho = 1/2 + i\gamma$, which are counted by $N(T)$.

There is a connection between **multiplicities of zeta-zeros** and the integral of $\zeta(s)$ over “**very short intervals**”, namely **lower bounds** of the form

$$(1) \quad \int_{\delta}^{2\delta} |\zeta(\beta + i\gamma + i\alpha)|^k d\alpha \geq \ell = \ell(\gamma, \delta, k) \quad (0 < \delta < \tfrac{1}{4}, k \in \mathbb{N}, \gamma \geq \gamma_0 > 0).$$

There is a connection between **multiplicities of zeta-zeros** and the integral of $\zeta(s)$ over “**very short intervals**”, namely **lower bounds** of the form

$$(1) \quad \int_{\delta}^{2\delta} |\zeta(\beta + i\gamma + i\alpha)|^k d\alpha \geq \ell = \ell(\gamma, \delta, k) \quad (0 < \delta < \tfrac{1}{4}, k \in \mathbb{N}, \gamma \geq \gamma_0 > 0).$$

For fixed β such that $\beta \geq \frac{1}{2}$, let \mathcal{D} be the rectangle with vertices

$$\tfrac{1}{4} - \beta \pm i \log^2 \gamma, 2 \pm i \log^2 \gamma, \zeta(\rho) = 0, \rho = \beta + i\gamma (\gamma \geq \gamma_0 > 0),$$

There is a connection between **multiplicities of zeta-zeros** and the integral of $\zeta(s)$ over “**very short intervals**”, namely **lower bounds** of the form

$$(1) \quad \int_{\delta}^{2\delta} |\zeta(\beta + i\gamma + i\alpha)|^k d\alpha \geq \ell = \ell(\gamma, \delta, k) \quad (0 < \delta < \tfrac{1}{4}, k \in \mathbb{N}, \gamma \geq \gamma_0 > 0).$$

For fixed β such that $\beta \geq \frac{1}{2}$, let \mathcal{D} be the rectangle with vertices

$$\tfrac{1}{4} - \beta \pm i \log^2 \gamma, 2 \pm i \log^2 \gamma, \zeta(\rho) = 0, \rho = \beta + i\gamma (\gamma \geq \gamma_0 > 0),$$

and let α be a parameter for which $0 < \alpha \leq 1$. By the **residue theorem**

There is a connection between **multiplicities of zeta-zeros** and the integral of $\zeta(s)$ over “**very short intervals**”, namely **lower bounds** of the form

$$(1) \quad \int_{\delta}^{\delta+2\delta} |\zeta(\beta+i\gamma+i\alpha)|^k d\alpha \geq \ell = \ell(\gamma, \delta, k) \quad (0 < \delta < \tfrac{1}{4}, k \in \mathbb{N}, \gamma \geq \gamma_0 > 0).$$

For fixed β such that $\beta \geq \frac{1}{2}$, let \mathcal{D} be the rectangle with vertices

$$\tfrac{1}{4} - \beta \pm i \log^2 \gamma, 2 \pm i \log^2 \gamma, \zeta(\rho) = 0, \rho = \beta + i\gamma (\gamma \geq \gamma_0 > 0),$$

and let α be a parameter for which $0 < \alpha \leq 1$. By the **residue theorem**

$$\frac{\zeta(\beta + i\gamma + i\alpha)}{(i\alpha)^r} = \frac{1}{2\pi i} \int_{\mathcal{D}} \Gamma(s - i\alpha) \frac{\zeta(s + \rho)}{s^r} ds.$$

There is a connection between **multiplicities of zeta-zeros** and the integral of $\zeta(s)$ over “**very short intervals**”, namely **lower bounds** of the form

$$(1) \quad \int_{\delta}^{2\delta} |\zeta(\beta + i\gamma + i\alpha)|^k d\alpha \geq \ell = \ell(\gamma, \delta, k) \quad (0 < \delta < \tfrac{1}{4}, k \in \mathbb{N}, \gamma \geq \gamma_0 > 0).$$

For fixed β such that $\beta \geq \frac{1}{2}$, let \mathcal{D} be the rectangle with vertices

$$\tfrac{1}{4} - \beta \pm i \log^2 \gamma, 2 \pm i \log^2 \gamma, \zeta(\rho) = 0, \rho = \beta + i\gamma (\gamma \geq \gamma_0 > 0),$$

and let α be a parameter for which $0 < \alpha \leq 1$. By the **residue theorem**

$$\frac{\zeta(\beta + i\gamma + i\alpha)}{(i\alpha)^r} = \frac{1}{2\pi i} \int_{\mathcal{D}} \Gamma(s - i\alpha) \frac{\zeta(s + \rho)}{s^r} ds.$$

Key fact: Since ρ is a zero of $\zeta(s)$ of multiplicity r , then the function $\zeta(s + \rho)s^{-r}$ is **regular** at $s = 0$. Its **only pole in \mathcal{D}** is $s = i\alpha$.

This gives

$$\zeta(\beta + i\gamma + i\alpha) \ll \alpha^r \left(\gamma(\beta - \tfrac{1}{4})^{-r} + 2^{-r} \right) \ll \alpha^r \gamma(\beta - \tfrac{1}{4})^{-r}.$$

This gives

$$\zeta(\beta + i\gamma + i\alpha) \ll \alpha^r \left(\gamma(\beta - \tfrac{1}{4})^{-r} + 2^{-r} \right) \ll \alpha^r \gamma(\beta - \tfrac{1}{4})^{-r}.$$

Integrating over α from δ to 2δ , and taking logarithms we obtain

This gives

$$\zeta(\beta + i\gamma + i\alpha) \ll \alpha^r (\gamma(\beta - \tfrac{1}{4})^{-r} + 2^{-r}) \ll \alpha^r \gamma(\beta - \tfrac{1}{4})^{-r}.$$

Integrating over α from δ to 2δ , and taking logarithms we obtain

THEOREM 1. *If $\beta \geq \frac{1}{2}$, $\gamma > \gamma_0 > 0$, $0 < \delta < 1/8$, $k \in \mathbb{N}$, then with the notation introduced above we have*

$$(2) \quad m(\beta + i\gamma) = r \leq \frac{1}{\log(\frac{1}{8\delta})} \left(\log \gamma - \frac{1}{k} \log \ell + O(1) \right) + O(1).$$

This gives

$$\zeta(\beta + i\gamma + i\alpha) \ll \alpha^r (\gamma(\beta - \tfrac{1}{4})^{-r} + 2^{-r}) \ll \alpha^r \gamma(\beta - \tfrac{1}{4})^{-r}.$$

Integrating over α from δ to 2δ , and taking logarithms we obtain

THEOREM 1. *If $\beta \geq \frac{1}{2}$, $\gamma > \gamma_0 > 0$, $0 < \delta < 1/8$, $k \in \mathbb{N}$, then with the notation introduced above we have*

$$(2) \quad m(\beta + i\gamma) = r \leq \frac{1}{\log(\frac{1}{8\delta})} \left(\log \gamma - \frac{1}{k} \log \ell + O(1) \right) + O(1).$$

We would like to **let $\delta \rightarrow 0+$** in (2) and obtain

$$(3) \quad m(\beta + i\gamma) = o(\log \gamma) \quad (\beta \geq \tfrac{1}{2}, \gamma \rightarrow \infty),$$

which is **not yet known unconditionally** in the **general case**, namely for the whole range $\frac{1}{2} \leq \beta < 1$.

This gives

$$\zeta(\beta + i\gamma + i\alpha) \ll \alpha^r (\gamma(\beta - \tfrac{1}{4})^{-r} + 2^{-r}) \ll \alpha^r \gamma(\beta - \tfrac{1}{4})^{-r}.$$

Integrating over α from δ to 2δ , and taking logarithms we obtain

THEOREM 1. *If $\beta \geq \frac{1}{2}$, $\gamma > \gamma_0 > 0$, $0 < \delta < 1/8$, $k \in \mathbb{N}$, then with the notation introduced above we have*

$$(2) \quad m(\beta + i\gamma) = r \leq \frac{1}{\log(\frac{1}{8\delta})} \left(\log \gamma - \frac{1}{k} \log \ell + O(1) \right) + O(1).$$

We would like to **let $\delta \rightarrow 0+$** in (2) and obtain

$$(3) \quad m(\beta + i\gamma) = o(\log \gamma) \quad (\beta \geq \tfrac{1}{2}, \gamma \rightarrow \infty),$$

which is **not yet known unconditionally** in the **general case**, namely for the whole range $\frac{1}{2} \leq \beta < 1$.

Obtaining (3) from (2) (or in any other way!) **seems very difficult**.

On the **Lindelöf Hypothesis** (that $\zeta(\frac{1}{2} + it) \ll_{\varepsilon} |t|^{\varepsilon}$) (3) does hold. Here and later ε denotes arbitrarily small positive constants, not necessarily the same ones at each occurrence.

On the **Lindelöf Hypothesis** (that $\zeta(\frac{1}{2} + it) \ll_{\varepsilon} |t|^{\varepsilon}$) (3) does hold. Here and later ε denotes arbitrarily small positive constants, not necessarily the same ones at each occurrence.

It is well known that **RH implies**

$$\zeta(\tfrac{1}{2} + it) \ll \exp\left(C \frac{\log t}{\log \log t}\right) \quad (C > 0).$$

On the **Lindelöf Hypothesis** (that $\zeta(\frac{1}{2} + it) \ll_{\varepsilon} |t|^{\varepsilon}$) (3) does hold. Here and later ε denotes arbitrarily small positive constants, not necessarily the same ones at each occurrence.

It is well known that **RH implies**

$$\zeta(\tfrac{1}{2} + it) \ll \exp\left(C \frac{\log t}{\log \log t}\right) \quad (C > 0).$$

Thus the **RH implies the LH**. It is **not known** whether the converse is true!

On the **Lindelöf Hypothesis** (that $\zeta(\frac{1}{2} + it) \ll_{\varepsilon} |t|^{\varepsilon}$) (3) does hold. Here and later ε denotes arbitrarily small positive constants, not necessarily the same ones at each occurrence.

It is well known that **RH implies**

$$\zeta(\tfrac{1}{2} + it) \ll \exp\left(C \frac{\log t}{\log \log t}\right) \quad (C > 0).$$

Thus the **RH implies the LH**. It is **not known** whether the converse is true!

On the RH one has a **small improvement** of (3), namely

$$(4) \quad m(\beta + i\gamma) \ll \frac{\log \gamma}{\log \log \gamma}.$$

On the **Lindelöf Hypothesis** (that $\zeta(\frac{1}{2} + it) \ll_{\varepsilon} |t|^{\varepsilon}$) (3) does hold. Here and later ε denotes arbitrarily small positive constants, not necessarily the same ones at each occurrence.

It is well known that **RH implies**

$$\zeta(\tfrac{1}{2} + it) \ll \exp\left(C \frac{\log t}{\log \log t}\right) \quad (C > 0).$$

Thus the **RH implies the LH**. It is **not known** whether the converse is true!

On the **RH** one has a **small improvement** of (3), namely

$$(4) \quad m(\beta + i\gamma) \ll \frac{\log \gamma}{\log \log \gamma}.$$

H.L. Montgomery (1977) proved (on RH) that **at least 2/3 of the zeros ρ are simple**, while **H.M. Bui** and **D.R. Heath-Brown** (2013) improved (also on RH) the constant 2/3 to $19/27 = 0.\overline{703}$.

The classical **Riemann–von Mangoldt** formula says that

$$N(T) = \frac{T}{2\pi} \log \left(\frac{T}{2\pi} \right) - \frac{T}{2\pi} + \frac{7}{8} + S(T) + O\left(\frac{1}{T}\right).$$

The classical **Riemann–von Mangoldt** formula says that

$$N(T) = \frac{T}{2\pi} \log \left(\frac{T}{2\pi} \right) - \frac{T}{2\pi} + \frac{7}{8} + S(T) + O\left(\frac{1}{T}\right).$$

Here $N(T)$ is the number of zeta zeros with $0 < \gamma \leq T$ and we have $S(T) = \frac{1}{\pi} \arg \zeta(\frac{1}{2} + iT)$. This function **regulates the finer behavior** of $\zeta(s)$. The term $O(1/T)$ is a **smooth function** of T .

The classical **Riemann–von Mangoldt** formula says that

$$N(T) = \frac{T}{2\pi} \log \left(\frac{T}{2\pi} \right) - \frac{T}{2\pi} + \frac{7}{8} + S(T) + O\left(\frac{1}{T}\right).$$

Here $N(T)$ is the number of zeta zeros with $0 < \gamma \leq T$ and we have $S(T) = \frac{1}{\pi} \arg \zeta(\frac{1}{2} + iT)$. This function **regulates the finer behavior of $\zeta(s)$** . The term $O(1/T)$ is a **smooth function** of T .

If T is the ordinate of a zeta-zero, then one defines $S(T) = S(T+0)$. Here $\arg \zeta(\frac{1}{2} + iT)$ is obtained by continuous variation along the segments joining the points $2, 2 + iT, \frac{1}{2} + iT$, starting with the value 0.

The classical **Riemann–von Mangoldt** formula says that

$$N(T) = \frac{T}{2\pi} \log \left(\frac{T}{2\pi} \right) - \frac{T}{2\pi} + \frac{7}{8} + S(T) + O\left(\frac{1}{T}\right).$$

Here $N(T)$ is the number of zeta zeros with $0 < \gamma \leq T$ and we have $S(T) = \frac{1}{\pi} \arg \zeta(\frac{1}{2} + iT)$. This function **regulates the finer behavior of $\zeta(s)$** . The term $O(1/T)$ is a **smooth function** of T .

If T is the ordinate of a zeta-zero, then one defines $S(T) = S(T+0)$. Here $\arg \zeta(\frac{1}{2} + iT)$ is obtained by continuous variation along the segments joining the points $2, 2 + iT, \frac{1}{2} + iT$, starting with the value 0.

One has the bounds

$$S(T) \ll \log T, \quad S(T) = o(\log T) \quad (\text{LH}), \quad S(T) \ll \frac{\log T}{\log \log T} \quad (\text{RH}).$$

Using the [trivial inequality](#)

$$m(\beta + i\gamma) \leq N(\gamma + H) - N(\gamma - H) \quad (0 < H \leq 1)$$

we obtain

Using the [trivial inequality](#)

$$m(\beta + i\gamma) \leq N(\gamma + H) - N(\gamma - H) \quad (0 < H \leq 1)$$

we obtain

$$m(\beta + i\gamma) \ll \log \gamma,$$

$$m(\beta + i\gamma) = o(\log \gamma) \quad (\text{LH}),$$

$$m(\beta + i\gamma) \ll \frac{\log \gamma}{\log \log \gamma} \quad (\text{RH}).$$

Using the **trivial inequality**

$$m(\beta + i\gamma) \leq N(\gamma + H) - N(\gamma - H) \quad (0 < H \leq 1)$$

we obtain

$$m(\beta + i\gamma) \ll \log \gamma,$$

$$m(\beta + i\gamma) = o(\log \gamma) \quad (\text{LH}),$$

$$m(\beta + i\gamma) \ll \frac{\log \gamma}{\log \log \gamma} \quad (\text{RH}).$$

Recall that the bound $m(\beta + i\gamma) \ll \log \gamma$ has **not been improved yet** in the whole range $\frac{1}{2} \leq \beta < 1$.

There is a **limit** for the estimation of $m(\beta + i\gamma)$ by direct **use of pointwise** estimates for $S(T)$. Namely one has

There is a **limit** for the estimation of $m(\beta + i\gamma)$ by direct **use of pointwise** estimates for $S(T)$. Namely one has

$$S(T) = \Omega_{\pm} \left(\left(\frac{\log T}{\log \log T} \right)^{1/3} \right), \quad S(T) = \Omega_{\pm} \left(\left(\frac{\log T}{\log \log T} \right)^{1/2} \right) \quad (\text{RH}),$$

but all **known values** of $S(T)$ satisfy $|S(T)| \leq 4$ (J.W. Bober-G.A. Hiary, 2016).

There is a **limit** for the estimation of $m(\beta + i\gamma)$ by direct **use of pointwise** estimates for $S(T)$. Namely one has

$$S(T) = \Omega_{\pm} \left(\left(\frac{\log T}{\log \log T} \right)^{1/3} \right), \quad S(T) = \Omega_{\pm} \left(\left(\frac{\log T}{\log \log T} \right)^{1/2} \right) \quad (\text{RH}),$$

but all **known values** of $S(T)$ satisfy $|S(T)| \leq 4$ (J.W. Bober-G.A. Hiary, 2016).

Key problem by this approach: **estimation of $S(T+H) - S(T)$.**

There is a **limit** for the estimation of $m(\beta + i\gamma)$ by direct **use of pointwise** estimates for $S(T)$. Namely one has

$$S(T) = \Omega_{\pm} \left(\left(\frac{\log T}{\log \log T} \right)^{1/3} \right), \quad S(T) = \Omega_{\pm} \left(\left(\frac{\log T}{\log \log T} \right)^{1/2} \right) \quad (\text{RH}),$$

but all **known values** of $S(T)$ satisfy $|S(T)| \leq 4$ (J.W. Bober-G.A. Hiary, 2016).

Key problem by this approach: **estimation of $S(T+H) - S(T)$** .

Here as usual $f(x) = \Omega_{\pm}(g(x))$ means that

$$\limsup_{x \rightarrow \infty} f(x)/g(x) > 0, \quad \liminf_{x \rightarrow \infty} f(x)/g(x) < 0,$$

and $g(x) > 0$ for $x \geq x_0 > 0$.

There is a **limit** for the estimation of $m(\beta + i\gamma)$ by direct **use of pointwise** estimates for $S(T)$. Namely one has

$$S(T) = \Omega_{\pm} \left(\left(\frac{\log T}{\log \log T} \right)^{1/3} \right), \quad S(T) = \Omega_{\pm} \left(\left(\frac{\log T}{\log \log T} \right)^{1/2} \right) \quad (\text{RH}),$$

but all **known values** of $S(T)$ satisfy $|S(T)| \leq 4$ (J.W. Bober-G.A. Hiary, 2016).

Key problem by this approach: **estimation of $S(T+H) - S(T)$** .

Here as usual $f(x) = \Omega_{\pm}(g(x))$ means that

$$\limsup_{x \rightarrow \infty} f(x)/g(x) > 0, \quad \liminf_{x \rightarrow \infty} f(x)/g(x) < 0,$$

and $g(x) > 0$ for $x \geq x_0 > 0$.

The Ω -results are due to **H.L. Montgomery** (1977) and **K.-M. Tsang** (1986), respectively.

A function closely related to the integral over short intervals is

$$F(T, \Delta) := \max_{t \in [T, T+\Delta]} \left| \zeta\left(\frac{1}{2} + it\right) \right| \quad (0 < \Delta \leq 1),$$

where Δ may depend on T .

A function closely related to the integral over short intervals is

$$F(T, \Delta) := \max_{t \in [T, T+\Delta]} \left| \zeta\left(\frac{1}{2} + it\right) \right| \quad (0 < \Delta \leq 1),$$

where Δ may **depend on** T .

The quantity $F(T, \Delta)$ was introduced and studied by **A.A. Karatsuba** (2001), who made the following **conjectures**.

A function closely related to the integral over short intervals is

$$F(T, \Delta) := \max_{t \in [T, T+\Delta]} \left| \zeta\left(\frac{1}{2} + it\right) \right| \quad (0 < \Delta \leq 1),$$

where Δ may depend on T .

The quantity $F(T, \Delta)$ was introduced and studied by A.A. Karatsuba (2001), who made the following conjectures.

Conjecture 1. There exists a positive function $\Delta = \Delta(T) \rightarrow 0$ as $T \rightarrow \infty$ such that, for some constant $A > 0$,

$$F(T, \Delta) \geq T^{-A}.$$

Conjecture 2. Conjecture 1 is valid for $\Delta = (\log \log T)^{-1}$.

Conjecture 2. Conjecture 1 is valid for $\Delta = (\log \log T)^{-1}$.

Conjecture 3. Conjecture 1 is valid for $\Delta = (\log T)^{-1}$.

Conjecture 2. Conjecture 1 is valid for $\Delta = (\log \log T)^{-1}$.

Conjecture 3. Conjecture 1 is valid for $\Delta = (\log T)^{-1}$.

These conjectures have not been proved **unconditionally** yet. Clearly Conjecture 3 implies Conjecture 2, which in turn implies Conjecture 1.

Conjecture 2. Conjecture 1 is valid for $\Delta = (\log \log T)^{-1}$.

Conjecture 3. Conjecture 1 is valid for $\Delta = (\log T)^{-1}$.

These conjectures have not been proved **unconditionally** yet. Clearly Conjecture 3 implies Conjecture 2, which in turn implies Conjecture 1.

M. Garaev (2002) proved that the RH implies Conjecture 3, while A.A. Karatsuba (2001) himself showed unconditionally that

$$F(T, \Delta) \geq e^{A \log \Delta \log T} \quad (0 < \Delta \leq 1/(\log T)).$$

Conjecture 2. Conjecture 1 is valid for $\Delta = (\log \log T)^{-1}$.

Conjecture 3. Conjecture 1 is valid for $\Delta = (\log T)^{-1}$.

These conjectures have not been proved **unconditionally** yet. Clearly Conjecture 3 implies Conjecture 2, which in turn implies Conjecture 1.

M. Garaev (2002) proved that the RH implies Conjecture 3, while A.A. Karatsuba (2001) himself showed unconditionally that

$$F(T, \Delta) \geq e^{A \log \Delta \log T} \quad (0 < \Delta \leq 1/(\log T)).$$

Shao-Ji Feng (2004) proved that the LH implies Conjecture 1 with an arbitrary constant $A > 0$. Other **relevant works** are due to M.E. Changa, B. Kerr and M.A. Korolev.

THEOREM 2. *If Conjecture 1 holds, then*

$$(5) \qquad m\left(\frac{1}{2} + i\gamma\right) = o(\log \gamma) \qquad (\gamma \rightarrow \infty).$$

THEOREM 2. *If Conjecture 1 holds, then*

$$(5) \qquad m\left(\frac{1}{2} + i\gamma\right) = o(\log \gamma) \qquad (\gamma \rightarrow \infty).$$

This shows again that the **LH** implies $m\left(\frac{1}{2} + i\gamma\right) = o(\log \gamma)$.

THEOREM 2. *If Conjecture 1 holds, then*

$$(5) \qquad m\left(\frac{1}{2} + i\gamma\right) = o(\log \gamma) \qquad (\gamma \rightarrow \infty).$$

This shows again that the **LH** implies $m\left(\frac{1}{2} + i\gamma\right) = o(\log \gamma)$.

Open questions: does (5) imply the **LH** or **Conjecture 1**?

THEOREM 2. *If Conjecture 1 holds, then*

$$(5) \quad m\left(\frac{1}{2} + i\gamma\right) = o(\log \gamma) \quad (\gamma \rightarrow \infty).$$

This shows again that the **LH** implies $m(\frac{1}{2} + i\gamma) = o(\log \gamma)$.

Open questions: does (5) imply the **LH** or **Conjecture 1**?

The **connection** between $F(T, \Delta)$ and integrals over short intervals is **easy**:

$$\int_{\delta}^{2\delta} |\zeta(\tfrac{1}{2} + i\gamma + i\alpha)|^k d\alpha = \int_0^{\delta} |\zeta(\tfrac{1}{2} + i\gamma + i\delta + ix)|^k dx \leq \delta F^k(\gamma + \delta, \delta),$$

where $k, \gamma > 0$.

THEOREM 2. *If Conjecture 1 holds, then*

$$(5) \qquad m\left(\frac{1}{2} + i\gamma\right) = o(\log \gamma) \qquad (\gamma \rightarrow \infty).$$

This shows again that the **LH** implies $m\left(\frac{1}{2} + i\gamma\right) = o(\log \gamma)$.

Open questions: does (5) imply the **LH** or **Conjecture 1**?

The **connection** between $F(T, \Delta)$ and integrals over short intervals is **easy**:

$$\int_{\delta}^{2\delta} |\zeta(\tfrac{1}{2} + i\gamma + i\alpha)|^k d\alpha = \int_0^{\delta} |\zeta(\tfrac{1}{2} + i\gamma + i\delta + ix)|^k dx \leq \delta F^k(\gamma + \delta, \delta),$$

where $k, \gamma > 0$.

Thus the **Karatsuba conjectures** have less stringent counterparts involving the **above integral**.

THEOREM 3. For $k > 0$, $\frac{1}{2} \leq \sigma \leq 1$, $0 < \delta \leq \frac{1}{2}$, $T \geq T_0 > 0$ and a suitable constant $C > 0$ we have

$$\int_{T-\delta}^{T+\delta} |\zeta(\sigma + it)|^k dt \geq 2\delta T^{-Ck \log(e/\delta)}.$$

THEOREM 3. For $k > 0$, $\frac{1}{2} \leq \sigma \leq 1$, $0 < \delta \leq \frac{1}{2}$, $T \geq T_0 > 0$ and a suitable constant $C > 0$ we have

$$\int_{T-\delta}^{T+\delta} |\zeta(\sigma + it)|^k dt \geq 2\delta T^{-Ck \log(e/\delta)}.$$

The starting point for the proof is the **classical formula**

$$\log \zeta(s) = \sum_{|t-\gamma| \leq 1} \log(s - \rho) + O(\log t),$$

THEOREM 3. For $k > 0$, $\frac{1}{2} \leq \sigma \leq 1$, $0 < \delta \leq \frac{1}{2}$, $T \geq T_0 > 0$ and a suitable constant $C > 0$ we have

$$\int_{T-\delta}^{T+\delta} |\zeta(\sigma + it)|^k dt \geq 2\delta T^{-Ck \log(e/\delta)}.$$

The starting point for the proof is the **classical formula**

$$\log \zeta(s) = \sum_{|t-\gamma| \leq 1} \log(s - \rho) + O(\log t),$$

which is valid **unconditionally** for

$$-1 \leq \sigma \leq 2, s \neq \rho, -\pi < \Im \log(s - \rho) \leq \pi,$$

where $\rho = \beta + i\gamma$ denotes complex zeros of $\zeta(s)$.

To get rid of the logarithms one uses (this is a consequence of the arithmetic-geometric means inequality)

$$\log \left\{ \frac{1}{b-a} \int_a^b f(t) dt \right\} \geq \frac{1}{b-a} \int_a^b \log f(t) dt$$

for $a < b$, $f(t) \in L[a, b]$ and $f(t) > 0$ in $[a, b]$.

To get rid of the logarithms one uses (this is a consequence of the **arithmetic-geometric means inequality**)

$$\log \left\{ \frac{1}{b-a} \int_a^b f(t) dt \right\} \geq \frac{1}{b-a} \int_a^b \log f(t) dt$$

for $a < b$, $f(t) \in L[a, b]$ and $f(t) > 0$ in $[a, b]$.

The integral

$$\int_{T-\delta}^{T+\delta} \sum_{|t-\gamma| \leq 1} \log |t - \gamma| dt$$

To get rid of the logarithms one uses (this is a consequence of the arithmetic-geometric means inequality)

$$\log \left\{ \frac{1}{b-a} \int_a^b f(t) dt \right\} \geq \frac{1}{b-a} \int_a^b \log f(t) dt$$

for $a < b$, $f(t) \in L[a, b]$ and $f(t) > 0$ in $[a, b]$.

The integral

$$\int_{T-\delta}^{T+\delta} \sum_{|t-\gamma| \leq 1} \log |t - \gamma| dt$$

is split into portions where $\sum_{|t-\gamma| \leq \delta} \log |t - \gamma|$ and $\sum_{\delta < |t-\gamma| \leq 1} \log |t - \gamma|$,

To get rid of the logarithms one uses (this is a consequence of the **arithmetic-geometric means inequality**)

$$\log \left\{ \frac{1}{b-a} \int_a^b f(t) dt \right\} \geq \frac{1}{b-a} \int_a^b \log f(t) dt$$

for $a < b$, $f(t) \in L[a, b]$ and $f(t) > 0$ in $[a, b]$.

The integral

$$\int_{T-\delta}^{T+\delta} \sum_{|t-\gamma| \leq 1} \log |t - \gamma| dt$$

is **split into portions** where $\sum_{|t-\gamma| \leq \delta} \log |t - \gamma|$ and $\sum_{\delta < |t-\gamma| \leq 1} \log |t - \gamma|$,

and each portion is estimated separately.

THEOREM 4. *Let $5/6 \leq \beta < 1$. Then we have, for $\gamma \geq \gamma_0(\varepsilon)$, a suitable constant $C > 0$ and any $\varepsilon > 0$,*

$$m(\beta + i\gamma) \leq C + \frac{13.35\beta}{3(1-\beta)\log 6 + \beta\log 2}(1-\beta)^{3/2}\log \gamma \\ + \frac{7(3-2\beta) + \varepsilon}{9(1-\beta)\log 6 + 3\beta\log 2}\log \log \gamma.$$

THEOREM 4. *Let $5/6 \leq \beta < 1$. Then we have, for $\gamma \geq \gamma_0(\varepsilon)$, a suitable constant $C > 0$ and any $\varepsilon > 0$,*

$$m(\beta + i\gamma) \leq C + \frac{13.35\beta}{3(1-\beta)\log 6 + \beta\log 2}(1-\beta)^{3/2}\log \gamma \\ + \frac{7(3-2\beta) + \varepsilon}{9(1-\beta)\log 6 + 3\beta\log 2}\log \log \gamma.$$

This result is relevant when β is close to unity, in which case the RH fails.

THEOREM 4. *Let $5/6 \leq \beta < 1$. Then we have, for $\gamma \geq \gamma_0(\varepsilon)$, a suitable constant $C > 0$ and any $\varepsilon > 0$,*

$$m(\beta + i\gamma) \leq C + \frac{13.35\beta}{3(1-\beta)\log 6 + \beta\log 2}(1-\beta)^{3/2}\log \gamma \\ + \frac{7(3-2\beta) + \varepsilon}{9(1-\beta)\log 6 + 3\beta\log 2}\log \log \gamma.$$

This result is relevant when β is close to unity, in which case the RH fails.

Corollary 1. For $5/6 \leq \beta < 1$ and $\gamma \geq \gamma_1$, we have

$$m(\beta + i\gamma) \leq 4\log \log \gamma + 20(1-\beta)^{3/2}\log \gamma.$$

THEOREM 4. *Let $5/6 \leq \beta < 1$. Then we have, for $\gamma \geq \gamma_0(\varepsilon)$, a suitable constant $C > 0$ and any $\varepsilon > 0$,*

$$m(\beta + i\gamma) \leq C + \frac{13.35\beta}{3(1-\beta)\log 6 + \beta\log 2}(1-\beta)^{3/2}\log \gamma \\ + \frac{7(3-2\beta) + \varepsilon}{9(1-\beta)\log 6 + 3\beta\log 2}\log \log \gamma.$$

This result is relevant when β is close to unity, in which case the RH fails.

Corollary 1. For $5/6 \leq \beta < 1$ and $\gamma \geq \gamma_1$, we have

$$m(\beta + i\gamma) \leq 4\log \log \gamma + 20(1-\beta)^{3/2}\log \gamma.$$

Corollary 2. If $m(\beta + i\gamma) \geq 8\log \log \gamma$ for $5/6 \leq \beta < 1$ and $\gamma \geq \gamma_2$, then

$$\beta \leq 1 - \left(\frac{m(\beta + i\gamma)}{40\log \gamma} \right)^{2/3}.$$

Starting point of proof: Let $\beta \geq 5/6$, $r = m(\beta + i\gamma)$ and \mathcal{E} be the rectangle with vertices $-2(1 - \beta) \pm 2i \log^2 \gamma$, $1 \pm 2i \log^2 \gamma$. If X , with $0 < X \ll \gamma^C$, is a parameter which will be suitably chosen, then by the residue theorem we obtain

$$\frac{\zeta(1 - \beta + \rho)}{(1 - \beta)^r} = \frac{1}{2\pi i} \int_{\mathcal{E}} X^{s-1+\beta} \Gamma(s-1+\beta) \frac{\zeta(s+\rho)}{s^r} ds \quad (\rho = \beta + i\gamma).$$

Starting point of proof: Let $\beta \geq 5/6$, $r = m(\beta + i\gamma)$ and \mathcal{E} be the rectangle with vertices $-2(1 - \beta) \pm 2i \log^2 \gamma$, $1 \pm 2i \log^2 \gamma$. If X , with $0 < X \ll \gamma^C$, is a parameter which will be suitably chosen, then by the residue theorem we obtain

$$\frac{\zeta(1 - \beta + \rho)}{(1 - \beta)^r} = \frac{1}{2\pi i} \int_{\mathcal{E}} X^{s-1+\beta} \Gamma(s-1+\beta) \frac{\zeta(s+\rho)}{s^r} ds \quad (\rho = \beta + i\gamma).$$

To bound the zeta-factor we shall use the **explicit inequality**

$$|\zeta(\sigma + it)| \leq A t^{B(1-\sigma)^{3/2}} \log^{2/3} t \quad (t \geq 3, \quad \frac{1}{2} \leq \sigma \leq 1).$$

Starting point of proof: Let $\beta \geq 5/6$, $r = m(\beta + i\gamma)$ and \mathcal{E} be the rectangle with vertices $-2(1 - \beta) \pm 2i \log^2 \gamma$, $1 \pm 2i \log^2 \gamma$. If X , with $0 < X \ll \gamma^C$, is a parameter which will be suitably chosen, then by the residue theorem we obtain

$$\frac{\zeta(1 - \beta + \rho)}{(1 - \beta)^r} = \frac{1}{2\pi i} \int_{\mathcal{E}} X^{s-1+\beta} \Gamma(s-1+\beta) \frac{\zeta(s+\rho)}{s^r} ds \quad (\rho = \beta + i\gamma).$$

To bound the zeta-factor we shall use the **explicit inequality**

$$|\zeta(\sigma + it)| \leq A t^{B(1-\sigma)^{3/2}} \log^{2/3} t \quad (t \geq 3, \quad \frac{1}{2} \leq \sigma \leq 1).$$

The currently **best known** values $A = 76.2$, $B = 4.45$, are due to [K. Ford](#) (2002). They are obtained by an elaboration of the method of [I.M. Vinogradov](#).

Starting point of proof: Let $\beta \geq 5/6$, $r = m(\beta + i\gamma)$ and \mathcal{E} be the rectangle with vertices $-2(1 - \beta) \pm 2i \log^2 \gamma$, $1 \pm 2i \log^2 \gamma$. If X , with $0 < X \ll \gamma^C$, is a parameter which will be suitably chosen, then by the residue theorem we obtain

$$\frac{\zeta(1 - \beta + \rho)}{(1 - \beta)^r} = \frac{1}{2\pi i} \int_{\mathcal{E}} X^{s-1+\beta} \Gamma(s-1+\beta) \frac{\zeta(s+\rho)}{s^r} ds \quad (\rho = \beta + i\gamma).$$

To bound the zeta-factor we shall use the **explicit inequality**

$$|\zeta(\sigma + it)| \leq A t^{B(1-\sigma)^{3/2}} \log^{2/3} t \quad (t \geq 3, \quad \frac{1}{2} \leq \sigma \leq 1).$$

The currently **best known** values $A = 76.2$, $B = 4.45$, are due to [K. Ford](#) (2002). They are obtained by an elaboration of the method of [I.M. Vinogradov](#).

To bound $\zeta(1 + i\gamma)$ we also use another consequence of Vinogradov's method (**zero-free region** for $\zeta(s)$):

$$\zeta(1 + it) \gg (\log |t|)^{-2/3} (\log \log |t|)^{-1/3}.$$

Thank you for your attention!