## **GEOMETRY HOMEWORK 2**

## B96201044 黃上恩, B98901182 時丕勳, K0020100x 劉士瑋

## September 25, 2011

**Problem 3** (P47: 5). If a closed plane curve C is contained inside a disk of radius r, prove that there exists a point  $p \in C$  such that the curvature  $\kappa$  of C at p satisfies  $|\kappa| \geq 1/r$ .

*Proof.* Let X(s) denote the curve C, where  $s \in [0, l]$  is an arc-length parameter, that is,  $||X'(s)|| \equiv 1$ . Since C is contained inside a disk of radius r, let A be the centre of the disk. So we have

$$||X(s) - A|| \le r \tag{1}$$

Consider  $f(s) = \langle X(s) - A, X(s) - A \rangle$ . Since [0, l] is compact, the maximum exists, denoting by  $f(s') = \max_{s \in [0, l]} f(s)$ . Therefore, we have f'(s') = 0 and  $f''(s') \leq 0$ . Now

$$f''(s) = 2(\|X'(s)\|^2 + \kappa(s)\langle X(s) - A, N(s)\rangle), \tag{2}$$

where  $X''(s) = \kappa(s)N(s)$  and N(s) is the normal vector. Take s = s' in (2) we have  $f''(s') \leq 0$  and hence

$$\kappa(s') \langle X(s) - A, N(s) \rangle \le -1 \tag{3}$$

This implies

$$|\kappa(s')\langle X(s) - A, N(s)\rangle| \ge 1$$
 (4)

By (1),  $|\langle X(s) - A, N(s) \rangle| \le ||X(s) - A|| \cdot ||N(s)|| \le r$ . We have  $|\kappa(s')| \ge 1/r$  as desired.

Problem 4 (P23: 4, 僅討論平面情形). Assume that all normals of a parametrized curve pass through a fixed point. Prove that the trace of the curve is contained in a circle.

*Proof.* Let P be the fixed point, and let X(s) be this curve. Then from description,  $\langle X(s) - P, X'(s) \rangle \equiv 0$  for all s. Let  $f(s) = \|X(s) - P\|^2$ , then we have  $f'(s) = 2 \langle X(s) - P, X'(s) \rangle = 0$  for all s. This implies the trace of the curve is contained in a circle centered at point P with radius  $\sqrt{f(s_0)}$  for some  $s_0$ .  $\square$ 

**Problem 5.** 以 t=0 開始將曲線  $(t^2,t^3)$  化成長度參數。並討論 t=0 時的曲率。

Problem 6.

- (a) 以原點為中心,將 y=f(x) 的圖形縮放  $\lambda$  倍,並說明新的圖形是  $y=\lambda f(\frac{x}{\lambda})$  的函數圖形。
- (b) 討論曲率的變化。

Problem 7. 如圖,有一橢圓,其焦點為  $O_1$  和  $O_2$ ,設 L 切橢圓於 P,且 L 與  $\overline{O_2P}$  之夾角為  $\theta$ 。以  $\theta$  為參數,說明曲率  $\kappa \propto \sin^3 \theta$ 

Problem 9. 如圖,有 regular curve  $\gamma(t)$ ,  $\gamma_0=\gamma(0)$ ,  $N_0=N(0)$ ,  $L_0=\{\gamma_0+vN_0\}$ 。 現考慮直線  $L_t=\{\gamma(t)+uN(t)\}$ ,令  $P(t)=L_t\cap L_0$  證明

$$\kappa(0) \neq 0 \Rightarrow \lim_{t \to 0} P(t) = \gamma_0 + \frac{1}{\kappa(0)} N_0$$