

Geometry Homework 1

B96201044 黃上恩, B98901182 時丕勳, K0020100x 劉士瑋

September 19, 2011

Problem 3 (P7: 4). Let $\alpha : (0, \pi) \rightarrow \mathbf{R}^2$ be given by

$$\alpha(t) = \left(\sin t, \cos t + \log \tan \frac{t}{2} \right),$$

where t is the angle that the y axis makes with the vector $\alpha(t)$. The trace of α is called the tractrix (Fig. 1-9). Show that

- (a) α is a differentiable parametrized curve, regular except at $t = \pi/2$.
- (b) The length of the segment of the tangent of the tractrix between the point of tangency and the y axis is constantly equal to 1.

Proof.

- (a) Let $x(t) = \sin t$, $y(t) = \cos t + \log \tan \frac{t}{2}$, then

$$x'(t) = \cos t; \quad y'(t) = -\sin t + \frac{1}{\sin t}.$$

It's trivial that both $x'(t)$ and $y'(t)$ are infinitely differentiable in $(0, \pi)$, so α is a differentiable parametrized curve.

$x'(t) = 0, y'(t) = 0 \iff t = \frac{\pi}{2}$, so α is regular except at $t = \pi/2$.

- (b) The intersection of y axis and the tangent of the tractrix is $\left(0, y(t) - \frac{y'(t)}{x'(t)}x(t)\right)$.
The length of the segment of the tangent of the tractrix between the point of tangency and the y axis is $\sqrt{x(t)^2 + \left(\frac{y'(t)}{x'(t)}x(t)\right)^2}$

$$\begin{aligned}
x(t)^2 + \left(\frac{y'(t)}{x'(t)} x(t) \right)^2 &= \sin^2 t \left(1 + \left(\frac{y'(t)}{x'(t)} \right)^2 \right) \\
&= \sin^2 t \left(1 + \left(\frac{-\sin t + \frac{1}{\sin t}}{\cos t} \right)^2 \right) \\
&= \sin^2 t \left(1 + \left(\frac{1 - \sin^2 t}{\sin t \cos t} \right)^2 \right) \\
&= \sin^2 t \left(\frac{1}{\sin^2 t} \right) \\
&= 1
\end{aligned}$$

So the length of the segment of the tangent of the tractrix between the point of tangency and the y axis $= \sqrt{x(t)^2 + \left(\frac{y'(t)}{x'(t)} x(t) \right)^2} = 1$.

□

Problem 5 (P47: 6). Let $\alpha(s)$, $s \in [0, l]$ be a closed convex plane curve positively oriented. The curve

$$\beta(s) = \alpha(s) - rn(s),$$

where r is a positive constant and n is the normal vector, is called a parallel curve to α (Fig. 1-37). Show that

(a) Length of β = length of α + $2\pi r$.

(b) $A(\beta) = A(\alpha) + rl + \pi r^2$.

(c) $\kappa_\beta(s) = \kappa_\alpha(s)/(1 + r\kappa_\alpha(s))$.

For (a)-(c), $A(\cdot)$ denotes the area bounded by the corresponding curve, and $\kappa_\alpha, \kappa_\beta$ are the curvatures of α and β , respectively.

Proof.

□

Problem 8 (Curvature is a geometric object I.). $X(s) = (x(s), y(s))$, where s is the arc-length parameter.

$$M = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, M^t = M^{-1}, \text{ i.e. } M \text{ is orthogonal.}$$

Let $\bar{X}(s) = M \cdot \begin{bmatrix} x(s) \\ y(s) \end{bmatrix} + \begin{bmatrix} \alpha \\ \beta \end{bmatrix}$, $\alpha, \beta \in \mathbf{R}$. What is the relation between $\kappa_X(s)$ and $\kappa_{\bar{X}}(s)$?

Proof. □

Problem 9 (Curvature is a geometric object II.). $X(t) = (x(t), y(t))$ be a regular curve. Let

$$\kappa(x(t), y(t)) \equiv \kappa(t) = \frac{\begin{vmatrix} x' & y' \\ x'' & y'' \end{vmatrix}}{(x'^2 + y'^2)^{\frac{3}{2}}}$$

Let $Y(u) = X(t(u))$, $t'(u) \neq 0$. Discuss the relation of $\kappa(x(t), y(t))$ and $\kappa(x(t(u)), y(t(u)))$ at the corresponding points.

Proof. □

Problem 10. Let $F(x, y) = c$ defines a plane curve. Prove that the curvature of the curve satisfies

$$|\kappa| = \left| \frac{\begin{bmatrix} F_y & -F_x \end{bmatrix} \begin{bmatrix} F_{xx} & F_{xy} \\ F_{xy} & F_{yy} \end{bmatrix} \begin{bmatrix} F_y \\ -F_x \end{bmatrix}}{(F_x^2 + F_y^2)^{\frac{3}{2}}} \right|$$

Where $F_x^2 + F_y^2 \neq 0$.

Proof. Let α be a point in the plane such that $F(\alpha) = c$. Consider the circle of curvature passing through α . If we observed the intersection of two line respectively perpendicular to the lines tangent to $F = c$ and passing through α and another point $\alpha' \in F = c$, the center of the circle o will be the limit as $\alpha' \rightarrow \alpha$. Thus,

$$|\alpha' - \alpha| = r \sin \theta$$

, where θ is the angle between the vectors $o - \alpha$ and $o - \alpha'$. By the formula of exterior product,

$$\sin \theta = \frac{|(o - \alpha) \times (o - \alpha')|}{|o - \alpha||o - \alpha'|}$$

Let n denote $\nabla F / |\nabla F|$ rotated counterclockwise by $\pi/2$. Since $\alpha' - \alpha$ is perpendicular to $\frac{\nabla F(\alpha)}{|\nabla F(\alpha)|}$, and since $o - \alpha$ and $o - \alpha'$ are respectively

parallel to $\frac{\nabla F(\alpha)}{|\nabla F(\alpha)|}$ and $\frac{\nabla F(\alpha')}{|\nabla F(\alpha')|}$, we obtained

$$\begin{aligned}
|\kappa| = 1/r &= \lim_{\alpha' \rightarrow \alpha} \frac{|(o - \alpha) \times (o - \alpha')|}{|\alpha' - \alpha||o - \alpha||o - \alpha'|} = \lim_{\alpha' \rightarrow \alpha} \frac{|\nabla F(\alpha) \times \nabla F(\alpha')|}{|\alpha' - \alpha||\nabla F(\alpha)||\nabla F(\alpha')|} \\
&= \lim_{t \rightarrow 0} \frac{|\nabla F(\alpha) \times \nabla F(\alpha + tn)|}{tn|\nabla F(\alpha)||\nabla F(\alpha + tn)|} \\
&= \lim_{t \rightarrow 0} \frac{|(F_x(\alpha)\vec{i} + F_y(\alpha)\vec{j}) \times (F_x(\alpha + tn)\vec{i} + F_y(\alpha + tn)\vec{j})|}{tn|\nabla F(\alpha)|^2} \\
&= \lim_{t \rightarrow 0} \frac{|F_x(\alpha)F_y(\alpha + tn) - F_x(\alpha + tn)F_y(\alpha)|}{tn|\nabla F(\alpha)|^2} \\
&= \lim_{t \rightarrow 0} \frac{|F_x(\alpha)[F_y(\alpha + tn) - F_y(\alpha)] - [F_x(\alpha + tn) - F_x(\alpha)]F_y(\alpha)|}{tn|\nabla F(\alpha)|^2} \\
&= \frac{|F_x(F_{yx}\vec{i} + F_{yy}\vec{j}) \cdot n - F_y(F_{xx}\vec{i} + F_{xy}\vec{j}) \cdot n|}{|\nabla F|^2} \\
&= \frac{|[(F_x F_{yx} - F_y F_{xx})\vec{i} + (F_x F_{yy} - F_y F_{xy})\vec{j}] \cdot n|}{|\nabla F|^2} \\
&= \frac{|[(F_x F_{yx} - F_y F_{xx})\vec{i} + (F_x F_{yy} - F_y F_{xy})\vec{j}] \cdot (F_y\vec{i} - F_x\vec{j})/\nabla F|}{|\nabla F|^2} \\
&= \frac{\left| \begin{bmatrix} F_y & -F_x \end{bmatrix} \begin{bmatrix} F_{xx} & F_{xy} \\ F_{xy} & F_{yy} \end{bmatrix} \begin{bmatrix} F_y \\ -F_x \end{bmatrix} \right|}{|\nabla F|^3} \\
&= \left| \frac{\begin{bmatrix} F_y & -F_x \end{bmatrix} \begin{bmatrix} F_{xx} & F_{xy} \\ F_{xy} & F_{yy} \end{bmatrix} \begin{bmatrix} F_y \\ -F_x \end{bmatrix}}{(F_x^2 + F_y^2)^{\frac{3}{2}}} \right|
\end{aligned}$$

□