

GEOMETRY HOMEWORK 2

B96201044 黃上恩, B98901182 時丕勳, K0020100x 劉士璋

September 28, 2011

Problem 3 (P47: 5). *If a closed plane curve C is contained inside a disk of radius r , prove that there exists a point $p \in C$ such that the curvature κ of C at p satisfies $|\kappa| \geq 1/r$.*

Proof. Let $X(s)$ denote the curve C , where $s \in [0, l]$ is an arc-length parameter, that is, $\|X'(s)\| \equiv 1$. Since C is contained inside a disk of radius r , let A be the centre of the disk. So we have

$$\|X(s) - A\| \leq r \quad (1)$$

Consider $f(s) = \langle X(s) - A, X(s) - A \rangle$. Since $[0, l]$ is compact, the maximum exists, denoting by $f(s') = \max_{s \in [0, l]} f(s)$. Therefore, we have $f'(s') = 0$ and $f''(s') \leq 0$. Now

$$f''(s) = 2 (\|X'(s)\|^2 + \kappa(s) \langle X(s) - A, N(s) \rangle), \quad (2)$$

where $X''(s) = \kappa(s)N(s)$ and $N(s)$ is the normal vector. Take $s = s'$ in (2) we have $f''(s') \leq 0$ and hence

$$\kappa(s') \langle X(s) - A, N(s) \rangle \leq -1 \quad (3)$$

This implies

$$|\kappa(s') \langle X(s) - A, N(s) \rangle| \geq 1 \quad (4)$$

By (1), $|\langle X(s) - A, N(s) \rangle| \leq \|X(s) - A\| \cdot \|N(s)\| \leq r$. We have $|\kappa(s')| \geq 1/r$ as desired. \square

Problem 4 (P23: 4, 僅討論平面情形). *Assume that all normals of a parametrized curve pass through a fixed point. Prove that the trace of the curve is contained in a circle.*

Proof. Let P be the fixed point, and let $X(s)$ be this curve. Then from description, $\langle X(s) - P, X'(s) \rangle \equiv 0$ for all s . Let $f(s) = \|X(s) - P\|^2$, then we have $f'(s) = 2 \langle X(s) - P, X'(s) \rangle = 0$ for all s . This implies the trace of the curve is contained in a circle centered at point P with radius $\sqrt{f(s_0)}$ for some s_0 . \square

Problem 5. 以 $t = 0$ 開始將曲線 (t^2, t^3) 化成長度參數。並討論 $t = 0$ 時的曲率。

Proof. Consider $t > 0$, the length of the curve of t is

$$\int_0^t 3t \sqrt{(4/9) + t^2} dt = \int_0^t \frac{3}{2} \sqrt{(4/9) + t^2} dt^2 = \left(\frac{4}{9} + t^2 \right)^{3/2} - \frac{8}{27} \quad (5)$$

Let $s > 0$ be the arc-length parameter, note that $s > 0$ equivalent to $t > 0$, so we have $s = (4/9 + t^2)^{3/2} - 8/27$ and hence

$$t = \sqrt{\left(s + \frac{8}{27} \right)^{2/3} - \frac{4}{9}} \quad (6)$$

Therefore the curve with arc-length parameter $s > 0$ is

$$\left((s + 8/27)^{2/3} - 4/9, \left((s + 8/27)^{2/3} - 4/9 \right)^{3/2} \right) \quad (7)$$

By symmetry, for the case $s < 0$ the corresponding curve is

$$\left((8/27 - s)^{2/3} - 4/9, - \left((8/27 - s)^{2/3} - 4/9 \right)^{3/2} \right) \quad (8)$$

We can write them together to get the result,

$$\left((8/27 + |s|)^{2/3} - 4/9, \text{sign}(s) \cdot \left((8/27 + |s|)^{2/3} - 4/9 \right)^{3/2} \right) \quad (9)$$

□

Problem 6.

(a) 以原點為中心，將 $y = f(x)$ 的圖形縮放 λ 倍，並說明新的圖形是 $y = \lambda f(\frac{x}{\lambda})$ 的函數圖形。

(b) 討論曲率的變化。

Proof.

□

Problem 7. 如圖，有一橢圓，其焦點為 O_1 和 O_2 ，設 L 切橢圓於 P ，且 L 與 $\overline{O_2P}$ 之夾角為 θ 。以 θ 為參數，說明曲率 $\kappa \propto \sin^3 \theta$

Proof.

□

Problem 9. 如圖，有 regular curve $\gamma(t)$ ， $\gamma_0 = \gamma(0)$ ， $N_0 = N(0)$ ， $L_0 = \{\gamma_0 + v N_0\}$ 。現考慮直線 $L_t = \{\gamma(t) + u N(t)\}$ ，令 $P(t) = L_t \cap L_0$ 證明

$$\kappa(0) \neq 0 \Rightarrow \lim_{t \rightarrow 0} P(t) = \gamma_0 + \frac{1}{\kappa(0)} N_0$$

$$\text{Proof. } \gamma_0 + v N_0 = \gamma(t) + u N(t) \Rightarrow v = \frac{\begin{vmatrix} \gamma(t) - \gamma_0 & N(t) \\ N_0 & N(t) \end{vmatrix}}{\begin{vmatrix} N_0 & N(t) \end{vmatrix}} \quad \square$$