

GEOMETRY HOMEWORK 5

B96201044 黃上恩, B98901182 時丕勳, K0020100x 劉士璋

October 17, 2011

Problem 1 (參見 P67 Ex16). 考慮

$$\begin{aligned} \mathbb{X}: \quad \mathbb{R}^2 &\rightarrow S^2 \setminus \{N\} \\ (u, v) &\mapsto \left(\frac{2u}{u^2+v^2+1}, \frac{2v}{u^2+v^2+1}, \frac{u^2+v^2-1}{u^2+v^2+1} \right), N = (0, 0, 1) \end{aligned}$$

- (a) 檢查這的確是 $S^2 \setminus \{N\}$ 的參數式
- (b) 計算 E, F, G , $E = G$ 嗎?
- (c) 計算 $\mathbb{X}_u, \mathbb{X}_v$
- (d) 若 W_1, W_2 是 \mathbb{R}^2 兩以 a 為起點的向量, 說明 W_1, W_2 的夾角 $= d\mathbb{X}(W_1)$ 與 $d\mathbb{X}(W_2)$ 的夾角

Proof.

□

Problem 2 (旋轉面). $\mathbb{X}(\theta, s) = (a(s) \cos \theta, a(s) \sin \theta, b(s))$, 其中 $(a(s), b(s))$ 為長度參數之平面曲線。計算 E, F, G 並討論其 *regular* 的條件。

Proof.

$$\begin{aligned} \mathbb{X}_\theta &= (-a(s) \sin \theta, a(s) \cos \theta, 0) \\ \mathbb{X}_s &= (a'(s) \cos \theta, a'(s) \sin \theta, b'(s)) \\ \Rightarrow E &= a(s)^2 \sin^2 \theta + a(s)^2 \cos^2 \theta \\ &= a(s)^2 \\ F &= -a(s)a'(s) \sin \theta \cos \theta + a(s)a'(s) \cos \theta \sin \theta \\ &= 0 \\ G &= a'(s)^2 \sin^2 \theta + a'(s)^2 \cos^2 \theta \\ &= a'(s)^2 \\ &= 1 \\ |\mathbb{X}_\theta \times \mathbb{X}_s| &= \sqrt{EG - F^2} \\ &= \sqrt{a(s)^2} \\ &= |a(s)| \end{aligned}$$

So \mathbb{X} is regular iff $a(s) \neq 0$. □

Problem 3 (管面). 設空間曲線 $\gamma(s)$, s 長度參數, $\vec{t}, \vec{n}, \vec{b}$ 為 *Frenet frame*. 令 $\mathbb{X}_l(s, \theta) = \gamma(s) + l \cos \theta \vec{n}(s) + l \sin \theta \vec{b}(s)$, $l > 0$, 計算 E, F, G 並討論其 *regular* 條件。

Proof.

$$\begin{aligned}
\mathbb{X}_{ls} &= \gamma'(s) + l \cos \theta \vec{n}'(s) + l \sin \theta \vec{b}'(s) \\
&= \vec{t}(s) + l \cos \theta \left(-\kappa(s) \vec{t}(s) - \tau(s) \vec{b}(s) \right) + l \tau(s) \sin \theta \vec{n}(s) \\
&= (1 - l\kappa(s) \cos \theta) \vec{t}(s) + l \tau(s) \sin \theta \vec{n}(s) - l \tau(s) \cos \theta \vec{b}(s) \\
\mathbb{X}_{l\theta} &= -l \sin \theta \vec{n}(s) + l \cos \theta \vec{b}(s) \\
\Rightarrow E &= (1 - l\kappa(s) \cos \theta)^2 + (l \tau(s) \sin \theta)^2 + (l \tau(s) \cos \theta)^2 \\
&= (1 - l\kappa(s) \cos \theta)^2 + l^2 \tau(s)^2 \\
F &= -l^2 \tau(s) \sin^2 \theta - l^2 \tau(s) \cos^2 \theta \\
&= -l^2 \tau(s) \\
G &= l^2 \sin^2 \theta + l^2 \cos^2 \theta \\
&= l^2 \\
|\mathbb{X}_{ls} \times \mathbb{X}_{l\theta}| &= \sqrt{EG - F^2} \\
&= \sqrt{l^2 ((1 - l\kappa(s) \cos \theta)^2 + l^2 \tau(s)^2) - l^4 \tau(s)^2} \\
&= \sqrt{l^2 (1 - l\kappa(s) \cos \theta)^2} \\
&= l |1 - l\kappa(s) \cos \theta|
\end{aligned}$$

So \mathbb{X} is regular iff $l\kappa(s) \cos \theta \neq 1 \forall s, \theta$

$$\Leftrightarrow \frac{1}{l\kappa(s)} \neq \cos \theta$$

$$\Leftrightarrow \left| \frac{1}{l\kappa(s)} \right| > 1$$

$$\Leftrightarrow |\kappa(s)| < \frac{1}{l}$$

□

Problem 6 (Ex6, p100). Show that

$$\mathbf{x}(u, v) = (u \sin \alpha \cos v, u \sin \alpha \sin v, u \cos \alpha)$$

where $0 < u < \infty, 0 < v < 2\pi, \alpha = \text{const.}$, is a parametrization of the cone with 2α as the angle of the vertex. In the corresponding coordinate neighborhood, prove that the curve

$$\mathbf{x}(c \exp(v \sin \alpha \cot \beta), v), \quad c = \text{const.}, \beta = \text{const.},$$

intersects the generators of the cone ($v = \text{const.}$) under the constant angle β .

Proof.

$$d\mathbf{x} = \begin{bmatrix} \sin \alpha \cos v & \sin \alpha \sin v & \cos \alpha \\ -u \sin \alpha \sin v & u \sin \alpha \cos v & 0 \end{bmatrix}$$

We have $\mathbf{x}_u \times \mathbf{x}_v = (-, -, u) \neq 0$ whenever $u > 0$. To show that the angle is β , define

$$E := \langle \mathbf{x}_u, \mathbf{x}_u \rangle = \sin^2 \alpha \cos^2 v + \sin^2 \alpha \sin^2 v + \cos^2 \alpha = 1$$

$$F := \langle \mathbf{x}_u, \mathbf{x}_v \rangle = -u \sin^2 \alpha \cos v \sin v + u \sin^2 \alpha \sin v \cos v + 0 = 0$$

$$G := \langle \mathbf{x}_v, \mathbf{x}_v \rangle = u^2 \sin^2 \alpha \sin^2 v + u^2 \sin^2 \alpha \cos^2 v + 0 = u^2 \sin^2 \alpha$$

$$\langle\langle (a, b), (c, d) \rangle\rangle := \begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} E & F \\ F & G \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix}$$

to be the first fundamental form of \mathbf{x} . Then

$$A := \partial(u, v)/\partial u = (1, 0)$$

$$B := \partial(c \exp(v \sin \alpha \cot \beta), v)/\partial v = (c \sin \alpha \cot \beta \exp(v \sin \alpha \cot \beta), 1).$$

Fixing the intersection at $(c \exp(v \sin \alpha \cot \beta), v)$, we got

$$\begin{aligned} \langle\langle A, B \rangle\rangle &= \begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} E & F \\ F & G \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & c^2 \exp(2v \sin \alpha \cot \beta) \sin^2 \alpha \end{bmatrix} \begin{bmatrix} c \sin \alpha \cot \beta \exp(v \sin \alpha \cot \beta) \\ 1 \end{bmatrix} \\ &= c \sin \alpha \cot \beta \exp(v \sin \alpha \cot \beta) \end{aligned}$$

$$\begin{aligned} \langle\langle A, A \rangle\rangle &= \begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} E & F \\ F & G \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & c^2 \exp(2v \sin \alpha \cot \beta) \sin^2 \alpha \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \\ &= 1 \end{aligned}$$

$$\begin{aligned} \langle\langle B, B \rangle\rangle &= \begin{bmatrix} c & d \end{bmatrix} \begin{bmatrix} E & F \\ F & G \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} \\ &= \begin{bmatrix} c \sin \alpha \cot \beta \exp(v \sin \alpha \cot \beta) & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & c^2 \exp(2v \sin \alpha \cot \beta) \sin^2 \alpha \end{bmatrix} \\ &\quad \begin{bmatrix} c \sin \alpha \cot \beta \exp(v \sin \alpha \cot \beta) \\ 1 \end{bmatrix} \\ &= c^2 \sin^2 \alpha \cot^2 \beta \exp(2v \sin \alpha \cot \beta) + c^2 \sin^2 \alpha \exp(2v \sin \alpha \cot \beta) \\ &= c^2 \sin^2 \alpha \csc^2 \beta \exp(2v \sin \alpha \cot \beta) \end{aligned}$$

$$\cos \theta = \frac{\langle\langle A, B \rangle\rangle}{\sqrt{\langle\langle A, A \rangle\rangle \langle\langle B, B \rangle\rangle}} = \frac{c \sin \alpha \cot \beta \exp(v \sin \alpha \cot \beta)}{|c \sin \alpha \csc \beta \exp(v \sin \alpha \cot \beta)|} = \pm \cos \beta$$

□