

# GEOMETRY HOMEWORK 12

B96201044 黃上恩, B98901182 時丕勳, K0020100x 劉士璋

December 22, 2011

**Problem 3** (Ex p294 3.). If  $p$  is a point of a regular surface  $S$ , prove that

$$K(p) = \lim_{r \rightarrow 0} \frac{12}{\pi} \frac{\pi r^2 - A}{r^4},$$

where  $K(p)$  is the Gaussian curvature of  $S$  at  $p$ ,  $r$  is the radius of a geodesic circle  $S_r(p)$  centered in  $p$ , and  $A$  is the area of the region bounded by  $S_r(p)$ .

*Proof.*

$$\begin{aligned} A_R &= \int_0^R \int_0^{2\pi} \sqrt{EG - F^2} d\theta dr \\ &= \int_0^R \int_0^{2\pi} \sqrt{G} d\theta dr \\ &\approx \int_0^R \int_0^{2\pi} r - \frac{K}{6} r^3 d\theta dr \\ &= \int_0^{2\pi} \left( \frac{1}{2} R^2 - \frac{K}{24} R^4 \right) d\theta \\ &= \pi R^2 - \frac{R^4}{24} \int_0^{2\pi} K d\theta \\ &\rightarrow \frac{1}{2\pi} \int_0^{2\pi} K d\theta = \frac{12}{r^4} \left( r^2 - \frac{1}{\pi} A_r \right) \\ &\rightarrow K(p) = \lim_{r \rightarrow 0} \frac{12}{r^4} \left( r^2 - \frac{1}{\pi} A_r \right) \\ &= \lim_{r \rightarrow 0} \frac{12}{\pi} \frac{\pi r^2 - A_r}{r^4} \end{aligned}$$

□

**Problem 4** (Ex p295 4.). Show that in a system of normal coordinates centered in  $p$ , all the Christoffel symbols are zero at  $p$ .

*Proof.* Let  $(u, v)$  be normal coordinate centered at  $p$ ,  $(r, \theta)$  be the geodesic polar coordinate centered at  $p$ .

Let  $\hat{E}, \hat{F}, \hat{G}$  be the first fundamental form of the coordinate  $(r, \theta)$ ,  $E, F, G$  be the first fundamental form of the coordinate  $(u, v)$ ,

$$\begin{aligned}\hat{E} &= 1, \hat{F} = 0 \\ \lim_{r \rightarrow 0} \hat{G} &= 0, \lim_{r \rightarrow 0} \sqrt{\hat{G}} = 1 \\ &\rightarrow \hat{G} = r^2 + o(r^3)\end{aligned}$$

$$\begin{aligned}r &= \sqrt{u^2 + v^2} \\ \theta &= \tan^{-1} \frac{v}{u} \\ \mathbb{X}_u &= \frac{u}{r} \mathbb{X}_r - \frac{v}{r^2} \mathbb{X}_\theta \\ \mathbb{X}_v &= \frac{v}{r} \mathbb{X}_r + \frac{u}{r^2} \mathbb{X}_\theta \\ \rightarrow E &= \frac{u^2}{r^2} + \frac{v^2}{r^4} \hat{G} \\ F &= \frac{uv}{r^2} - \frac{uv}{r^4} \hat{G} \\ G &= \frac{v^2}{r^2} + \frac{u^2}{r^4} \hat{G}\end{aligned}$$

When  $r \rightarrow 0$ :

$$\begin{aligned}\hat{G} &\rightarrow r^2 \\ \hat{G}_u &\rightarrow 2u \\ \hat{G}_v &\rightarrow 2v\end{aligned}$$

$$\begin{aligned}E_u &= \frac{2uv^2}{r^4} - \frac{4uv^2}{r^6} \hat{G} + \frac{v^2}{r^4} \hat{G}_u \\ &= \frac{2uv^2}{r^4} - \frac{4uv^2}{r^4} + \frac{2uv^2}{r^4} = 0 \\ E_v &= -\frac{2u^2v}{r^4} + \frac{2v(u^2 - v^2)}{r^6} \hat{G} + \frac{v^2}{r^4} \hat{G}_v \\ &= -\frac{2u^2v}{r^4} + \frac{2v(u^2 - v^2)}{r^4} + \frac{2v^3}{r^4} = 0 \\ F_u &= \frac{v^3 - vu^2}{r^4} - \frac{v^3 - 3u^2v}{r^6} \hat{G} - \frac{uv}{r^4} \hat{G}_u \\ &= \frac{v^3 - vu^2}{r^4} - \frac{v^3 - 3u^2v}{r^4} - \frac{2u^2v}{r^4} = 0 \\ F_v &= \frac{u^3 - uv^2}{r^4} - \frac{u^3 - 3v^2u}{r^6} \hat{G} - \frac{uv}{r^4} \hat{G}_v \\ &= \frac{u^3 - uv^2}{r^4} - \frac{u^3 - 3v^2u}{r^4} - \frac{2uv^2}{r^4} = 0\end{aligned}$$

$$\begin{aligned}
G_u &= -\frac{2v^2u}{r^4} + \frac{2u(v^2 - u^2)}{r^6}\hat{G} + \frac{u^2}{r^4}\hat{G}_u \\
&= -\frac{2v^2u}{r^4} + \frac{2u(v^2 - u^2)}{r^4} + \frac{2u^3}{r^4} = 0 \\
G_v &= \frac{2vu^2}{r^4} - \frac{4vu^2}{r^6}\hat{G} + \frac{u^2}{r^4}\hat{G}_v \\
&= \frac{2vu^2}{r^4} - \frac{4vu^2}{r^4} + \frac{2vu^2}{r^4} = 0
\end{aligned}$$

So  $[i, j, k] = 0$  and  $\Gamma_{ij}^k = 0$ . □

**Problem 5** (Ex p295 5.). *For which of the pair of surfaces given below does there exist a local isometry?*

(a) *Torus of revolution and cone.*

(b) *Cone and sphere.*

(c) *Cone and cylinder.*

**Problem 8.**

(a) 在半徑  $R$  的球面上，計算 *geodesic circle* 的長度，並驗證 P292 課文中間  $K(p)$  的公式。

(b) 用一樣的精神，檢驗 P294 3. 的公式。

*Proof.* (a) WLOG, let  $p = (0, 0, R)$ , If  $q \in T_p$  with  $q = (l, \theta)$ , then

$$\exp(q) = \left( R \sin \frac{l}{R} \cos \theta, R \sin \frac{l}{R} \sin \theta, R \cos \frac{l}{R} \right),$$

and thus the length of the image of the circle  $\{q \in T_p : d(q, p) = l\}$  is

$$2\pi \left\langle \frac{\partial \exp(q)}{\partial \theta}, \frac{\partial \exp(q)}{\partial \theta} \right\rangle^{1/2} = 2\pi \left| R \sin \frac{l}{R} \right|$$

. When  $l \rightarrow 0$ , it is

$$2\pi R \sin \frac{l}{R}.$$

, which is the length of the geodesic circle. By the formula,

$$\begin{aligned}
K(p) &= \lim_{r \rightarrow 0} \frac{3}{\pi} \frac{2\pi r - L}{r^3} = \lim_{r \rightarrow 0} \frac{3}{\pi} \frac{2\pi r - 2\pi R \sin \frac{l}{R}}{r^3} \\
&\approx \lim_{r \rightarrow 0} \frac{3}{\pi} \frac{2\pi r - 2\pi R \left( r/R - \left( \frac{r}{R} \right)^3 / 6 \right)}{r^3} \\
&= \lim_{r \rightarrow 0} \frac{3}{\pi} \frac{2\pi r \left( \frac{r}{R} \right)^3 / 6}{r^3} = \frac{1}{R^2}.
\end{aligned}$$

(b) The area bounded by the geodesic circle is

$$2\pi R \int_0^l \left| \sin \frac{r}{R} \right| dr.$$

When  $l \rightarrow 0$ , it is

$$2\pi R^2 - 2\pi R^2 \cos \frac{l}{R}.$$

By the formula,

$$\begin{aligned} K(p) &= \lim_{r \rightarrow 0} \frac{12}{\pi} \frac{\pi r^2 - A}{r^4} = \lim_{r \rightarrow 0} \frac{12}{\pi} \frac{\pi r^2 + 2\pi R^2 \cos \frac{r}{R} - 2\pi R^2}{r^4} \\ &\approx \lim_{r \rightarrow 0} \frac{12}{\pi} \frac{\pi r^2 + 2\pi R^2 \left( 1 - \left( \frac{r}{R} \right)^2 / 2 + \left( \frac{r}{R} \right)^4 / 24 \right) - 2\pi R^2}{r^4} \\ &= \lim_{r \rightarrow 0} \frac{12}{\pi} \frac{\pi R^2 \left( \frac{r}{R} \right)^4 / 12}{r^4} = \frac{1}{R^2} \end{aligned}$$

□