






Children (<5 years)
p=0.87 n pairs = 31
a
a
Clinical outcome rate (%)
100
80
60
40
Monoinfection
Co-infection



Adolescents (5-17 years)
p=0.038 n pairs = 50
a
b
Clinical outcome rate (%)
120
100
80
60
40
20
Monoinfection
Co-infection



Adults (18-60 years)
p=0.15 n pairs = 35
a
a
Clinical outcome rate (%)
100
80
60
40
20
Monoinfection
Co-infection



Older adults (>60 years)
p=0.0014 n pairs = 46
a
b
Clinical outcome rate (%)
100
80
60
Monoinfection
Co-infection



Hospitalization
p=0.23 n pairs = 31
a
a
Clinical outcome rate (%)
120
100
80
60
40
20
Monoinfection
Co-infection



Hospitalization
p=0.44 n pairs = 50
a
a
Clinical outcome rate (%)
120
100
80
60
40
20
Monoinfection
Co-infection



Hospitalization
p=0.81 n pairs = 35
a
a
Clinical outcome rate (%)
100
80
60
40
20
Monoinfection
Co-infection



Hospitalization
p=0.44 n pairs = 46
a
a
Clinical outcome rate (%)
110
100
90
80
70
60
50
Monoinfection
Co-infection



ICU admission
p=0.02 n pairs = 30
a
b
Clinical outcome rate (%)
10
8
6
4
2
0
Monoinfection
Co-infection



ICU admission
p=0.74 n pairs = 49
a
a
Clinical outcome rate (%)
10
8
6
4
2
0
Monoinfection
Co-infection



ICU admission
p=0.15 n pairs = 35
a
a
Clinical outcome rate (%)
20
15
10
5
0
Monoinfection
Co-infection


ICU admission
p=0.00029 n pairs = 44
a
b
Clinical outcome rate (%)
50
40
30
20
10
0
Monoinfection
Co-infection


Death
p=0.012 n pairs = 30
a
b
Clinical outcome rate (%)
0.30
0.25
0.20
0.15
0.10
0.05
0.00
Monoinfection
Co-infection


Death
p=0.11 n pairs = 48
a
a
Clinical outcome rate (%)
0.15
0.10
0.05
0.00
Monoinfection
Co-infection


Death
p<0.0001 n pairs = 35
a
b
Clinical outcome rate (%)
5
4
3
2
1
0
Monoinfection
Co-infection


Death
p=0.4 n pairs = 44
a
a
Clinical outcome rate (%)
20
15
10
5
0
Monoinfection
Co-infection