

A
LRTI, 5-17 years
p=0.038 n = 50
ICU admission, >60 years
p=0.00029 n = 44
ICU admission, <5 years
p=0.020 n = 30
Death, 18-60 years
p<0.0001 n = 35
Clinical outcome rate (%)
Mono-detection
Co-detection
B
LRTI
Candida + HMPV 99.0% (n=96)
Candida + RSV 97.9% (n=890)
EBV + S maltophilia 97.3% (n=148)
RSV + S pneumoniae 97.1% (n=170)
CP + EBV 96.4% (n=112)
Hospitalization
Candida + HBoV 100.0% (n=100)
Candida + RSV 99.6% (n=890)
Candida + MP 99.4% (n=174)
Candida + IFV 99.1% (n=532)
Candida + HMPV 99.0% (n=96)
ICU
EBV + S pneumoniae 34.2% (n=114)
EBV + P aeruginosa 31.9% (n=77)
EBV + S maltophilia 30.8% (n=148)
K pneumoniae + P aeruginosa 29.4% (n=390)
EBV + H influenzae 26.4% (n=141)
Death
EBV + S maltophilia 8.9% (n=148)
MTB + RSV 8.3% (n=62)
HMPV + MTB 7.5% (n=53)
EBV + HSV 7.5% (n=68)
A baumannii + SARS-CoV-2 7.0% (n=116)
Rate (%)
C
Clinical outcome risk ratio across age groups
All ages
Aged < 5 years
Aged 5–17 years
Aged 18–60 years
Aged > 60 years
Pathogen pair
CP + HAdV
MP + RSV
HAdV + HMPV
H influenzae + M catarrhalis
MTB + RSV
MP + MTB
S pneumoniae + HRV
MTB + HRV
HMPV + HRV
CP + IFV
H influenzae + S aureus
MP + HAdV
MTB + IFV
HMPV + RSV
MP + HPIV
LRTI Hosp. ICU Death
Clinical outcome
log2(Clinical outcome risk ratio)

Figure 2