
High sequential-detection-rate pathogen pairs by age group
All ages
K pneumoniae → P aeruginosa
98.18%
S maltophilia → A baumannii
94.19%
A baumannii → S maltophilia
89.64%
IFV → Candida
78.99%
MP → Candida
72.07%
0 20 40 60 80 100
Children (<5 years)
IFV → H influenzae
MP → H influenzae
IFV → M tuberculosis
HCoV → SARS-CoV-2
SARS-CoV-2 → HCoV
0 5 10 15 20 25 30
Adolescents (5-17 years)
64.63%
62.59%
52.78%
50.83%
49.59%
0 20 40 60 80 100
Adults (18-60 years)
IFV → H influenzae
64.63%
MP → H influenzae
62.59%
IFV → M tuberculosis
52.78%
HCoV → SARS-CoV-2
50.83%
SARS-CoV-2 → HCoV
49.59%
0 20 40 60 80 100
Older adults (>60 years)
IFV → H influenzae
64.63%
MP → H influenzae
62.59%
IFV → M tuberculosis
52.78%
HCoV → SARS-CoV-2
50.83%
SARS-CoV-2 → HCoV
49.59%
0 20 40 60 80 100
Unknown
IFV → H influenzae
64.63%
MP → H influenzae
62.59%
IFV → M tuberculosis
52.78%
HCoV → SARS-CoV-2
50.83%
SARS-CoV-2 → HCoV
49.59%
0 20 40 60 80 100
Sequential detection rate: B(t+1)+ among A(