Directional sequential-detection associations

Pathogen A (preceding window)

CMV, SARS-CoV-2, EBV, HAdV, HBoV, HCoV, HMPV, HPIV, HRV, HSV, IFV, RSV, MP, CP, *B pertussis*, *H influenzae*, *K pneumoniae*, *Legionella*, *M catarrhalis*, *M tuberculosis*, *P aeruginosa*, *S aureus*, *S maltophilia*, *S pneumoniae*, *A baumannii*, Aspergillus, Candida, Cryptococcus, Pneumocystis

Pathogen B (subsequent window)

Pathogen type: Virus, Atypical, Bacteria, Fungi

log2(odds ratio): 6, 4, 2, 0, -2, -4, -6

Hospitalization

SARS-CoV-2 → *M tuberculosis*: * 0.65

A baumannii → *M tuberculosis*: * 0.66

IFV → *M tuberculosis*: * 0.87

S maltophilia → *M tuberculosis*: * 0.64

M tuberculosis → *H influenzae*: * 0.51

IFV → MP: * 0.84

S aureus → *M tuberculosis*: * 0.62

HAdV → MP: * 0.66

Risk ratio (95% CI): 0, 1, 2

ICU

IFV → EBV: * 4.25

Candida → *H influenzae*: * 0.11

IFV → MP: * 0.28

IFV → *H influenzae*: * 0.24

S pneumoniae → *M tuberculosis*: * 0.05

IFV → *M tuberculosis*: * 0.34

H influenzae → *M tuberculosis*: * 0.05

EBV → *H influenzae*: * 0.39

RSV → IFV: * 0.14

Risk ratio (95% CI): 0, 1, 2, 3