

Experimental design (respirationAdj1)

Trial design




Figure 1. Experimental design of respirationAdj1 (3 conditions; rhythmic fast = RhyF, rhythmic slow = RhyS, random slow = RanS). Estimated (est.) trial length equals the mean ITI, time window of the red fixation cross (1 s), and grand median detection and confidence RTs from Forster et al. (2025, Exp 2). Blocks consist of 80 trials (RhyS, RanS) or 160 trials (RhyF) to approximate similarity in duration. At the beginning of each block, starting periods are implemented that last about 30 s with ITIs matching the mean ITI of the respective other faster or slower condition (RhyF blocks: 6 trials with $t_{\text{Starting ITI}}$ of 3.5 s; RhyS and RanS blocks: 10 trials with $t_{\text{Starting ITI}}$ of 1 s). Excluding those starting period trials, the total trial count per condition is 320 trials (voltage symbol: 240 signal-present; empty set symbol: 80 signal-absent). Offline threshold assessments are used (and repeated prior to each block) to identify individual near-threshold intensities (mA) of approximately 60% detectability (similar procedure as in Forster et al., 2025; Grund et al., 2022). Near-threshold signals are delivered to the left index finger; button responses are given with the right index finger.

All time parameters (t) displayed in s. J/N equals Yes/No in German; S/U equals Confident/Unconfident in German.

Forster, C., Stephani, T., Grund, M., Panagoulas, E., Al, E., Hofmann, S. M., ... & Villringer, A. (2025). Pre-stimulus beta power mediates explicit and implicit perceptual biases in distinct cortical areas. *Communications Psychology*, 3(1), 93.

Grund, M., Al, E., Pabst, M., Dabbagh, A., Stephani, T., Nierhaus, T., ... & Villringer, A. (2022). Respiration, heartbeat, and conscious tactile perception. *Journal of Neuroscience*, 42(4), 643-656.