





1 QhX: A Python package for periodicity detection in red 2 noise

3 **Andjelka B. Kovačević** ^{1*}, **Dragana Ilić** ¹, **Momčilo Tošić**^{1*}, **Marina**
4 **Pavlović** ², **Aman Raju** ¹, **Mladen Nikolić** ¹, **Saša Simić** ³, **Iva Čvorović**
5 **Hajdinjak** ¹, and **Luka Č. Popović** ⁴

6 ¹ University of Belgrade-Faculty of Mathematics, Studentski trg 16, Belgrade, Serbia ² Mathematical
7 Institute of Serbian Academy of Science and Arts, Serbia ³ Faculty of sciences, University of Kragujevac,
8 Radoja Domanovića 12, Serbia ⁴ Astronomical Observatory, Belgrade, Serbia  Corresponding author *
9 These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright,
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

10 Summary

11 QhX is a Python-based package developed for periodicity detection in red noise time series data,
12 with a focus on the Vera C. Rubin Observatory Legacy Survey of Space and Time (LSST,
13 [Ivezić et al., 2019](#)). Traditional methods, such as those based on the Fourier Transform, often
14 struggle with red noise signals, which are prevalent in real-world datasets. The core of QhX is a
15 2D Hybrid method (cross-correlation of wavelet transforms of light curves, [Kovačević et al.,](#)
16 [2018](#)) that operates in a “period-period” phase space, capable of detecting oscillations in one
17 or more light curves. The QhX pipeline first transforms input data into a time-period plane
18 via wavelets and then (auto)correlates the resulting wavelet matrices to obtain a correlation
19 density in the period-period plane. After integrating the correlation density, the final decision
20 on detected periods is made by a statistical robovetter based on the significance, upper and
21 lower errors of detected periods, and the Intersection over Union (IoU) metric for measuring
22 the proximity and overlap of periods across bands. Beyond compiling the numerical catalog of
23 vetted periods, QhX offers visualization across photometric bands.

Statement of need




Figure 1: The left panel shows a 1D light curve with observational data (black error bars) and a model (blue line). QhX applies wavelet transforms to convert the time-series data (either observed or modeled) into the time-frequency domain, then cross-correlates wavelet transform matrices generating a correlation density map (right) in 2D period-period phase space where clusters reveal consistent periodic signals. After applying statistical procedures, the package generates a numerical catalog of flagged periodic objects (bottom left) and a dynamical view of detected periods across objects (bottom right).

Periodic variability can be encountered across a wide range of astronomical objects, from asteroids and stars to quasars. However, identifying meaningful signals in these variable sources is often complicated by red noise (see, e.g., Figure 1 in Gaia Collaboration et al., 2019; Kasliwal et al., 2015; Kovačević, Radović, et al., 2022), which shows fractal-like patterns across time scales (Belete et al., 2018; Vio et al., 1991). This, along with the non-stationary nature of signals and often unfavorable sampling (Brandt et al., 2018; D’Orazio & Charisi, 2023), makes identifying coherent signals challenging. Traditional time-frequency analysis, limited by the Fourier uncertainty principle (i.e., Gabor limit, Gabor, 1947), struggles with these complex signals, highlighting the need for a nonlinear approach (Abry et al., 1995; Cohen, 1995) for analyzing astronomical signals like quasar light curves.

QhX provides a robust framework with features specifically designed to address these challenges. The first feature of QhX is its core 2D Hybrid method (see Figure 1), detailed in (Kovačević et al., 2018), with an analogy to 2D Correlation Spectroscopy (Noda, 2018) discussed in Kovačević (2024). 2D Hybrid method enables a nonlinear approach (via cross-correlation), expanding detection into a two-dimensional period-period phase space. By applying wavelet transforms, QhX maps time-series data into a time-frequency domain, then (auto)correlates it to produce a correlation density in the period-period plane, enhancing detection in red noise-dominated data.

To further ensure robust detection, QhX uses an innovative Intersection over Union (IoU) metric alongside standard statistical measures (significance and upper and lower error bounds) to assess the overlap of detected periods across optical bands and objects. In addition to correlation density maps, each period is visualized as the center of an “IoU ball,” with a radius that represents relative error, calculated as the mean of the upper and lower error bounds—similar to a circular aperture in photometry (Saxena et al., 2024).

The third feature of QhX introduces a novel approach beyond traditional periodogram and wavelet transform plots by generating both numerical and interactive visual catalogs. These catalogs rank periodicity candidates by reliability, allowing for interactive inspection of signal consistency—a level of interpretability that static plots cannot achieve.

53 QhX structure




Figure 2: Schematic representation of the QhX package architecture.

The QhX package is a modular and extensible API (see Figure 2) designed for efficient detection and analysis of periodic signals in astronomical time series data, particularly for projects like LSST. Given that astronomical data analysis often requires rapid prototyping and experimentation with various algorithms, the modular design of QhX enables users to easily swap or modify functions, supporting diverse research needs without the constraints of a fixed class structure.

60 The package is organized into interconnected modules, each fulfilling a specialized role:

1. Core Algorithms:

- The algorithms module provides essential signal-processing techniques, including the Weighted Wavelet Z-Transform (wwtz) and prototype superlet transforms, which enable robust analysis of continuous time series data.

- 65 ▪ The correlation function within the `utils` module supports the 2D Hybrid method
66 by converting light curve data into wavelet matrices and performing (auto)correla-
67 tion, creating correlation density maps that highlight periodicity as main diagonal
68 clusters in the period-period phase space.
- 69 2. **Signal Detection and Validation:**
- 70 ▪ The detection module identifies candidate periodic signals and assesses their valid-
71 ity using statistical measures, including significance testing and error calculations,
72 informed by methods developed with the LSST community ([Johnson et al., 2019](#)).
73 The Intersection over Union (IoU) metric is applied to ensure robust detection
74 across bands and objects.
- 75 ▪ Robovetters, or statistical validation tools, finalize the reliability of detected periods,
76 adding an additional layer of quality control.
- 77 3. **Data Management:**
- 78 ▪ The `data_manager` and `data_manager_dynamical` modules manage data flow, han-
79 dling tasks like data loading, outlier removal, and format compatibility. They also
80 support custom data loaders to process various data formats.
- 81 ▪ The `batch_processor` and `parallelization_solver` modules optimize task dis-
82 tribution across multiple processors, boosting computational efficiency for large
83 datasets.
- 84 4. **Visualization and Output:**
- 85 ▪ The `plots` module includes tools for creating interactive visualizations, such as
86 `interactive_plot`, which allows for exploring detected periodicities across bands
87 and objects. For large datasets, `interactive_plot_large_files` enables in-depth
88 inspection of signal consistency.
- 89 ▪ The `output` and `output_parallel` modules handle result storage, supporting both
90 single-threaded and parallelized workflows.
- 91 5. **Testing and Validation:**
- 92 ▪ The `tests` module, with functions like `test_parallel` and `test_integrated`, vali-
93 dates the functionality of different components, ensuring robustness across various
94 processing setups.
- 95 QhX(Version 0.2.0) is a standalone, open-source package designed for handling datasets with
96 varying numbers of filters across surveys. It offers both dynamic and fixed modes, along
97 with parallel processing capabilities for large datasets. The `dynamical_mode.py` module offers
98 optional inclusion of observational errors, enhancing the accuracy of periodic signal detection
99 across multiple bands.

100 Representative Applications

101 The QhX method has been applied to various studies, including:

- 102 ▪ **Quasar Periodicity:** Investigating periodic signals in various quasars ([Fatović et al., 2023](#);
103 [Kovačević et al., 2018, 2019](#); [Kovačević, Popović, et al., 2020](#)).
- 104 ▪ **Quasi-Periodic Oscillations:** Analyzing oscillations in quasars ([Kovačević, Yi, et al.,](#)
105 [2020](#)).
- 106 ▪ **Very Low-Frequency Signals:** Detecting VLF signals variability before, during and after
107 earthquakes ([Kovačević, Nina, et al., 2022](#)).

108 Additionally, the QhX pipeline is listed as a [directable software in-kind contribution](#) to the LSST
109 project, highlighting its role in the LSST.

110 Documentation and Tutorials

111 Comprehensive documentation for QhX, available at [Github Pages](#), includes several example
112 notebooks:

- **Basic Tutorial:** Introduces the fundamentals of QhX using a mock light curve, helping new users get started quickly.
- **Parallel Processing Example:** Demonstrates how to perform parallel processing with quasar light curves from the [LSST AGN Data Challenge database](#), showcasing the software's capability to handle large datasets efficiently.
- **Task Distribution:** Showcases how to distribute tasks across multiple processors using, enhancing computational performance.
- **Merging Large Files:** Provides guidance on handling extensive datasets by merging large files, which is essential for high-volume data analysis.
- **Visualization of Large Datasets:** Illustrates how to visualize large files obtained from High-Performance Computing (HPC) environments, enabling effective interpretation of results.

Acknowledgements

Authors acknowledge funding provided by the University of Belgrade - Faculty of Mathematics (the contract 451-03-66/2024-03/200104), Faculty of Sciences University of Kragujevac (451-03-65/2024-03/200122), and Astronomical Observatory Belgrade (contract 451-03-66/2024-03/200002), through grants by the Ministry of Education, Science, and Technological Development of the Republic of Serbia) through the grants by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia.

References

- Abry, P., Gonçalves, P., & Flandrin, P. (1995). *Wavelets, spectrum analysis and 1/f processes* (pp. 15–29). Springer New York.
- Belete, A. B., Bravo, J. P., Canto Martins, B. L., Leo, I. C., De Araujo, J. M., & De Medeiros, J. R. (2018). Multifractality Signatures in Quasars Time Series – I. 3C 273. *Monthly Notices of the Royal Astronomical Society*, 478(3), 3976–3986.
- Brandt, W. N., Ni, Q., Yang, G., Anderson, S. F., Assef, R. J., Barth, A. J., Bauer, F. E., Bongiorno, A., Chen, C.-T., De Cicco, D., Gezari, S., Grier, C. J., Hall, P. B., Hoenig, S. F., Lacy, M., Li, J., Luo, B., Paolillo, M., Peterson, B. M., ... Yu, Z. (2018). Active Galaxy Science in the LSST Deep-Drilling Fields: Footprints, Cadence Requirements, and Total-Depth Requirements. *arXiv e-Prints*, arXiv:1811.06542. <https://doi.org/10.48550/arXiv.1811.06542>
- Cohen, L. (1995). *Time-frequency analysis*. Englewood Cliffs, NJ.
- D'Orazio, D. J., & Charisi, M. (2023). Observational Signatures of Supermassive Black Hole Binaries. *arXiv e-Prints*, arXiv:2310.16896. <https://doi.org/10.48550/arXiv.2310.16896>
- Fatović, M., Palaversa, L., Tisanić, K., Thanjavur, K., Ivezić, Ž., & Kovačević, A. B. et al. (2023). Detecting Long-period Variability in the SDSS Stripe 82 Standards Catalog. *Astrophysical Journal*, 165(4), 1–13.
- Gabor, D. (1947). Acoustical Quanta and the Theory of Hearing. *Nat*, 159(4044), 591–594.
- Gaia Collaboration, Eyer, L., Rimoldini, L., Audard, M., Anderson, R. I., Nienartowicz, K., Glass, F., Marchal, O., Grenon, M., Mowlavi, N., Holl, B., Clementini, G., Aerts, C., Mazeh, T., Evans, D. W., Szabados, L., Brown, A. G. A., Vallenari, A., Prusti, T., ... Zwitter, T. (2019). Gaia Data Release 2. Variable stars in the colour-absolute magnitude diagram. *Astronomy & Astrophysics*, 623, A110. <https://doi.org/10.1051/0004-6361/201833304>
- Ivezić, Ž., Kahn, S. M., Tyson, J. A., Abel, B., Acosta, E., & Allsman, R. et al. (2019). LSST: From Science Drivers to Reference Design and Anticipated Data Products. *Astrophysical*

- 158 *Journal*, 873(2), 111–151.
- 159 Johnson, M. A. C., Gandhi, P., Chapman, A. P., Moreau, L., Charles, P. A., Clarkson, W.
160 I., & Hill, A. B. (2019). Prospecting for periods with LSST - low-mass X-ray binaries
161 as a test case. *Monthly Notices of the Royal Astronomical Society*, 484(1), 19–30.
162 <https://doi.org/10.1093/mnras/sty3466>
- 163 Kasliwal, V. P., Vogeley, M. S., & Richards, G. T. (2015). Are the variability properties of the
164 Kepler AGN light curves consistent with a damped random walk? *Monthly Notices of the*
165 *Royal Astronomical Society*, 451(4), 4328–4345. <https://doi.org/10.1093/mnras/stv1230>
- 166 Kovačević, A. B. (2024). Two-dimensional (2D) hybrid method: Expanding 2D corre-
167 lation spectroscopy (2D-COS) for time series analysis. *Applied Spectroscopy*, 0(0),
168 00037028241241308. <https://doi.org/10.1177/00037028241241308>
- 169 Kovačević, A. B., Nina, A., Popović, L. Č., & Radovanović, M. (2022). Two-Dimensional
170 Correlation Analysis of Periodicity in Noisy Series: Case of VLF Signal Amplitude Variations
171 in the Time Vicinity of an Earthquake. *Mathematics*, 10(22), 1–14.
- 172 Kovačević, A. B., Pérez-Hernández, E., Popović, L. Č., Shapovalova, A. I., Kollatschny, W.,
173 & Ilić, D. (2018). Oscillatory Patterns in the Light curves of Five Long-Term Monitored
174 Type 1 Active Galactic Nuclei. *Monthly Notices of the Royal Astronomical Society*, 475(2),
175 2051–2066.
- 176 Kovačević, A. B., Popović, L. Č., & Ilić, D. (2020). Two-Dimensional Correlation Analysis of
177 Periodicity in Active Galactic Nuclei Time series. *Open Astronomy*, 29(1), 51–55.
- 178 Kovačević, A. B., Popović, L. Č., Simić, S., & Ilić, D. (2019). The Optical Variability of
179 Supermassive Black Hole Binary Candidate PG 1302-102: Periodicity and Perturbation in
180 the Light Curve. *Astrophysical Journal*, 871(1), 1–27.
- 181 Kovačević, A. B., Radović, V., Ilić, D., Popović, L. Č., Assef, R. J., Sánchez-Sáez, P.,
182 Nikutta, R., Raiteri, C. M., Yoon, I., Homayouni, Y., Li, Y.-R., Caplar, N., Czerny,
183 B., Panda, S., Ricci, C., Jankov, I., Landt, H., Wolf, C., Kovačević-Dojčinović, J., ...
184 Marčeta-Mandić, S. (2022). The LSST era of supermassive black hole accretion disk
185 reverberation mapping. *Astrophysical Journal Supplement Series*, 262(2), 49. <https://doi.org/10.3847/1538-4365/ac88ce>
- 186
187 Kovačević, A. B., Yi, T., Dai, X., Yang, X., Čvorović-Hajdinjak, I., & Popović, L. Č. (2020).
188 Confirmed Short Periodic Variability of Subparsec Supermassive Binary Black Hole Candi-
189 date Mrk 231. *Monthly Notices of the Royal Astronomical Society*, 494(3), 4069–4076.
- 190 Noda, I. (2018). Advances in Two-Dimensional Correlation Spectroscopy (2DCOS). In J. Laane
191 (Ed.), *Frontiers and Advances in Molecular Spectroscopy* (pp. 47–75). Elsevier.
- 192 Saxena, A., Salvato, M., Roster, W., Shirley, R., Buchner, J., Wolf, J., Kohl, C., Starck, H.,
193 Dwelly, T., Comparat, J., & al., et. (2024). CIRCLEZ : Reliable photometric redshifts for
194 active galactic nuclei computed solely using photometry from Legacy Survey Imaging for
195 DESI. 690, A365. <https://doi.org/10.1051/0004-6361/202450886>
- 196 Vio, R., Cristiani, S., Lessi, O., & Salvadori, L. (1991). 3C 345: Is the Variability of Quasars
197 Nonlinear? *Astrophysical Journal*, 380, 351–356.