





1 QhX: A Python package for periodicity detection in red 2 noise

3 **Andjelka B. Kovačević** ^{1*}, **Dragana Ilić** ¹, **Momčilo Tošić**^{1*}, **Marina**
4 **Pavlović** ², **Aman Raju** ¹, **Mladen Nikolić** ¹, **Saša Simić** ³, **Iva Čvorović**
5 **Hajdinjak** ¹, and **Luka Č. Popović** ⁴

6 **1** University of Belgrade-Faculty of Mathematics, Studentski trg 16, Belgrade, Serbia **2** Mathematical
7 Institute of Serbian Academy of Science and Arts, Serbia **3** Faculty of sciences, University of Kragujevac,
8 Radoja Domanovića 12, Serbia **4** Astronomical Observatory, Belgrade, Serbia  Corresponding author *
9 These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#)).

10 Summary

11 QhX is a Python package for detecting periodicity in red noise time series, developed as an
12 in-kind contribution to the Vera C. Rubin Observatory Legacy Survey of Space and Time
13 (LSST, [Ivezić et al., 2019](#)). Traditional Fourier-based methods often struggle with red noise,
14 which is common in quasar light curves and other accreting objects. QhX addresses these
15 challenges with its core 2D Hybrid method ([Kovačević et al., 2018](#)). Input data are mapped
16 into a time-period plane via wavelet transforms, which are (auto)correlated to produce a
17 correlation density map in a “period-period” plane. Statistical vetting incorporates significance,
18 upper and lower error bounds, and the novel Intersection over Union (IoU) metric to evaluate
19 the proximity and overlap of detected periods across bands and objects. In addition to a vetted
20 numerical catalog, QhX dynamically visualizes periodicity across photometric bands and objects.

Statement of need




Figure 1: The left panel shows a 1D light curve with observational data (black error bars) and a model (blue line). QhX transforms the time-series into the time-frequency domain and cross-correlates wavelet matrices to produce a 2D period-period correlation map (right), where clusters indicate periodic signals. After map integration, statistical vetting generates a numerical catalog of flagged periodic objects (bottom left) and a dynamic view of detected periods across objects and bands (bottom right).

Periodic variability spans a range of astronomical objects, from asteroids to quasars. Identifying meaningful signals is complicated by red noise (see, e.g., Figure 1 in Gaia Collaboration et al., 2019; Kasliwal et al., 2015; Kovačević, Radović, et al., 2022), which exhibits fractal-like patterns across time scales (Belete et al., 2018; Vio et al., 1991). Non-stationary signals and unfavorable sampling (Brandt et al., 2018; D’Orazio & Charisi, 2023) further obscure coherent patterns. Traditional time-frequency methods, constrained by the Fourier uncertainty principle (i.e., Gabor limit, Gabor, 1947), often fail with such complex signals, highlighting the need for nonlinear approaches (Abry et al., 1995; Cohen, 1995).

QhX provides features specifically designed to address these challenges. The first feature is its core 2D Hybrid method (see Figure 1), detailed in (Kovačević et al., 2018), inspired by 2D Correlation Spectroscopy (Kovačević, 2024; Noda, 2018). By applying wavelet transforms, QhX maps time-series data into the time-frequency domain and (auto)correlates it, generating a period-period correlation density that enhances signal detection. Secondly, QhX introduces an Intersection over Union (IoU) metric, combined with standard statistical measures (significance, upper and lower error bounds), to evaluate the overlap of detected periods across bands and objects. Each period is represented as the center of an “IoU ball,” with its radius reflecting relative error, calculated as the mean of the upper and lower error bounds—analogous to a circular aperture in photometry (Saxena et al., 2024). Thirdly, QhX enhances traditional analysis by generating numerical and interactive visual catalogs that rank periodicity candidates by reliability. These interactive catalogs enable detailed inspection of signal consistency, offering greater interpretability than traditional static plots.

43 QhX structure




Figure 2: Schematic representation of the QhX package architecture.

44 QhX (Version 0.2.0) is an open-source package optimized for gappy quasar light curves but
 45 adaptable to other datasets. It supports both dynamic and fixed modes, with parallel processing
 46 capabilities for large-scale data. The modular design facilitates rapid experimentation by
 47 enabling easy swapping or modification of functions (see Figure 2), addressing diverse research
 48 needs. For fixed-only workflows, specialized functions such as `data_manager` offer minimal
 49 overhead and optimal performance, while `data_manager_dynamical` supports both dynamic
 50 and fixed configurations to handle more complex scenarios involving dynamic filters.

51 The package is organized as follows:

52 1. Core:

- 53 ■ `algorithms` module provides essential signal-processing techniques, including the
- 54 Weighted Wavelet Z-Transform (`wwtz`) and prototype `superlet` transforms.
- 55 ■ `correlation` function within the `utils` module supports the 2D Hybrid method by
- 56 converting light curve data into wavelet matrices and performing (auto)correlation,
- 57 creating correlation density.

58 2. Signal Detection and Validation:

- 59 ■ `detection` module identifies candidate periodic signals and assesses their validity
- 60 using statistical measures (significance and upper and lower error (Johnson et al.,

- 61 2019)). The Intersection over Union (IoU) metric identifies overlapping periods
62 across bands and objects.
- 63 ■ Statistical vetting categorizes detected periods for each object and band as reliable,
64 medium, or poor.
- 65 3. **Data Management:**
- 66 ■ `data_manager` and `data_manager_dynamical` modules manage data flow, data
67 loading, outlier removal, and format compatibility.
 - 68 ■ `batch_processor` and `parallelization_solver` modules optimize task distribution
69 across multiple processors.
- 70 4. **Visualization and Output:**
- 71 ■ `plots` module includes tools for creating interactive visualizations, such as
72 `interactive_plot`, which allows for exploring detected periodicities across bands
73 and objects. For large datasets, `interactive_plot_large_files` enables in-depth
74 inspection of signal consistency.
 - 75 ■ `output` and `output_parallel` modules handle result storage, supporting both
76 single-threaded and parallelized workflows.
- 77 5. **Testing:**
- 78 ■ `tests` module, containing `test_parallel` and `test_integrated`, validates the
79 functionality across various processing setups.

80 Representative Applications

81 The QhX method has been applied to:

- 82 -Quasar periodicity detection (Fatović et al., 2023; Kovačević et al., 2018, 2019; Kovačević,
83 Popović, et al., 2020).
- 84 -Quasi-Periodic Oscillations detection (Kovačević, Yi, et al., 2020).
- 85 -Very Low-Frequency (VLF) signals variability in the vicinity of earthquakes (Kovačević, Nina,
et al., 2022).
- QhX is the [LSST directable software in-kind contribution](#).

86 Acknowledgements

87 Funding was provided by the University of Belgrade - Faculty of Mathematics (the contract
88 451-03-66/2024-03/200104), Faculty of Sciences University of Kragujevac (451-03-65/2024-
89 03/200122), and Astronomical Observatory Belgrade (contract 451-03-66/2024- 03/200002),
90 through grants by the Ministry of Education, Science, and Technological Development of the
91 Republic of Serbia.

92 References

- 93 Abry, P., Gonçalves, P., & Flandrin, P. (1995). *Wavelets, spectrum analysis and 1/f processes*
94 (pp. 15–29). Springer New York.
- 95 Belete, A. B., Bravo, J. P., Canto Martins, B. L., Leo, I. C., De Araujo, J. M., & De Medeiros,
96 J. R. (2018). Multifractality Signatures in Quasars Time Series – I. 3C 273. *Monthly*
97 *Notices of the Royal Astronomical Society*, 478(3), 3976–3986.
- 98 Brandt, W. N., Ni, Q., Yang, G., Anderson, S. F., Assef, R. J., Barth, A. J., Bauer, F. E.,
99 Bongiorno, A., Chen, C.-T., De Cicco, D., Gezari, S., Grier, C. J., Hall, P. B., Hoenig,
100 S. F., Lacy, M., Li, J., Luo, B., Paolillo, M., Peterson, B. M., ... Yu, Z. (2018). Active
101 Galaxy Science in the LSST Deep-Drilling Fields: Footprints, Cadence Requirements, and
102 Total-Depth Requirements. *arXiv e-Prints*, arXiv:1811.06542. [https://doi.org/10.48550/
103 arXiv.1811.06542](https://doi.org/10.48550/arXiv.1811.06542)
- 104 Cohen, L. (1995). *Time-frequency analysis*. Englewood Cliffs, NJ.

- 105 D'Orazio, D. J., & Charisi, M. (2023). Observational Signatures of Supermassive Black Hole
106 Binaries. *arXiv e-Prints*, arXiv:2310.16896. <https://doi.org/10.48550/arXiv.2310.16896>
- 107 Fatović, M., Palaversa, L., Tisanić, K., Thanjavur, K., Ivezić, Ž., & Kovačević, A. B. et
108 al. (2023). Detecting Long-period Variability in the SDSS Stripe 82 Standards Catalog.
109 *Astrophysical Journal*, 165(4), 1–13.
- 110 Gabor, D. (1947). Acoustical Quanta and the Theory of Hearing. *Nat*, 159(4044), 591–594.
- 111 Gaia Collaboration, Eyer, L., Rimoldini, L., Audard, M., Anderson, R. I., Nienartowicz, K.,
112 Glass, F., Marchal, O., Grenon, M., Mowlavi, N., Holl, B., Clementini, G., Aerts, C., Mazeh,
113 T., Evans, D. W., Szabados, L., Brown, A. G. A., Vallenari, A., Prusti, T., ... Zwitter, T.
114 (2019). Gaia Data Release 2. Variable stars in the colour-absolute magnitude diagram.
115 *Astronomy & Astrophysics*, 623, A110. <https://doi.org/10.1051/0004-6361/201833304>
- 116 Ivezić, Ž., Kahn, S. M., Tyson, J. A., Abel, B., Acosta, E., & Allsman, R. et al. (2019). LSST:
117 From Science Drivers to Reference Design and Anticipated Data Products. *Astrophysical*
118 *Journal*, 873(2), 111–151.
- 119 Johnson, M. A. C., Gandhi, P., Chapman, A. P., Moreau, L., Charles, P. A., Clarkson, W.
120 I., & Hill, A. B. (2019). Prospecting for periods with LSST - low-mass X-ray binaries
121 as a test case. *Monthly Notices of the Royal Astronomical Society*, 484(1), 19–30.
122 <https://doi.org/10.1093/mnras/sty3466>
- 123 Kasliwal, V. P., Vogeley, M. S., & Richards, G. T. (2015). Are the variability properties of the
124 Kepler AGN light curves consistent with a damped random walk? *Monthly Notices of the*
125 *Royal Astronomical Society*, 451(4), 4328–4345. <https://doi.org/10.1093/mnras/stv1230>
- 126 Kovačević, A. B. (2024). Two-dimensional (2D) hybrid method: Expanding 2D corre-
127 lation spectroscopy (2D-COS) for time series analysis. *Applied Spectroscopy*, 0(0),
128 00037028241241308. <https://doi.org/10.1177/00037028241241308>
- 129 Kovačević, A. B., Nina, A., Popović, L. Č., & Radovanović, M. (2022). Two-Dimensional
130 Correlation Analysis of Periodicity in Noisy Series: Case of VLF Signal Amplitude Variations
131 in the Time Vicinity of an Earthquake. *Mathematics*, 10(22), 1–14.
- 132 Kovačević, A. B., Pérez-Hernández, E., Popović, L. Č., Shapovalova, A. I., Kollatschny, W.,
133 & Ilić, D. (2018). Oscillatory Patterns in the Light curves of Five Long-Term Monitored
134 Type 1 Active Galactic Nuclei. *Monthly Notices of the Royal Astronomical Society*, 475(2),
135 2051–2066.
- 136 Kovačević, A. B., Popović, L. Č., & Ilić, D. (2020). Two-Dimensional Correlation Analysis of
137 Periodicity in Active Galactic Nuclei Time series. *Open Astronomy*, 29(1), 51–55.
- 138 Kovačević, A. B., Popović, L. Č., Simić, S., & Ilić, D. (2019). The Optical Variability of
139 Supermassive Black Hole Binary Candidate PG 1302-102: Periodicity and Perturbation in
140 the Light Curve. *Astrophysical Journal*, 871(1), 1–27.
- 141 Kovačević, A. B., Radović, V., Ilić, D., Popović, L. Č., Assef, R. J., Sánchez-Sáez, P.,
142 Nikutta, R., Raiteri, C. M., Yoon, I., Homayouni, Y., Li, Y.-R., Caplar, N., Czerny,
143 B., Panda, S., Ricci, C., Jankov, I., Landt, H., Wolf, C., Kovačević-Dojčinović, J., ...
144 Marčeta-Mandić, S. (2022). The LSST era of supermassive black hole accretion disk
145 reverberation mapping. *Astrophysical Journal Supplement Series*, 262(2), 49. <https://doi.org/10.3847/1538-4365/ac88ce>
- 146
- 147 Kovačević, A. B., Yi, T., Dai, X., Yang, X., Čvorović-Hajdinjak, I., & Popović, L. Č. (2020).
148 Confirmed Short Periodic Variability of Subparsec Supermassive Binary Black Hole Candi-
149 date Mrk 231. *Monthly Notices of the Royal Astronomical Society*, 494(3), 4069–4076.
- 150 Noda, I. (2018). Advances in Two-Dimensional Correlation Spectroscopy (2DCOS). In J. Laane
151 (Ed.), *Frontiers and Advances in Molecular Spectroscopy* (pp. 47–75). Elsevier.

- 152 Saxena, A., Salvato, M., Roster, W., Shirley, R., Buchner, J., Wolf, J., Kohl, C., Starck, H.,
153 Dwelly, T., Comparat, J., & al., et. (2024). CIRCLEZ : Reliable photometric redshifts for
154 active galactic nuclei computed solely using photometry from Legacy Survey Imaging for
155 DESI. *690*, A365. <https://doi.org/10.1051/0004-6361/202450886>
- 156 Vio, R., Cristiani, S., Lessi, O., & Salvadori, L. (1991). 3C 345: Is the Variability of Quasars
157 Nonlinear? *Astrophysical Journal*, *380*, 351–356.

DRAFT