





QhX: A Python package for periodicity detection in red noise

Andjelka B. Kovačević^{1*}, Dragana Ilić¹, Momčilo Tošić^{1*}, Marina Pavlović², Aman Raju¹, Mladen Nikolić¹, Saša Simić³, Iva Čvorović Hajdinjak¹, and Luka Č. Popović⁴

¹ University of Belgrade-Faculty of Mathematics, Studentski trg 16, Belgrade, Serbia ² Mathematical Institute of Serbian Academy of Science and Arts, Serbia ³ Faculty of sciences, University of Kragujevac, Radoja Domanovića 12, Serbia ⁴ Astronomical Observatory, Belgrade, Serbia  Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Summary

QhX is a Python package for detecting periodicity in red noise time series, developed as an in-kind contribution to the Vera C. Rubin Observatory Legacy Survey of Space and Time (LSST, [Ivezić et al., 2019](#)). Traditional Fourier-based methods often struggle with red noise, which is common in quasar light curves and other accreting objects. QhX addresses these challenges with its core 2D Hybrid method ([Kovačević et al., 2018](#)). Input data are mapped into a time-period plane via wavelet transforms, which are (auto)correlated to produce a correlation density map in a “period-period” plane. Statistical vetting incorporates significance, upper and lower error bounds, and the novel Intersection over Union (IoU) metric to evaluate the proximity and overlap of detected periods across bands and objects. In addition to a vetted numerical catalog, QhX dynamically visualizes periodicity across photometric bands and objects.

Statement of need




Figure 1: The left panel shows a 1D light curve with observational data (black error bars) and a model (blue line). QhX transforms the time-series into the time-frequency domain and cross-correlates wavelet matrices to produce a 2D period-period correlation map (right), where clusters indicate periodic signals. After map integration, statistical vetting generates a numerical catalog of flagged periodic objects (bottom left) and a dynamic view of detected periods across objects and bands (bottom right).

Periodic variability spans a range of astronomical objects, from asteroids to quasars. Identifying meaningful signals is complicated by red noise (see, e.g., Figure 1 in [Gaia Collaboration et al., 2019](#); [Kasliwal et al., 2015](#); [Kovačević, Radović, et al., 2022](#)), which exhibits fractal-like patterns across time scales ([Belete et al., 2018](#); [Vio et al., 1991](#)). Non-stationary signals and unfavorable sampling ([Brandt et al., 2018](#); [D’Orazio & Charisi, 2023](#)) further obscure coherent patterns. Traditional time-frequency methods, constrained by the Fourier uncertainty principle (i.e., Gabor limit, [Gabor, 1947](#)), often fail with such complex signals, highlighting the need for nonlinear approaches ([Abry et al., 1995](#); [Cohen, 1995](#)).

QhX provides features specifically designed to address these challenges. The first feature is its core 2D Hybrid method (see Figure 1), detailed in ([Kovačević et al., 2018](#)), inspired by 2D Correlation Spectroscopy ([Kovačević, 2024](#); [Noda, 2018](#)). By applying wavelet transforms, QhX maps time-series data into the time-frequency domain and (auto)correlates it, generating a period-period correlation density that enhances signal detection. Secondly, QhX introduces an Intersection over Union (IoU) metric, combined with standard statistical measures (significance, upper and lower error bounds), to evaluate the overlap of detected periods across bands and objects. Each period is represented as the center of an “IoU ball,” with its radius reflecting relative error, calculated as the mean of the upper and lower error bounds—analogous to a circular aperture in photometry ([Saxena et al., 2024](#)). Thirdly, QhX enhances traditional analysis by generating numerical and interactive visual catalogs that rank periodicity candidates by reliability. These interactive catalogs enable detailed inspection of signal consistency, offering greater interpretability than traditional static plots.

43 QhX structure




Figure 2: Schematic representation of the QhX package architecture.

44 QhX (Version 0.2.0) is an open-source package optimized for gappy quasar light curves but
 45 adaptable to other datasets. It supports both dynamic and fixed modes, with parallel processing
 46 capabilities for large-scale data. The modular design facilitates rapid experimentation by
 47 enabling easy swapping or modification of functions (see Figure 2), addressing diverse research
 48 needs. For fixed-only workflows, specialized functions such as `data_manager` offer minimal
 49 overhead and optimal performance, while `data_manager_dynamical` supports both dynamic
 50 and fixed configurations to handle more complex scenarios involving dynamic filters.

51 The package is organized as follows:

52 1. Core:

- 53 ■ `algorithms` module provides essential signal-processing techniques, including the
- 54 Weighted Wavelet Z-Transform (`wwtz`) and prototype `superlet` transforms.
- 55 ■ `correlation` function within the `utils` module supports the 2D Hybrid method by
- 56 converting light curve data into wavelet matrices and performing (auto)correlation,
- 57 creating correlation density.

58 2. Signal Detection and Validation:

- 59 ■ `detection` module identifies candidate periodic signals and assesses their validity
- 60 using statistical measures (significance and upper and lower error (Johnson et al.,

- 2019)). The Intersection over Union (IoU) metric identifies overlapping periods across bands and objects.
- Statistical vetting categorizes detected periods for each object and band as reliable, medium, or poor.
3. **Data Management:**
- `data_manager` and `data_manager_dynamical` modules manage data flow, data loading, outlier removal, and format compatibility.
 - `batch_processor` and `parallelization_solver` modules optimize task distribution across multiple processors.
4. **Visualization and Output:**
- `plots` module includes tools for creating interactive visualizations, such as `interactive_plot`, which allows for exploring detected periodicities across bands and objects. For large datasets, `interactive_plot_large_files` enables in-depth inspection of signal consistency.
 - `output` and `output_parallel` modules handle result storage, supporting both single-threaded and parallelized workflows.
5. **Testing:**
- `tests` module, containing `test_parallel` and `test_integrated`, validates the functionality across various processing setups.

Representative Applications

The QhX method has been applied to:

- Quasar periodicity detection (Fatović et al., 2023; Kovačević et al., 2018, 2019; Kovačević, Popović, et al., 2020).
- Quasi-Periodic Oscillations detection (Kovačević, Yi, et al., 2020).
- Very Low-Frequency (VLF) signals variability in the vicinity of earthquakes (Kovačević, Nina, et al., 2022).
- QhX is the [LSST directable software in-kind contribution](#).

Acknowledgements

Funding was provided by the University of Belgrade - Faculty of Mathematics (the contract 451-03-66/2024-03/200104), Faculty of Sciences University of Kragujevac (451-03-65/2024-03/200122), and Astronomical Observatory Belgrade (contract 451-03-66/2024-03/200002), through grants by the Ministry of Education, Science, and Technological Development of the Republic of Serbia.

References

- Abry, P., Gonçalves, P., & Flandrin, P. (1995). *Wavelets, spectrum analysis and 1/f processes* (A. Antoniadis & G. Oppenheim, Eds.; pp. 15–29). Springer New York. https://doi.org/10.1007/978-1-4612-2544-7_2
- Belete, A. B., Bravo, J. P., Canto Martins, B. L., Leão, I. C., De Araujo, J. M., & De Medeiros, J. R. (2018). Multifractality signatures in quasars time series - I. 3C 273. 478(3), 3976–3986. <https://doi.org/10.1093/mnras/sty1316>
- Brandt, W. N., Ni, Q., Yang, G., Anderson, S. F., Assef, R. J., Barth, A. J., Bauer, F. E., Bongiorno, A., Chen, C.-T., De Cicco, D., Gezari, S., Grier, C. J., Hall, P. B., Hoenig, S. F., Lacy, M., Li, J., Luo, B., Paolillo, M., Peterson, B. M., ... Yu, Z. (2018). Active Galaxy Science in the LSST Deep-Drilling Fields: Footprints, Cadence Requirements, and Total-Depth Requirements. *arXiv e-Prints*, arXiv:1811.06542. <https://doi.org/10.48550/arXiv.1811.06542>
- Cohen, L. (1995). *Time-frequency analysis*. Prentice Hall PTR. ISBN: 9780135945322

- 106 D'Orazio, D. J., & Charisi, M. (2023). Observational Signatures of Supermassive Black Hole
107 Binaries. *arXiv e-Prints*, arXiv:2310.16896. <https://doi.org/10.48550/arXiv.2310.16896>
- 108 Fatović, M., Palaversa, L., Tisanić, K., Thanjavur, K., Ivezić, Ž., Kovačević, A. B., Ilić, D., &
109 Č. Popović, L. (2023). Detecting Long-period Variability in the SDSS Stripe 82 Standards
110 Catalog. *165*(4), 138. <https://doi.org/10.3847/1538-3881/acb596>
- 111 Gabor, D. (1947). Acoustical quanta and the theory of hearing. *Nature*, *159*(4044), 591–594.
112 <https://doi.org/10.1038/159591a0>
- 113 Gaia Collaboration, Eyer, L., Rimoldini, L., Audard, M., Anderson, R. I., Nienartowicz, K.,
114 Glass, F., Marchal, O., Grenon, M., Mowlavi, N., Holl, B., Clementini, G., Aerts, C., Mazeh,
115 T., Evans, D. W., Szabados, L., Brown, A. G. A., Vallenari, A., Prusti, T., ... Zwitter, T.
116 (2019). Gaia Data Release 2. Variable stars in the colour-absolute magnitude diagram.
117 *Astronomy & Astrophysics*, *623*, A110. <https://doi.org/10.1051/0004-6361/201833304>
- 118 Ivezić, Ž., Kahn, S. M., Tyson, J. A., Abel, B., Acosta, E., Allsman, R., Alonso, D., AlSayyad,
119 Y., Anderson, S. F., Andrew, J., Angel, J. R. P., Angeli, G. Z., Ansari, R., Antilogus, P.,
120 Araujo, C., Armstrong, R., Arndt, K. T., Astier, P., Aubourg, É., ... Zhan, H. (2019). LSST:
121 From Science Drivers to Reference Design and Anticipated Data Products. *873*(2), 111.
122 <https://doi.org/10.3847/1538-4357/ab042c>
- 123 Johnson, M. A. C., Gandhi, P., Chapman, A. P., Moreau, L., Charles, P. A., Clarkson, W.
124 I., & Hill, A. B. (2019). Prospecting for periods with LSST - low-mass X-ray binaries
125 as a test case. *Monthly Notices of the Royal Astronomical Society*, *484*(1), 19–30.
126 <https://doi.org/10.1093/mnras/sty3466>
- 127 Kasliwal, V. P., Vogeley, M. S., & Richards, G. T. (2015). Are the variability properties of the
128 Kepler AGN light curves consistent with a damped random walk? *Monthly Notices of the*
129 *Royal Astronomical Society*, *451*(4), 4328–4345. <https://doi.org/10.1093/mnras/stv1230>
- 130 Kovačević, A. B. (2024). Two-dimensional (2D) hybrid method: Expanding 2D corre-
131 lation spectroscopy (2D-COS) for time series analysis. *Applied Spectroscopy*, *0*(0),
132 00037028241241308. <https://doi.org/10.1177/00037028241241308>
- 133 Kovačević, A. B., Nina, A., Popović, L. Č., & Radovanović, M. (2022). Two-dimensional
134 correlation analysis of periodicity in noisy series: Case of VLF signal amplitude variations
135 in the time vicinity of an earthquake. *Mathematics*, *10*(22). <https://doi.org/10.3390/math10224278>
- 136
137 Kovačević, A. B., Pérez-Hernández, E., Popović, L. Č., Shapovalova, A. I., Kollatschny, W., &
138 Ilić, D. (2018). Oscillatory patterns in the light curves of five long-term monitored type 1
139 active galactic nuclei. *475*(2), 2051–2066. <https://doi.org/10.1093/mnras/stx3137>
- 140 Kovačević, A. B., Popović, L. Č., & Ilić, D. (2020). Two-dimensional correlation analysis of
141 periodicity in active galactic nuclei time series. *Open Astronomy*, *29*(1), 51–55. <https://doi.org/10.1515/astro-2020-0007>
- 142
143 Kovačević, A. B., Popović, L. Č., Simić, S., & Ilić, D. (2019). The Optical Variability of
144 Supermassive Black Hole Binary Candidate PG 1302-102: Periodicity and Perturbation in
145 the Light Curve. *871*(1), 32. <https://doi.org/10.3847/1538-4357/aaf731>
- 146 Kovačević, A. B., Radović, V., Ilić, D., Popović, L. Č., Assef, R. J., Sánchez-Sáez, P., Nikutta,
147 R., Raiteri, C. M., Yoon, I., Homayouni, Y., Li, Y.-R., Caplar, N., Czerny, B., Panda, S.,
148 Ricci, C., Jankov, I., Landt, H., Wolf, C., Kovačević-Dojčinović, J., ... Marčeta-Mandić, S.
149 (2022). The LSST Era of Supermassive Black Hole Accretion Disk Reverberation Mapping.
150 *262*(2), 49. <https://doi.org/10.3847/1538-4365/ac88ce>
- 151 Kovačević, A. B., Yi, T., Dai, X., Yang, X., Čvorović-Hajdinjak, I., & Popović, L. Č. (2020).
152 Confirmed short periodic variability of subparsec supermassive binary black hole candidate
153 Mrk 231. *494*(3), 4069–4076. <https://doi.org/10.1093/mnras/staa737>

- 154 Noda, I. (2018). Chapter 2 - advances in two-dimensional correlation spectroscopy (2DCOS).
155 In J. Laane (Ed.), *Frontiers and advances in molecular spectroscopy* (pp. 47–75). Elsevier.
156 <https://doi.org/https://doi.org/10.1016/B978-0-12-811220-5.00002-2>
- 157 Saxena, A., Salvato, M., Roster, W., Shirley, R., Buchner, J., Wolf, J., Kohl, C., Starck, H.,
158 Dwelly, T., Comparat, J., & al., et. (2024). CIRCLEZ : Reliable photometric redshifts for
159 active galactic nuclei computed solely using photometry from Legacy Survey Imaging for
160 DESI. *690*, A365. <https://doi.org/10.1051/0004-6361/202450886>
- 161 Vio, R., Cristiani, S., Lessi, O., & Salvadori, L. (1991). 3C 345: Is the Variability of Quasars
162 Nonlinear? *380*, 351. <https://doi.org/10.1086/170594>

DRAFT