

Spectral analyses of digital deterministic signals

Signal processing

Dell'Aria Dorian

Goffaux Lionel


October 7, 2020

Academic Year 2020-2021

Bachelor's degree in Computer Science

1 Influence of the Parameter N

Question 1




As expected, on the first graph, the maximum value is one and the minimum value is minus one corresponding to the amplitude=1 of the signal. The sine wave also achieves 6 periods in 10 ms meaning that its frequency is $1000 \frac{6}{10} = 600\text{Hz}$. On the second graph, we observe a spike at $\phi \approx 0.376$ which after computation gives us a frequency of $f = 600\text{Hz}$.

2 Influence of the Sampling Frequency

Question 2

For $f = 600\text{Hz}$ and $f = 4400\text{Hz}$, we see that the Fourier transform correspond to the reality, because $2f \leq Fe$ (Shannon's theorem). For $f = 5000\text{Hz}$, samples represents the sum of the errors. We should see a graph with 0 everywhere if errors were equals to 0, because $2f = Fe$. For $f = 5600\text{Hz}$ and $f = 10600\text{Hz}$, we observe a stroboscopic effect because $2f > Fe$. So the Fourier transform doesn't correspond to the reality.




3 Influence of the Parameter NTFD

Question 3

1. (a) The Position of the Fourier spike is at $2\pi * \frac{f}{F_e} = 1.66$.
 (b) The maximum of the Fourier spike is equal to $\frac{N}{2} = 16$.
 (c) There are two frequency samples on each secondary lobe.
2. (a) The position of the Fourier spike is at $2\pi * \frac{f}{F_e} = 1.66$.
 (b) The maximum of the Fourier spike is equal to $\frac{N}{2} = 16$.
 (c) There are sixteen frequency samples on each secondary lobe.
3. (a) The position of the Fourier spike is at $2\pi * \frac{f}{F_e} = 1.57$.
 (b) The maximum of the Fourier spike is equal to $\frac{N}{2} = 16$.
 (c) There are two frequency samples on each secondary lobe.
4. (a) The position of the Fourier spike is at $2\pi * \frac{f}{F_e} = 1.57$.

- (b) The maximum of the Fourier spike is equal to $\frac{N}{2} = 16$.
- (c) There are sixteen frequency samples on each secondary lobe.




4 Spectral Analysis of Several Sine Waves

Question 4


We find that the closer the sample frequency is to twice the higher frequency, the closer the spikes are to π . So in order to increase the accuracy, it's better to set the sample rate as close as possible to two times the higher frequency that we want to observe. For the NTFD parameter, that only change the space between the frequency samples in the spectrum. So, the higher it is the most continuous the plot seems. We choose $N = 45$ because N must be greater than $2d$ where d is difference between the two closest normalized frequency

to avoid lobes overlapping. For the number of samples in the time's space, we observe that the shorter the signal lasts in time the larger the spike corresponding to that signal frequency is and vice versa. There is probably a link with the uncertainty principle in quantum mechanics.



5 Weighting Window Effect

Question 5



We observe that the main lobe is getting larger and larger and the other lobes are getting lower and lower from one window to another. That fits well with the given table.

Question 6

For the first graph, we observe that the rectangular window is the best because the size of the main lobe is smaller than the other ones. For the second graph the difference between the two frequency is bigger, so the main lobe of the smallest frequency is drowned into the secondary lobes of the biggest one. So we choose the Blackman window because the secondary lobes of this window are smaller.

