

# Weekly Meeting

Qi Li, qi.li@cern.ch

(IHEP, CAS)

June 1, 2016



中国科学院高能物理研究所

Institute of High Energy Physics Chinese Academy of Sciences

# Introduction

# Object Definition

- photons
- Muons
- Electrons
- Jets
- Missing ET
- Overlap Removal

Begin with  $h \rightarrow \gamma\gamma$  selections. The additional cuts follow as:

- At least one lepton
- At least two jets
- B-veto
- Tight mass window
- Missing ET significance

# Event Yields

All cuts	SM Higgs pair
all	100.0%
Trigger	73.9%
GRL	73.9%
Detector Quality	73.9%
has PV	73.9%
2 loose photons	60.3%
$e - \gamma$ ambiguity	59.8%
Tight ID	50.4%
Isolation	44.7%
Rel.Pt cuts	40.9%
$105 < m_{\gamma\gamma} < 160$ GeV	40.7%
At least 2 jets	34.6%
At least 1 lepton	16.1%
b-veto	14.1%
Tight mass window	11.4%
MET Significance	9.9%

**Table:** Cut efficiencies for non-resonance

Signal yields = 0.15

# Background Yields

Channel	ggh	VBF	Wh	Zh	tth
Events yields	negligible	negligible	0.078	0.018	0.054
Run 1 results	negligible	negligible	0.14	0.025	0.08

**Table:** Event yields for SM Higgs productions

- The Continuum bkg estimated from sideband:  
 $N^{\text{continuum}} = 0.79$

# Sensitivities

$$Z = 0.15$$

# Summary



