

VU Research Portal

The effects of psychotherapy for depression on anxiety symptoms: a meta-analysis

Weitz, Erica; Kleiboer, Annet; van Straten, Annemieke; Cuijpers, Pim

published in

Psychological Medicine

2018

DOI (link to publisher)

[10.1017/S0033291717003622](https://doi.org/10.1017/S0033291717003622)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Weitz, E., Kleiboer, A., van Straten, A., & Cuijpers, P. (2018). The effects of psychotherapy for depression on anxiety symptoms: a meta-analysis. *Psychological Medicine*, 48(13), 2140-2152.
<https://doi.org/10.1017/S0033291717003622>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Original Article

Cite this article: Weitz E, Kleiboer A, van Straten A, Cuijpers P (2018). The effects of psychotherapy for depression on anxiety symptoms: a meta-analysis. *Psychological Medicine* **48**, 2140–2152. <https://doi.org/10.1017/S0033291717003622>

Received: 1 June 2017
Revised: 14 November 2017
Accepted: 14 November 2017
First published online: 24 January 2018

Key words:

Anxiety; depression; psychotherapy; meta-analysis

Author for correspondence:

Erica Weitz, E-mail: e.weitz@vu.nl

The effects of psychotherapy for depression on anxiety symptoms: a meta-analysis

Erica Weitz, Annet Kleiboer, Annemieke van Straten and Pim Cuijpers

Department of Clinical, Neuro and Developmental Psychology, EMGO Institute for Health and Care Research, Vrije Universiteit Amsterdam, Van der Boerhorststraat 1, 1081 BT Amsterdam, The Netherlands

Background. More than half of patients who present with depressive disorders also have elevated comorbid anxiety symptoms. Given the high comorbidity between these disorders, it is important to understand the extent that psychotherapies for depression additionally ameliorate symptoms of anxiety.

Methods. Systematic searches were conducted in PubMed, PSYCinfo, EMBASE, and the Cochrane Registry of Controlled Trials. Included studies were randomized controlled trials that compared psychotherapy compared with a control condition for the treatment of adults with a primary diagnosis or elevated symptoms of depression and that examined the effects of treatment on anxiety outcomes. Acute phase depression and anxiety (continuous measure) outcomes were extracted. Effect sizes were calculated by subtracting the average post-treatment scores of the psychotherapy group from the average post-treatment scores of the comparison group divided by the pooled standard deviation.

Results. Fifty-two studies of varying quality met the inclusion criteria. Pooled effect sizes showed that anxiety outcomes were significantly lower in the psychotherapy conditions than in control conditions at post-treatment [$g = 0.52$; 95% confidence interval (CI) 0.44–0.60; NNT (numbers-needed-to-treat) = 3.50]. Moderate heterogeneity was observed ($I^2 = 55\%$, 95% CI 40–66). Bivariate metaregression analysis revealed a significant association between depression and anxiety effect sizes at post-treatment. Longer-term follow-ups of up to 14 months post-baseline showed indications for a small lasting effect of psychotherapy on anxiety outcomes ($g = 0.27$).

Conclusions. This meta-analysis provides evidence that psychotherapy aimed at depression can also reduce anxiety symptoms in relation to control conditions.

Introduction

It is well established that depressive and anxiety symptoms often co-occur in patients with a primary diagnosis of either a depressive or anxiety disorder (Brown *et al.* 2001; Wiethoff *et al.* 2010). This particular co-morbid symptomatology can complicate treatment and lead to worsening outcomes (Joffe *et al.* 1993; Zajecka & Ross 1995; Fava *et al.* 2004; Wiethoff *et al.* 2010). Given the high overlap of depressive and anxiety symptoms, it is important to understand the extent to which psychotherapies designed for the treatment of depression can additionally ameliorate symptoms of anxiety.

Although depression and anxiety disorders are classified separately in the DSM, high rates of co-occurring symptoms of anxiety and depression, genetic links, and similarities in the etiology and course of the disorders have led some to consider anxiety and depression as part of a coherent spectrum (Goldberg *et al.* 2009). Despite the evidence of co-occurrence, few studies have examined the efficacy of psychological treatments designed for targeting comorbid anxiety and depressive disorders simultaneously (Smits *et al.* 2009). This may be due to treatment efficacy literature being mainly structured by a psychiatric disorder. Thus, effective psychological and pharmacologic therapies have been developed for the treatment of depression or anxiety disorders separately and the effects of these treatments cannot be extended to symptoms of comorbid disorders.

Previous meta-analyses have shown that varied psychotherapies such as cognitive behavioral therapy (CBT), interpersonal psychotherapy (IPT), problem-solving therapy (PST), and possibly psychodynamic psychotherapy, as well as pharmacotherapies such as SSRIs and MAOIs are effective for treating depression (De Maat *et al.* 2006; Malouff *et al.* 2007; Cuijpers *et al.* 2011, 2013b). Meanwhile, among anxiety disorders, CBT administered alone or in combination with pharmacotherapy have been the most widely-studied and proven efficacious treatments for anxiety symptoms (Furukawa *et al.* 2004; Hunot *et al.* 2007; Acarturk *et al.* 2009; Sanchez-Meca *et al.* 2010; Cuijpers *et al.* 2014).

The high prevalence of co-morbid depression and anxiety spurred several meta-analyses examining whether CBT for the treatment of anxiety disorders reduces comorbid depressive symptoms. These meta-analyses demonstrate that CBT for Generalized Anxiety Disorder (GAD), Social Anxiety Disorder, and Panic Disorder significantly reduces depressive

symptoms compared with control conditions, with effect sizes ranging from 0.60 to 1.00 (Mitte, 2005; Hunot *et al.* 2007; Hofmann & Smits, 2008; Acarturk *et al.* 2009; Sanchez-Meca *et al.* 2010; Cuijpers *et al.* 2014). In addition, a recent meta-analysis comparing the effects of treatments designed for anxiety disorders with depression treatments directly, found no differences between the two treatments on depression outcomes (Cuijpers *et al.* 2016).

Although this provides robust meta-analytic evidence that treatments designed for and studied in populations with anxiety disorders also significantly reduce depressive symptoms, data on whether typical psychotherapies for depression also reduce symptoms of anxiety have not yet been aggregated. Individual randomized controlled trials (RCTs) have shown mixed results with equal numbers of studies indicating a significant reduction in anxiety symptoms between depression psychotherapy and control conditions (e.g., Serfaty *et al.* 2009; Bohlmeijer *et al.* 2011; Buntrock *et al.* 2015; Milgrom *et al.* 2015) as demonstrating no significant difference between depression psychotherapies and control conditions (e.g. Chesney *et al.* 2003; Lynch *et al.* 2004; Chiesa *et al.* 2012; Lemma & Fonagy, 2013; Buhrman, *et al.* 2015), and a few studies even finding negative effects (see Fig. 2). Since individual RCTs have limited statistical power, aggregating these data would provide more robust evidence for whether depression psychotherapies ameliorate anxiety symptoms. This information could be valuable to clinicians to better support their treatment selections. Thus, this meta-analysis aims to examine whether the effects of psychotherapies for depression compared with control conditions reduces post-treatment anxiety symptoms, in addition to post-treatment depressive symptoms. It also examines the relationship between depression and anxiety symptoms after psychotherapeutic treatment.

Methods

Identification and study selection

A database of RCTs on psychological treatment of adult depression was utilized for study identification and selection. This database has been described elsewhere (Cuijpers *et al.* 2008) and has been used in a series of previously published meta-analyses (<http://www.evidencebasedpsychotherapies.org>). It was developed through comprehensive literature searches (from 1966 to January, 1 2016). During these searches, 16 365 abstracts were examined for inclusion from PubMed, PSYCinfo, EMBASE, and the Cochrane Registry of Controlled Trials, and previously conducted meta-analyses. Studies on the treatment of adult depression were examined for inclusion. From the 16 407 abstracts identified and examined against inclusion and exclusion criteria (13 384 after duplicate removal), 1885 full-text articles were retrieved for potential inclusion in the database. A total of 645 studies met criteria, were incorporated in the database, and were then checked for inclusion in this meta-analysis.

This study includes RCTs comparing psychotherapeutic treatment aimed at treating depression compared with control conditions (wait-list, pill placebo, care-as-usual, other) for adults with a primary diagnosis or elevated symptoms of depression established by a standardized diagnostic interview or a standardized clinician or self-report measure of depressive symptoms, which utilize a continuous measure of anxiety (general symptoms). No language restrictions were applied. Psychotherapeutic treatment was defined according to previously delineated criteria (Cuijpers

et al. 2008) and included interventions in which either verbal communication between a therapist and client was central to the psychotherapy, or psychological treatment in book or internet format which clients work through individually supported by a therapist (by telephone or e-mail). Usual care was defined as patients receiving the same care they would have received if they never entered the trial.

Studies of patients under 18 years, those with an intellectual disability, or that included less than three treatment sessions were excluded. RCTs not reporting outcome scores on a continuous, validated measure of anxiety symptoms were excluded as effect sizes that pertain to this studies' main question could not be calculated. Studies of patients with co-morbid medical diagnoses or post-partum depression (special populations) were not excluded.

Quality assessment and data extraction

Quality of the included studies was assessed using four criteria of the Cochrane collaboration's 'risk of bias' tool (eds Higgins & Green, 2011) which assesses study validity by examining possible sources of bias within the RCTs. This included assessing whether (1) randomization was adequately generated, (2) and properly concealed, (3) if appropriate measures were taken to prevent knowledge of the treatment allocation to parties, and (4) if appropriate methods for handling missing data were utilized. A negative score was given to a study when quality criteria were not handled adequately or there was not enough information to rate the item.

Data extracted from the published papers included continuous outcomes on depression or anxiety symptom scales and reported characteristics of the studies such as recruitment method (community or clinical populations) target group of the study (adults or other populations like comorbid medical diagnosis or older adults), and depression inclusion criteria. We also reported several facets of the psychological treatment: treatment delivery (group, individual, guided self-help), treatment length and type of psychotherapy (CBT or other). 'Other' psychotherapy includes IPT, PST, psychodynamic psychotherapy, and others. These were combined since too few studies are available within each treatment to analyze them separately. Two independent raters performed the data extraction (EW, AK).

Meta-analysis

Effect sizes indicating the difference between psychotherapy and controls on post-treatment anxiety or depression outcome measure scores were calculated (Hedges' *g*) by subtracting the average post-treatment anxiety score of the psychotherapy group from the average post-treatment anxiety score of the comparison group and dividing by the pooled standard deviation. The same calculation was conducted for post-treatment depression scores. If post-treatment effect sizes were not reported, mean depression or anxiety symptom change from pre-treatment to post-treatment was utilized. Effect sizes of 0.8 are considered large, 0.5 moderate, and 0.2 and below are considered to be small (Cohen, 1988).

Whenever possible, effect sizes associated with the intention-to-treat samples were utilized, however when not available, completer samples effect sizes were used. If two depression or anxiety measures were utilized, an average effect size was computed. In the event that two control groups were utilized in a single RCT, care-as-usual was chosen as the control as it mimics real-world practice. If CAU was not available, then another

control condition was chosen as the comparison. In order to minimize overestimation of the effect sizes, wait-list controls were not chosen as the comparison condition when another control was available because they produce significantly larger effect sizes (Furukawa *et al.* 2014).

The Comprehensive Meta-analysis program (CMA version 3) was utilized to calculate pooled mean effect sizes. Random effects models were chosen for all analyses as we expected some heterogeneity between studies. In addition to standardized effect sizes, we also calculated the numbers-needed-to-treat (NNT) using the Kraemer & Kupfer (2006) formula. The NNT designates the number of patients that would need to be treated with psychotherapy to have one additional positive outcome compared to a control condition.

Heterogeneity between included studies was examined by calculating I^2 , which quantifies heterogeneity uncovered by the Q-statistic and reports (in percentages) how much overall variance is attributed to between-study variance. An I^2 of 25% indicates low heterogeneity, 50% moderate heterogeneity, and 75% indicated high heterogeneity. The 95% confidence interval (CI) around I^2 is also calculated using the non-central chi-square approach in the heterogi module of STATA (Ioannidis *et al.* 2007; von Hippel, 2015).

Subgroup analyses were conducted to determine how varying sample and study characteristics influence the difference between psychotherapy and control conditions in treating anxiety symptoms. Subgroup analysis was conducted using the mixed effect model, which pools subgroups of studies using the random effects model, and tests for subgroup differences by using the fixed effects model. Subgroups were pooled according to the type of psychotherapy utilized, method of recruitment, format of the psychotherapy, study eligibility criteria (including depressive symptoms *v.* diagnosis of depression), type of control condition, study quality, and inclusion of special populations.

In order to understand the relationship between depression and anxiety outcomes, additional metaregression analyses were conducted in CMA. Bivariate analysis examined the relationship between depression effect sizes and anxiety effect sizes (with anxiety ES as the dependent variable). Multivariate metaregression analysis examined this relationship while controlling for relevant study and clinical characteristics. Because studies included varying lengths of treatments, metaregression analysis examined number of treatment sessions as a moderator of treatment outcomes.

Publication bias was assessed in several ways. First, a funnel plot of effect sizes was visually inspected to see whether a larger number of trials clustered in the bottom right, which would indicate publication bias. Duval and Tweedie's trim-and-fill procedure was utilized in order to calculate the approximate number of studies missing from the funnel plot and transform this into an effect size corrected for publication bias. Egger's test of the intercept, which quantifies bias detected in the funnel plot, was performed using procedures outlined by Hedges & Olkin (1985).

Results

From the 645 studies included in the database, a total of 52 studies met the inclusion criteria for this meta-analysis. Three hundred eighty-three studies were excluded for not having control conditions, 186 studies did not have anxiety measurements, and 24 were excluded for other reasons (e.g. could not extract data).

The PRISMA flowchart outlining the inclusion process is presented in Fig. 1. Specific design and clinical characteristics of the studies are presented in Table 1.

Characteristics of included studies

A total of 52 RCT trials with 62 comparisons between psychotherapy (3072 patients) *v.* control conditions (2665 patients) and reporting post-treatment anxiety symptom inventories were included in this meta-analysis. Out of the 52 included trials, 22 were aimed at adults in general, nine trials focused on older adults, 13 trials included adults with medical diseases, and eight trials were aimed at another specific group such as post-partum depression. A total of 30 studies recruited participants from community sources, 12 trials recruited strictly from clinical populations, and 10 trials recruited in another manner or used a combination of methods.

Various psychotherapies for depression were examined in the included trials. Of the 62 comparisons about half ($n = 33$) utilized CBT as the psychotherapy and 29 examined other types of psychotherapy (Table 1). The control condition comparison groups also varied: care-as-usual was examined in 29 comparisons, wait-list in 23 comparisons, and another type of control such as attention-controls or psychoeducation in 10 comparisons. In terms of psychotherapy format, 26 comparisons delivered treatment via individual psychotherapy, 17 comparisons used group therapy (one additional trial used both individual and group treatment), and 17 comparisons utilized guided self-help. The number of treatment sessions ranged from three to 20 in the active interventions (mode: 8). Most studies were conducted in Europe or North America, and seven were conducted in other countries.

Measurements included in the trials included commonly used self-report measures of anxiety symptoms. Fourteen studies (15 comparisons) administered the anxiety subscale of the Hospital Anxiety and Depression Scale (HADS), 16 studies (19 comparisons) administered the Beck Anxiety Inventory (BAI) alone or in combination with another anxiety scale, 20 studies used other valid anxiety scales. Most of the included measures target general symptoms of anxiety and do not pertain to specific anxiety disorders, allowing for thorough comparison. Two studies used the Penn State Worry Scale (PSWS), which is often used in studies of GAD, but assesses more general symptoms of anxiety. Two other studies utilized scales specifically for the measurement of PTSD symptoms (Meyer *et al.* 1990). For this reason, a sensitivity analysis was conducted excluding these studies.

Measurements of depression included common self-report and clinician-rated measurements with a majority of trials utilizing the CES-D, HAM-D, or BDI alone or in combination with another depression measure. Ten studies used other validated depression measures.

Quality assessment

Study quality of the included trials varied: 34 out of 52 studies used an adequate generation of the randomization sequence, 33 studies reported allocating participants to conditions by an independent party, 50 studies reported using self-report outcomes and two blind assessors, and 37 studies utilized intention-to-treat samples indicating that missing data were handled appropriately. Thirty-five studies scored positive on three or four items of the risk of bias, and the remaining 17 studies scored positive on 0–2 items.




Fig. 1. Flowchart of study inclusion.

Anxiety outcomes

Differences between psychotherapy and control conditions on post-treatment anxiety symptoms were examined in 62 comparisons and the pooled effect size showed significantly lower anxiety symptoms in psychotherapy conditions than in control conditions at post-treatment ($g = 0.52$; 95% CI 0.44–0.60; NNT = 3.50) (Table 2, Fig. 2). Moderate heterogeneity was observed ($I^2 = 55\%$, 95% CI 40–66). Outliers (nine comparisons), with CIs positioned outside of the pooled effect CIs were omitted from the analysis and an effect size of $g = 0.49$ (95% CI 0.42–0.55; $I^2 = 14$, 95% CI 0–40) was observed (Table 2). Multiple comparisons from a single study were included in 10 trials, thereby violating the independence assumption. Because this may artificially reduce heterogeneity, two additional sensitivity analyses including only one effect per study (the highest, then the lowest) were conducted. In both sensitivity analyses, effect sizes and heterogeneity differed little from the original analysis (Table 2). A further sensitivity

analysis excluding studies utilizing PTSD measures found similar effect sizes ($g = 0.52$; 95% CI 0.44–0.60).

Additional analyses were conducted to examine effect sizes by specific anxiety measures. Examining only studies using the BAI revealed an effect size of $g = 0.40$ (95% CI 0.27–0.53). Studies that utilized the HADS-A produced an effect size of $g = 0.58$ (95% CI 0.43–0.73).

The funnel plot and Duval and Tweedie's trim and fill procedure showed evidence for little publication bias. In addition, Egger's test of the intercept was not significant ($p = 0.32$). When adjusting for missing studies/small sample bias the effect size estimate was similar to the observed effect size ($g = 0.50$; 95% CI 0.42–0.59).

Depression outcomes

Secondary analysis examined the overall effect size of psychotherapy compared with control conditions on outcome depression

Table 1. Selected characteristics of studies examining the effects of psychotherapy for depression on anxiety

Study	Target population	Depression	Psycho-therapy	Control	Format	N _{ses}	N post PST	N post CTRL	Anxiety outcome	Qual ^a	Coun-try
1. Ammerman <i>et al.</i> (2013)	Mothers	Dep symptoms	CBT	CAU	Indiv	15	47	46	BSL-A	— + sr +	US
2. Andersson <i>et al.</i> (2002)	Adults	MDD	CBT	Other	Gsh	5	36	49	BAI	+ + sr —	SE
3. Batink <i>et al.</i> (2013)	Adults	Dep symptoms	MBCT	TAU	Grp	8	64	66	PSWQ	+ + sr +	NL
4. Bohlmeijer <i>et al.</i> (2011)	Adults	Dep symptoms	ACT	WL	Grp	8	49	44	HADS-A	+ + sr +	NL
5. Buhrman <i>et al.</i> (2015)	Adults w/chronic pain	Dep symptoms	CBT	WL	Gsh	8	28	24	BAI	+ + sr +	SE
6. Buntrock <i>et al.</i> (2015)	Adults	Dep symptoms	CBT	Other	Gsh	6	202	204	HADS-A	+ + sr +	DE
7. Carlbring <i>et al.</i> (2013)	Adults	Dep symptoms	BA + ACT	WL	Gsh	8	40	40	BAI	+ + sr +	SE
8. Chesney <i>et al.</i> (2003)	Adults with HIV	Dep symptoms	Other	Other	Grp	10	46	44	STAI	— — sr —	US
9. Chiesa <i>et al.</i> (2012)	Adults	MDD	MBCT	Other	Indiv	8	23	20	BAI	+ + sr —	IT
10. Cramer <i>et al.</i> (2011)	Adults	Dep symptoms	CBT	CAU	Grp	12	46	19	BAI	+ + sr —	US
11. Dobkin <i>et al.</i> (2011)	Adults w/parkinsons	Mood disorder	CBT	Other	Indiv	10	41	39	HADS-A	+ + sr +	US
12. Evans & Connis (1995)	Adults w/Cancer	Dep symptoms	CBT	CAU	Grp	8	27	24	SCL-A	— — sr —	US
			SUP*				21				
13. Faramarzi <i>et al.</i> (2008)	Women w/infertility	MDD	CBT	CAU	Grp	10	29	30	ASQ	— — sr —	IR
14. Fledderus <i>et al.</i> (2012)	Adults	Dep symptoms	ACT-E	WL	Gsh	9	125	126	HADS-A	+ — sr +	NL
			ACT-M				125				
15. Freedland <i>et al.</i> (2010)	Adults w/coronary bypass	Mood disorder	CBT	CAU	Indiv	9	41	40	BAI	+ + sr +	US
			SUP				42				
16. Freedland <i>et al.</i> (2015)	Adults w/heart failure	MDD	CBT	e-CAU	Indiv	24	79	79	BAI	+ + sr +	US
17. Gitlin <i>et al.</i> (2014)	Older African American adults	Dep symptoms	Other	WL	Indiv	10	106	102	STAI (state)	+ + sr +	US
18. Grote <i>et al.</i> (2009)	Pre/Post-partum	Dep symptoms	IPT	e-CAU	Indiv	8	25	28	BAI	— — sr +	US
19. Haringsma <i>et al.</i> (2006)	Older adults	Dep symptoms	CBT	WL	Grp	10	52	58	HADS-A	— — sr +	NL
20. Hautzinger & Welz (2004)	Older adults	Dep symptoms	CBT	WL	Grp	12	55	30	SCL-A	+ + sr +	DE
21. Hsiao <i>et al.</i> (2014)	Adjustment disorder w/dep mood	Dep symptoms	Other (BMS)	CAU	Grp	8	33	37	STAI	+ + sr +	TW
22. Johansson <i>et al.</i> (2012a)	Adults	MDD	DYN	Other	Gsh	9	46	46	BAI/GAD-7	+ + sr +	SE
23. Johansson <i>et al.</i> (2012b)	Adults	MDD	CBT std	Other	Gsh	8	34	42	BAI	+ + sr +	SE
			CBT tail				36				
24. Jonkers <i>et al.</i> (2012)	Older adults w/chronic illness	Dep symptoms	Other (Life Review)	CAU	Gsh	10	183	178	SCL-A	+ + sr +	NL
25. Kelly <i>et al.</i> (1993)	Older adults	Dep symptoms	CBT	CAU	Grp	8	27	27	SCL-A	— — sr —	US
			SUP				14				
26. Kivi <i>et al.</i> (2014)	Adults	Dep symptoms	CBT	CAU	Indiv	7	45	47	BAI	— — sr —	SE
27. Korte <i>et al.</i> (2012)	Older adults	Dep symptoms	Other (Life Review)	CAU	Grp	8	100	102	HADS-A	+ + sr +	NL
28. Laidlaw <i>et al.</i> (2008)	Older adults	MDD	CBT	CAU	Indiv	8	20	20	PSWI	+ + sr —	UK
29. Lamers <i>et al.</i> (2015)	Older adults	Dep symptoms	Other (Life Review)	WL	Gsh	7	58	58	HADS-A	+ + sr +	NL

30. Lemma & Fonagy (2013)	Adults	Dep symptoms	DYN	Other	Grp	8	8	8	GAD-7	-- sr +	UK
31. Lynch <i>et al.</i> (2004)	Adults	Dep symptoms	PST	CAU	Gsh	6	9	13	DHP-A	-- sr -	US
32. Maina <i>et al.</i> (2005)	Adults	Mood disorder	PST	WL	Indiv	20	10	10	HAM-A	-- + +	IT
			SUP				10				
33. Martin <i>et al.</i> (2015)	Adults w/migraine	MDD	CBT	CAU	Indiv	12	18	26	BAI	-- sr -	AU
34. Milgrom <i>et al.</i> (2015)	Women w/antenatal depression	MDD	CBT	CAU	Indiv	8	27	27	BAI	+ + sr +	AU
35. Naeem <i>et al.</i> (2014)	Adults	MDD	CBT	CAU	Gsh	7	94	89	HADS-A	-- sr -	PK
36. Naeem <i>et al.</i> (2015)	Adults	MDD	CBT	CAU	Indiv	7	69	68	HADS-A	+ + sr +	PK
37. Pot <i>et al.</i> (2010)	Older adults	Dep symptoms	Other (Life Review)	Other	Grp	12	83	88	HADS-A	+ + sr +	NL
38. Pots <i>et al.</i> (2014)	Adults	Dep symptoms	MBCT	WL	Grp	12	76	75	HADS-A	+ + sr +	NL
39. Qiu <i>et al.</i> (2013)	Adults w/breast cancer	MDD	CBT	WL	Grp	10	31	31	SAS	+ + sr +	CN
40. Richards <i>et al.</i> (2015)	Adults	Dep symptoms	CBT	WL	Gsh	7	96	92	GAD-7	+ + sr +	UK
41. Savard <i>et al.</i> (2006)	Adults w/breast cancer	Dep symptoms	CBT	WL	Indiv	8	21	16	HADS-A	+ + sr -	CA
42. Serfaty <i>et al.</i> (2009)	Older adults	Mood disorder	CBT	CAU	Indiv	12	60	53	BAI	+ + sr -	UK
43. Simson <i>et al.</i> (2008)	Adults w/diabetes	Dep symptoms	SUP	CAU	Indiv	5	15	15	HADS-A	-- sr +	DE
44. Strong <i>et al.</i> (2008)	Adults w/cancer	MDD	PST	CAU	Indiv	10	97	99	SCL-A	+ + sr +	UK
45. Swartz <i>et al.</i> (2008)	Mothers w/depression	MDD	IPT	CAU	Indiv	8	23	17	BAI	-- sr +	US
46. Talbot <i>et al.</i> (2011)	Women w/ h_x sexual abuse	MDD	IPT	CAU	Indiv	16	31	22	PTSD-SSS	-- sr +	US
47. Tovote <i>et al.</i> (2014)	Adults w/diabetes	Dep symptoms	MBCT	WL	Indiv	8	31	31	GAD-7	+ - sr +	NL
			CBT								
48. Vernmark <i>et al.</i> (2010)	Adults	MDD	CBT std	WL	Gsh	7	29	29	BAI	+ + sr +	SE
			CBT tail				27				
49. Vitriol <i>et al.</i> (2009)	Women w/severe dep & h_x child abuse	Mood disorder	DYN	CAU	Indiv	12	44	43	PTO	-- + +	CL
50. Warmerdam <i>et al.</i> (2008)	Adults	Dep symptoms	CBT	WL	Gsh	8	88	87	HADS-A	+ + sr +	NL
			PST				88				
51. Watkins <i>et al.</i> (2012)	Adults	Dep symptoms	Other (CNT)	CAU	Gsh	3	33	37	GAD-7	+ + sr +	UK
52. Wierzbicki & Bartlett (1987)	Adults	Mood disorder	CBT (ind)	WL	Indiv & Grp	6	9	20	A- state/ A-trait	-- sr -	US
			CBT (grp)								
			CBT (grp)								

^aIn this column a positive or negative sign is given for four quality criteria, respectively: allocation sequence; concealment of allocation to conditions; blinding of assessors for anxiety outcomes; and intention-to-treat analyses.

N at post-treat, N for anxiety analysis; Guided self-help, online or book intervention with contact from therapist.

Clin, Clinical Sample; Comm, Community Sample; CBT, Cognitive Behavior Therapy; ACT, Acceptance and Commitment Therapy; MBCT, Mindfulness Based Cognitive Therapy; SUP, Supportive psychotherapy; IPT, Interpersonal Psychotherapy; BMS, Body, Mind, Spirit Psychotherapy; DYN, Psychodynamic Psychotherapy; Indiv, Individual; Grp, Group; Gsh, guided self-help; CAU, Care-as-Usual; WL, Wait-list; BSL, Brief Symptom Inventory; BDI, Beck Depression Inventory; MADRS, Montgomery Åsberg Depression Scale; CES-D, Center for Epidemiologic Studies Depression Scale; HAM-D, Hamilton Rating Scale for Depression; PHQ-9, Patient Health Questionnaire; EPDS, Edinburgh Post-natal Depression Scale; HADS, Hospital Anxiety Depression Scale; GDS, Geriatric Depression Scale; IDS, Inventory of Depressive Symptomatology; SCL, Symptom Checklist; DHP, Duke Health Profile (depression); D30, Depression scale from the Minnesota Multiphasic Personality Inventory. BAI, Beck Anxiety Inventory; STAI, State-trait Anxiety Inventory; ASQ, Anxiety Screening Questionnaire; GAD-7, Generalized Anxiety Disorder 7; PSWI, Penn State Worry Inventory; HAM-A, Hamilton Rating Scale for Anxiety; PTSD-SSS, Modified PTSD Symptom Scale Self-Report; PTO, Post-traumatic Stress Disorder Assessment.

Table 2. Results of meta-analysis of psychotherapy v. control conditions

	<i>N</i>	<i>g</i>	95% CI	<i>Z</i>	<i>I</i> ²	95% CI	<i>p</i>
Anxiety acute outcomes							
All comparisons	62	0.52	0.44–0.60	12.58	55	40–66	
One ES per study (only highest)	52	0.52	0.43–0.61	11.39	58	43–69	
One ES per study (only lowest)	52	0.49	0.40–0.58	11.11	55	38–67	
ES w/o outliers (9 comparisons)	53	0.49	0.42–0.55	15.09	14	0–40	
BAI only	19	0.40	0.27–0.53	5.86	33	0–62	
HADS-A only	15	0.58	0.43–0.73	7.50	69	48–82	
Subgroup analyses							
Type of psychotherapy							
CBT	33	0.54	0.44–0.65	10.21	50	25–66	0.57
Other	29	0.49	0.36–0.62	7.53	61	41–74	
Recruitment							
Community	36	0.49	0.41–0.57	11.70	31	0–54	0.55
Clinical	14	0.62	0.36–0.87	4.73	73	54–84	
Other	12	0.56	0.36–0.76	5.47	64	33–81	
Format ^a							
Individual	26	0.50	0.34–0.65	6.31	66	48–77	0.24
Group	17	0.46	0.33–0.59	7.04	30	0–61	
Guided self-help	17	0.60	0.49–0.71	10.68	39	0–66	
Depression diagnosis							
MDD	19	0.60	0.45–0.76	7.71	56	26–74	0.40
Mood disorder	10	0.55	0.28–0.83	4.01	68	38–83	
Depressive symptoms	33	0.48	0.39–0.58	9.89	44	16–63	
Type of control group							
WL	23	0.59	0.48–0.70	10.48	37	0–62	0.08
CAU	29	0.51	0.38–0.65	7.32	65	48–76	
Other	10	0.38	0.23–0.53	5.05	26	0–64	
Study quality							
High	41	0.49	0.41–0.57	11.50	50	28–65	0.23
Low	21	0.62	0.42–0.83	5.97	62	40–76	
Special population							
Yes	33	0.42	0.33–0.51	8.94	32	0–55	0.01
No	29	0.61	0.49–0.73	10.08	57	35–72	
Long-term effects post-baseline							
Up to 7 month FU	11	0.25	0.13–0.37	4.15	0	0–60	
Up to 14 month FU	10	0.27	0.17–0.38	5.05	0	0–62	
Depression acute outcomes							
All comparisons	62	0.63	0.54–0.71	14.45	59	46–69	
One ES per study (only highest)	52	0.64	0.54–0.74	12.69	65	53–74	
One ES per study (only lowest)	52	0.59	0.50–0.69	12.33	62	48–72	
ES without outliers (eight comparisons)	54	0.61	0.55–0.67	19.44	13	0–38	

^aThe Wierzbicki & Bartlett (1987) study was removed from the analysis because both group and individual psychotherapy formats were utilized.




Fig. 2. Forest plot of included studies.

scores in 62 comparisons (Table 2). The pooled effect size was $g = 0.63$ (95% CI 0.54–0.71; NNT = 2.82) and heterogeneity between studies was moderate ($I^2 = 59\%$; CI 46–69). Eight outliers were omitted from the meta-analysis and an effect size of $g = 0.61$ (95% CI 0.55–0.67; $I^2 = 13\%$, 95% CI 0–38) was observed. Inspection of the funnel plot and Duval and Tweedie's trim and fill procedure indicated minor publication bias. Egger's test of the intercept was not significant ($p = 0.07$). After adjustment for missing studies the effect size was similar to the observed effect size ($g = 0.57$; 95% CI 0.47–0.66).

In addition, eight studies examined multiple psychotherapy interventions in one trial. Analyses utilizing only the highest and then the lowest effect sizes per study were conducted (Table 2).

Subgroup and meta-regression analysis

Subgroup analyses (see Table 2) showed no significant differences in effect sizes when examining differences in recruitment

method, depression inclusion criteria, psychotherapy format, or study quality. The analysis did indicate a trend toward studies with wait-list control groups displaying higher effect sizes on anxiety than studies utilizing CAU or other control conditions ($p = 0.08$). In addition, studies on the general population had a significantly higher effect size than those studies with special populations, such as comorbid medical disorders ($p = 0.01$).

Bivariate meta-regression analyses examining associations between anxiety effect sizes and continuous variables revealed a significant association between depression and anxiety effect sizes at post-treatment (Table 3, Fig. 3). Additional multivariate meta-regression analysis showed depression effect sizes remained significantly associated with anxiety effect sizes at post-treatment after controlling for study and clinical characteristics (Table 3). Additionally, a number of treatment sessions did not predict anxiety outcomes between psychotherapies and control conditions ($b = 0.00$; $p = 0.78$).

Table 3. Results of metaregression analysis

		<i>b</i>	95% CI	<i>p</i> value
Bivariate				
Intercept		0.18	0.05–0.30	0.01
Depression ES		0.55	0.37–0.72	0.00
Multivariate				
Intercept		0.36	0.04–0.67	0.03
Depression ES		0.52	0.33–0.72	0.00
Psychotherapy (CBT = 0)		0.02	–0.11–0.14	0.82
Recruitment	– Clinical	Ref		
	– Community	–0.05	–0.26–0.15	0.61
	– Other	–0.07	–0.28–0.15	0.55
Format	– Individual	Ref		
	– Group	–0.06	–0.25–0.13	0.55
	– Gsh	–0.01	–0.23–0.20	0.91
Dep type	– MDD	Ref		
	– Mood Dis	0.01	–0.21–0.23	0.92
	– Symptoms	0.02	–0.15–0.20	0.78
Control	– WL	Ref		
	– CAU	–0.01	–0.21–0.19	0.92
Other	– Other	–0.18	–0.36–0.01	0.04
Study quality (high = ref)		0.08	–0.09–0.25	0.33
Special pop (no = ref)		–0.21	–0.38–0.04	0.02

Long-term follow-up

Studies without wait-list control conditions, indicating participants never received the intervention, and that reported long-term outcomes on anxiety measures were included in follow-up analysis. The pooled effect size signifying the difference between psychotherapy and control conditions in anxiety symptoms up to 7 months post-baseline was $g = 0.25$ (95% CI 0.13–0.37), and up to 14 months post-baseline was $g = 0.27$ (95% CI 0.17–0.38) (Table 2).

Discussion

Our results indicate that, when comparing effect sizes, depression treatment (psychotherapy) is almost as effective at reducing comorbid anxiety symptoms as it is at reducing depressive symptoms. Moderate to large effect sizes were observed for anxiety symptoms ($g = 0.52$; NNT = 3.50) and for depression symptoms ($g = 0.63$; NNT = 2.82) with moderate heterogeneity and little evidence that publication bias affected the results. Long-term follow-up assessments revealed a small lasting effect up to 1 year after baseline assessment ($g = 0.27$), although this should be interpreted with some caution since studies used naturalistic follow-up. For patients with commonly comorbid depression and anxiety symptoms, choosing which treatment to utilize may be challenging. However, these results suggest that common evidence-based psychotherapies for depression can ameliorate anxiety symptoms, and may be sufficient for reducing anxiety symptoms without adjunctive treatments.

However, this meta-analysis was not able to discern if a specific psychotherapy for depression is more effective at treating anxiety symptoms than others or if these psychotherapies are equivalent to utilizing psychotherapies designed to treat anxiety symptoms.

In comparison with previous meta-analyses, anxiety and depression effect sizes found in this sample are comparable to those reported in meta-analyses specifically examining the effects of psychotherapy for anxiety or psychotherapy for depression. Anxiety symptom effect sizes reported here are similar, although slightly lower, than those previously reported in meta-analyses of psychotherapies specifically targeting anxiety which have shown large effect sizes (between 0.71 and 0.84) (Hunot *et al.* 2007; Hofmann & Smits, 2008; Acarturk *et al.* 2009; Sanchez-Meca *et al.* 2010; Cuijpers *et al.* 2014). This study found moderate effect sizes of $g = 0.52$; however, the prior meta-analyses focused on patients diagnosed with anxiety disorders and may include patients with more severe anxiety symptoms. This indicates the validity of the finding that psychotherapy targeting depression also ameliorates anxiety symptoms.

Similarly, depression effect sizes reported here were equivalent to previous meta-analyses which reported effect sizes of psychotherapy for depression *v.* control conditions between 0.56 and 0.82 (Gloaguen *et al.* 1998; Ekers *et al.* 2008; Sanchez-Meca *et al.* 2010; Cuijpers *et al.* 2011, 2012, 2013a). While not all depression trials comparing psychotherapies and controls provided anxiety measures and could be included in this meta-analysis, the sample of trials included here are representative in terms of effect sizes.




Fig. 3. Metaregression of anxiety effect sizes on depression effect size.

Subgroup analysis revealed that studies without special populations exhibited significantly higher effect sizes than studies that included special populations, such as those with a comorbid medical diagnosis or post-partum depression. This may be due to the varying populations included or high statistical heterogeneity between the studies. Consistent with previous research, studies with a wait-list control group displayed a trend towards higher effect sizes than studies utilizing other control conditions (Furukawa *et al.* 2014). No other significant subgroup analyses were identified (including no difference in effect sizes based on the type of psychotherapy, depression inclusion criteria, and control group utilized). It was not possible to examine the effectiveness of the included psychotherapies separately due to the low number of studies meeting criteria per psychotherapy included, therefore providing insufficient statistical power to examine this adequately. Examining whether specific types of psychotherapy were equally effective for the treatment of anxiety symptoms was not the main aim of the study. However, subgroup analysis depicted similar outcomes between CBT and the other psychotherapies included. Future research should examine which specific psychotherapies are optimal for reducing anxiety symptoms.

Separate analysis on anxiety measures found that the BAI produced lower effect sizes than the HADS-A. Although these were both moderate effect sizes, the differences may be related to study characteristics or to differences in the specific measures; however, studies including both measures are needed to examine this further. In addition, there is some heterogeneity among all anxiety measurements included. Although only validated measures of general anxiety symptoms were utilized (with the exception of the two studies utilizing PTSD measures), the potential

effect of including specific measures could not be assessed in this meta-analysis due to insufficient numbers of studies per anxiety measurement.

In addition multivariate metaregression, controlling for clinical and study characteristics, demonstrated a significant association between outcome depression and anxiety scores. This provides support that anxiety and depression are highly connected, however the mechanisms of how depression psychotherapies treat anxiety symptoms, whether certain components of the psychotherapies are more effective than others, and whether depression and anxiety symptoms remit sequentially or simultaneously, is still unclear and should be explored further using RCTs and individual patient data meta-analyses.

Although this meta-analysis provides robust evidence that psychotherapies for depression significantly reduce anxiety symptoms compared with controls, there are several additional limitations to consider when interpreting these results. Only half of the studies met all quality criteria, thus, a majority of the trials included had a considerable risk of bias. The studies scoring as lower quality had a higher effect size (although not significantly) than those scoring as higher quality studies. Thus, study results should be interpreted with some caution. Second, although the meta-analysis includes considerable comparisons, several of the subgroup analyses were conducted with few studies. Thus, finding few significant differences in subgroup analyses might be due to low statistical power and may not indicate that effect sizes for these groups are equivalent. Further analysis of differences in subgroups should be conducted as the evidence accumulates.

Furthermore, because this meta-analysis was conducted on secondary outcome measures, no anxiety symptom inclusion

criteria were required. This implies that baseline anxiety scores for some of the sample may not be in the clinical range. However, because of the high levels of comorbidity between anxiety and depressive symptoms, it is expected that a majority of patients included in these studies had elevated symptoms of anxiety. Nonetheless, these results may not pertain to patients with diagnosed comorbid mood and anxiety disorders and further analysis including patients with elevated symptoms of both should be conducted when enough evidence warrants analysis.

Despite these limitations, this meta-analysis provides evidence for utilizing psychotherapies for depression to treat patients with comorbid anxiety symptoms. When patients present with a main complaint of depression, but exhibit general comorbid anxiety symptoms, psychotherapies for depression can reduce symptoms of anxiety without supplementary treatments. Although this establishes that psychotherapies for depression reduce anxiety symptoms on average (possibly through common factors), not all participants with comorbid anxiety symptoms or disorders will benefit from psychotherapy for depression and certain treatments may be more efficacious for treating anxiety symptoms than others (specific factors). Thus, future research should continue to examine which treatments, study, and individual characteristics may affect treatment response for participants exhibiting comorbid depression and anxiety.

Acknowledgements. The authors would like to thank Eirini Karyotaki for her helpful comments and input.

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of Interest. None.

References

- Acarturk C, Cuijpers P, van Straten A and de Graaf R (2009) Psychological treatment of social anxiety disorder: a meta-analysis. *Psychological Medicine* **39**, 241–254.
- Ammerman RT, Putnam FW, Altaye M, Teeters AR, Stevens J and Van Ginkel JB (2013) Treatment of depressed mothers in home visiting: impact on psychological distress and social functioning. *Child Abuse & Neglect* **37**, 544–554.
- Andersson G, Begstrom J, Hollandare F, Carlbring P, Kaldo V and Ekselius L (2002) Internet-based self-help for depression: randomised controlled trial. *British Journal of Psychiatry* **187**, 456–461.
- Batink T, Peeters F, Geschwind N, van Os J and Wichers M (2013) How does MBCT for depression work? Studying cognitive and affective mediation pathways. *PLoS ONE* **8**, 1–13.
- Bohlmeijer ET, Fledderus M, Roxk TAJJ and Pieterse ME (2011) Efficacy of an early intervention based on acceptance and commitment therapy for adults with depressive symptomatology: evaluation in a randomized controlled trial. *Behaviour Research and Therapy* **49**, 62–67.
- Brown TA, Campbell LA, Lehman CL, Grisham JR and Mancill RB (2001) Current and lifetime comorbidity of the DSM-IV anxiety and mood disorders in a large clinical sample. *Journal of Abnormal Psychology* **110**, 585–599.
- Buhrman M, Syk M, Burvall O, Hartig T, Gordh T and Andersson G (2015) Individualized guided internet-delivered cognitive behaviour therapy for chronic pain patients with Comorbid depression and anxiety: a randomized controlled trial. *Clinical Journal of Pain* **31**, 504–516.
- Buntrock C, Ebert D, Lehr D, Riper H, Smit F, Cuijpers P et al. (2015) Effectiveness of a web-based cognitive behavioural intervention for sub-threshold depression: pragmatic randomised controlled trial. *Psychotherapy and Psychosomatics* **84**, 348–358.
- Carlbring P, Hagglund M, Luthstrom A, Dahlin M, Kadowaki A, Vernmark K et al. (2013) Internet-based behavioral activation and acceptance-based treatment for depression: a randomized controlled trial. *Journal of Affective Disorders* **148**, 331–337.
- Chesney MA, Chambers DB, Taylor JM, Johnson LM and Folkman S (2003) Coping effectiveness training for men living with HIV: results from a randomized clinical trial testing a group-based intervention. *Psychosomatic Medicine* **65**, 1038–1046.
- Chiesa A, Mandelli L and Serretti A (2012) Mindfulness-based cognitive therapy versus psycho-education for patients with major depression who did not achieve remission following antidepressant treatment: a preliminary analysis. *Journal of Alternative and Complementary Medicine* **18**, 756–760.
- Cohen J (1988) *Statistical Power Analysis for the Behavioral Sciences*. 2nd edn. Hillsdale, NJ: Erlbaum.
- Cramer H, Salisbury C, Conrad J, Eldred J and Araya R (2011) Group cognitive behavioural therapy for women with depression: pilot and feasibility study for a randomised controlled trial using mixed methods. *BMC Psychiatry* **11**, 82.
- Cuijpers P, Berking M, Andersson G, Quigley L, Kleiboer A and Dobson KS (2013a). A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments. *The Canadian Journal of Psychiatry/La Revue Canadienne de Psychiatrie* **58**, 376–385.
- Cuijpers P, Cristea IA, Weitz E, Gentili C and Berking M (2016) The effects of cognitive and behavioral therapies for anxiety disorders on depression: a meta-analysis. *Psychological Medicine* **46**, 3451–3462.
- Cuijpers P, Driessen E, Hollon SD, van Oppen P, Barth J and Andersson G (2012) The efficacy of non-directive supportive therapy for adult depression: a meta-analysis. *Clinical Psychology Review* **32**, 280–291.
- Cuijpers P, Geraedts AS, van Oppen P, Andersson G, Markowitz JC and van Straten A (2011) Interpersonal psychotherapy for depression: a meta-analysis. *The American Journal of Psychiatry* **168**, 581–592.
- Cuijpers P, Sijbrandij M, Koole S, Huibers M, Berking M and Andersson G (2014) Psychological treatment of generalized anxiety disorder: a meta-analysis. *Clinical Psychology Review* **34**, 130–140.
- Cuijpers P, Sijbrandij M, Koole SL, Andersson G, Beekman AT and Reynolds CFI (2013b). The efficacy of psychotherapy and pharmacotherapy in treating depressive and anxiety disorders: a meta-analysis of direct comparisons. *World Psychiatry* **12**, 137–148.
- Cuijpers P, van Straten A, Warmerdam L and Andersson G (2008) Psychological treatment of depression: a meta-analytic database of randomized studies. *BMC Psychiatry* **8**, 36.
- De Maat S, Dekker J, Schoevers R and De Jonghe F (2006) Relative efficacy of psychotherapy and pharmacotherapy in the treatment of depression: a meta-analysis. *Psychotherapy Research* **16**, 566–578.
- Dobkin RD, Menza M, Allen LA, Gara MA, Mark MH, Tiu J et al. (2011) Cognitive-Behavioral therapy for depression in Parkinson's disease: a randomized, controlled trial. *American Journal of Psychiatry* **168**, 1066–1074.
- Ekers D, Richards D and Gilbody S (2008) A meta-analysis of randomized trials of behavioural treatment of depression. *Psychological Medicine* **38**, 611–623.
- Evans RL and Connis RT (1995) Comparison of brief group therapies for depressed cancer patients receiving radiation treatment. *Public Health Reports* **110**, 306–311.
- Faramarzi M, Alipor A, Esmaelzadeh S, Kheirkhah F, Poladi K and Pash H (2008) Treatment of depression and anxiety in infertile women: cognitive behavioral therapy versus fluoxetine. *Journal of Affective Disorders* **108**, 159–164.
- Fava M, Alpert JE, Carmin CN, Wisniewski SR, Trivedi MH, Biggs MM et al. (2004) Clinical correlates and symptom patterns of anxious depression among patients with major depressive disorder in STAR*D. *Psychological Medicine* **34**, 1299–1308.
- Fledderus M, Bohlmeijer ET, Pieterse ME and Schreurs KM (2012) Acceptance and commitment therapy as guided self-help for psychological distress and positive mental health: a randomized controlled trial. *Psychological Medicine* **42**, 485–495.
- Freedland KE, Carney RM, Rich MW, Steinmeyer BC and Rubin EH (2015) Cognitive behavior therapy for depression and self-care in heart failure patients: a randomized clinical trial. *JAMA Internal Medicine* **175**, 1773–1782.

- Freedland KE, Skala JA, Carney RM, Rubin EH, Lustman PJ, Davila-Roman VG *et al.* (2010) Treatment of depression after coronary artery bypass surgery. *Archives of General Psychiatry* **66**, 387–396.
- Furukawa TA, Noma H, Caldwell DM, Honyashiki M, Shinohara K, Imai H *et al.* (2014) Waiting list may be a Nocebo condition in psychotherapy trials: a contribution from network meta-analysis. *Acta Psychiatrica Scandinavica* **130**, 181–192.
- Furukawa TA, Watanabe N and Churchill RA (2004) Psychotherapy plus antidepressant for panic disorder with or without agoraphobia systematic review. *British Journal of Psychiatry* **188**, 305–312.
- Gitlin LN, Harris LF, McCoy MC, Chernett NL, Pizzi LT, Jutkowitz E *et al.* (2014) A home-based intervention to reduce depressive symptoms and improve quality of life in older African Americans. *Annals of Internal Medicine* **159**, 243–252.
- Gloaguen V, Cottraux J, Cucherat M and Blackburn I (1998) A meta-analysis of the effects of cognitive therapy in depressed patients. *Journal of Affective Disorders* **49**, 59–72.
- Goldberg DP, Krueger RF, Andrews G and Hobbs MJ (2009) Emotional disorders: cluster 4 of the proposed meta-structure for DSM-V and ICD-11. *Psychological Medicine* **39**, 2043–2059.
- Grote NK, Swartz HA, Geibel SL, Zuckoff A, Houck PR and Frank E (2009) A randomized controlled trial of culturally relevant, brief interpersonal psychotherapy for perinatal depression. *Psychiatric Services* **60**, 313–321.
- Haringsma R, Engels GI, Cuijpers P and Spinhoven P (2006) Effectiveness of the Coping with Depression (CWD) course for older adults provided by the community-based mental health care system in the Netherlands: a randomized controlled field trial. *International Psychogeriatrics* **18**, 307–325.
- Hautzinger M and Welz S (2004) Kognitive verhaltenstherapie bei depressionen im alter: Ergebnisse einer kontrollierten vergleichsstudie unter ambulanten bedingungen an depressionen mittleren schweregrads. *Zeitschrift fur Gerontologie und Geriatrie* **37**, 427–435.
- Hedges L and Olkin I (1985) *Statistical Methods for Meta-analysis*. Academic Press: Orlando.
- Higgins JP, Green S (eds) (2011) Chapter 8: assessing risk of bias in included studies. In *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5. <http://handbook.cochrane.org/>
- Hofmann SG and Smits JAJ (2008) Cognitive-behavioral therapy for adult anxiety disorders: a meta-analysis of randomized placebo-controlled trials. *The Journal of Clinical Psychiatry* **69**, 621–632.
- Hsiao FH, Lai YM, Chen YT, Yang TT, Liao SC, Ho RTH *et al.* (2014) Efficacy of psychotherapy on diurnal cortisol patterns and suicidal ideation in adjustment disorder with depressed mood. *General Hospital Psychiatry* **36**, 214–219.
- Hunot V, Churchill R, Silva de Lima M and Teixeira V (2007) Psychological therapies for generalised anxiety disorder. *Cochrane Database of Systematic Reviews*, CD001848, 1–93.
- Ioannidis JPA, Patsopoulos NA and Evangelou E (2007) Uncertainty in heterogeneity estimates in meta-analyses. *British Medical Journal* **335**, 914–916.
- Joffe R, Bagby R and Levitt A (1993) Anxious and nonanxious depression. *American Journal of Psychiatry* **150**, 1257–1258.
- Johansson R, Ekbladh S, Hebert A, Lindström M, Möller S, Petitt E *et al.* (2012a). Psychodynamic guided self-help for adult depression through the internet: a randomised controlled trial. *PLoS ONE* **7**.
- Johansson R, Sjöberg E, Sjögren M, Johnsson E, Carlbring P, Andersson T *et al.* (2012b). Tailored vs. standardized internet-based cognitive behavior therapy for depression and comorbid symptoms: a randomized controlled trial. *PLoS ONE* **7**, 1–9.
- Jonkers CCM, Lamers F, Bosma H, Metsemakers JFM and van Eijk JTM (2012) The effectiveness of a minimal psychological intervention on self-management beliefs and behaviors in depressed chronically ill elderly persons: a randomized trial. *International Psychogeriatrics* **24**, 288–297.
- Kelly JA, Murphy DA, Bahr R, Kalichman SC, Morgan MG, Stevenson Y *et al.* (1993) Outcome of cognitive-behavioral and support group brief therapies for depressed, HIV-infected persons. *American Journal of Psychiatry* **150**, 1679–1686.
- Kivi M, Eriksson MCM, Hange D, Peterson E-L, Vernmark K, Johansson B *et al.* (2014) Internet-based therapy for mild to moderate depression in Swedish primary care: short term results from the PRIM-NET randomized controlled trial. *Cognitive Behaviour Therapy* **43**, 289–298.
- Korte J, Bohlmeijer ET, Cappeliez P, Smit F and Westerhof GJ (2012) Life review therapy for older adults with moderate depressive symptomatology: a pragmatic randomized controlled trial. *Psychological Medicine* **42**, 1163–1173.
- Kraemer HC and Kupfer DJ (2006) Size of treatment effects and their importance to clinical research and practice. *Biological Psychiatry* **59**, 990–996.
- Laidlaw K, Davidson K, Toner H, Jackson G, Clark S, Law J *et al.* (2008) A randomised controlled trial of cognitive behaviour therapy vs treatment as usual in the treatment of mild to moderate late life depression. *International Journal of Geriatric Psychiatry* **23**, 843–850.
- Lamers SMA, Bohlmeijer ET, Korte J and Westerhof GJ (2015) The efficacy of life-review as online-guided self-help for adults: a randomized trial. *Journals of Gerontology – Series B Psychological Sciences and Social Sciences* **70**, 24–34.
- Lemma A and Fonagy P (2013) Feasibility study of a psychodynamic online group intervention for depression. *Psychoanalytic Psychology* **30**, 367–380.
- Lynch D, Tamburrino M, Nagel R and Smith MK (2004) Telephone-based treatment for family practice patients with mild depression. *Psychological Reports* **94**, 785–792.
- Maina G, Forner F and Bogetto F (2005) Randomized controlled trial comparing brief dynamic and supportive therapy with waiting list condition in minor depressive disorders. *Psychotherapy and Psychosomatics* **74**, 43–50.
- Malouff JM, Thorsteinsson EB and Schutte NS (2007) The efficacy of problem solving therapy in reducing mental and physical health problems: a meta-analysis. *Clinical Psychology Review* **27**, 46–57.
- Martin PR, Aiello R, Gilson K, Meadows G, Milgrom J and Reece J (2015) Cognitive behavior therapy for comorbid migraine and/or tension-type headache and major depressive disorder: an exploratory randomized controlled trial. *Behaviour Research and Therapy* **73**, 8–18.
- Meyer TJ, Miller ML, Metzger RL and Borkovec TD (1990) Development and validation of the penn state worry questionnaire. *Behaviour Research and Therapy* **28**, 487–495.
- Milgrom J, Holt C, Holt CJ, Ross J, Ericksen J and Gemmill AW (2015) Feasibility study and pilot randomised trial of an antenatal depression treatment with infant follow-up. *Archives of Women's Mental Health* **18**, 717–730.
- Mitte K (2005) Meta-analysis of cognitive-behavioral treatments for generalized anxiety disorder: a comparison with pharmacotherapy. *Psychological Bulletin* **131**, 785–795.
- Naeem F, Gul M, Irfan M, Munshi T, Asif A, Rashid S *et al.* (2015) Brief culturally adapted CBT (CaCBT) for depression: a randomized controlled trial from Pakistan. *Journal of Affective Disorders* **177**, 101–107.
- Naeem F, Sarhandi I, Gul M, Khalid M, Aslam M, Anbrin A *et al.* (2014) A multicentre randomised controlled trial of a carer supervised culturally adapted CBT (CaCBT) based self-help for depression in Pakistan. *Journal of Affective Disorders* **156**, 224–227.
- Pot AM, Bohlmeijer ET, Onrust S, Melenhorst A-S, Veerbeek M and De Vries W (2010) The impact of life review on depression in older adults: a randomized controlled trial. *International Psychogeriatrics* **22**, 572–581.
- Pots WTM, Meulenbeek PAM, Veehof MM, Klungers J and Bohlmeijer ET (2014) The efficacy of mindfulness-based cognitive therapy as a public mental health intervention for adults with mild to moderate depressive symptomatology: a randomized controlled trial. *PLoS ONE* **9**, 1–12.
- Qiu J, Chen W, Gao X, Xu Y, Tong H, Yang M *et al.* (2013) A randomized controlled trial of group cognitive behavioral therapy for Chinese breast cancer patients with major depression. *Psychosomatic Obstetrics & Gynecology* **34**, 60–67.
- Richards D, Timulak L, O'Brien E, Hayes C, Vigano N, Sharry J *et al.* (2015) A randomized controlled trial of an internet-delivered treatment: its potential as a low-intensity community intervention for adults with symptoms of depression. *Behaviour Research and Therapy* **75**, 20–31.
- Sanchez-Meca J, Rosa-Alcazar AI, Marin-Martinez F and Gomez-Conesa A (2010) Psychological treatment of panic disorder with or without agoraphobia: a meta-analysis. *Clinical Psychology Review* **30**, 37–50.
- Savard J, Simard S, Giguère I, Ivers H, Morin CM, Maunsell E *et al.* (2006) Randomized clinical trial on cognitive therapy for depression in women

- with metastatic breast cancer: psychological and immunological effects. *Palliative & Supportive Care* **4**, 219–237.
- Serfaty MA, Haworth D and Buszewicz M** (2009) Clinical effectiveness of individual cognitive behavioral therapy for depressed older people in primary care. *Archives of General Psychiatry* **66**, 1332–1340.
- Simson U, Nawarotzky U, Friese G, Porck W, Schottenfeld-Naor Y, Hahn S *et al.*** (2008) Psychotherapy intervention to reduce depressive symptoms in patients with diabetic foot syndrome. *Diabetic Medicine* **25**, 206–212.
- Smits JAJ, Minhajuddin A and Jarrett RB** (2009) Cognitive therapy for depressed adults with comorbid social phobia. *Journal of Affective Disorders* **114**, 271–278.
- Strong V, Waters R, Hibberd C, Murray G, Wall L, Walker J *et al.*** (2008) Management of depression for people with cancer 4 (SMaRT oncology 1): a randomised trial. *Lancet* **372**, 40–48.
- Swartz HA, Frank E, Zuckoff A, Cyranowski JM, Houck PR, Cheng Y *et al.*** (2008) Brief interpersonal psychotherapy for depressed mothers whose children are receiving psychiatric treatment. *American Journal of Psychiatry* **165**, 1155–1162.
- Talbot N, Chaudron L, Ward EA, Duberstein PR, Conwell Y, O'Hara MW *et al.*** (2011) A randomized effectiveness trial of interpersonal psychotherapy for depressed women with sexual abuse histories. *Psychiatric Services* **62**, 374–380.
- Tovote KA, Flee J, Snippe E, Peeters ACTM, Emmelkamp PMG, Sanderman R *et al.*** (2014) Individual mindfulness-based cognitive therapy and cognitive behavior therapy for treating depressive symptoms in patients with diabetes: results of a randomized controlled trial. *Diabetes Care* **37**, 2427–2434.
- Vernmark K, Lenndin J, Bjärehed J, Carlsson M, Karlsson J, Öberg J *et al.*** (2010) Internet administered guided self-help versus individualized e-mail therapy: a randomized trial of two versions of CBT for major depression. *Behaviour Research and Therapy* **48**, 368–376.
- Vitriol VG, Ballesteros ST, Florenzano RU, Weil KP and Benadof DF** (2009) Evaluation of an outpatient intervention for women with severe depression and a history of childhood trauma. *Psychiatric Services* **60**, 936–942.
- von Hippel PT** (2015) The heterogeneity statistic I (2) can be biased in small meta-analyses. *BMC Medical Research Methodology* **15**, 2–8.
- Warmerdam L, van Straten A, Twisk J, Riper H and Cuijpers P** (2008) Internet-Based treatment for adults with depressive symptoms: randomized controlled trial. *Journal of Medical Internet Research* **10**, 1–16.
- Watkins ER, Taylor RS, Byng R, Baeyens C, Read R, Pearson K *et al.*** (2012) Guided self-help concreteness training as an intervention for major depression in primary care: a phase II randomized controlled trial. *Psychological Medicine* **42**, 1359–1371.
- Wierzbicki M and Bartlett TS** (1987) The efficacy of group and individual cognitive therapy for mild depression. *Cognitive Therapy and Research* **11**, 337–342.
- Wiethoff K, Bauer M, Baghai TC, Möller H-J, Fisher R, Hollinde D *et al.*** (2010) Prevalence and treatment outcome in anxious versus nonanxious depression: results from the German Algorithm Project. *The Journal of Clinical Psychiatry* **71**, 1047–1054.
- Zajecka JM and Ross JS** (1995) Management of comorbid anxiety and depression. *Journal of Clinical Psychiatry* **56**, 10–13.