cambridge.org/psm

Original Article

Cite this article: Chaplin AB, Daniels NF, Ples D, Anderson RZ, Gregory-Jones A, Jones PB, Khandaker GM (2023). Longitudinal association between cardiovascular risk factors and depression in young people: a systematic review and meta-analysis of cohort studies. *Psychological Medicine* **53**, 1049–1059. https://doi.org/10.1017/S0033291721002488

Received: 19 January 2021 Revised: 21 May 2021 Accepted: 3 June 2021

First published online: 25 June 2021

Keywords:

Body mass index; cardiovascular risk factor; depression; depressive symptoms; meta-analysis; smoking

Author for correspondence:

Anna B. Chaplin,

E-mail: ac2114@medschl.cam.ac.uk

© The Author(s), 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (http://creativecommons.org/licenses/by/4.0), which permits unrestricted re- use, distribution and reproduction, provided the original article is properly cited.



Longitudinal association between cardiovascular risk factors and depression in young people: a systematic review and meta-analysis of cohort studies

Anna B. Chaplin¹ , Natasha F. Daniels² , Diana Ples² , Rebecca Z. Anderson^{3,4}, Amy Gregory-Jones², Peter B. Jones^{1,5} and Golam M. Khandaker^{1,5,6,7,8}

¹Department of Psychiatry, University of Cambridge, Cambridge, UK; ²School of Clinical Medicine, University of Cambridge, Cambridge, UK; ³Royal Liverpool University Hospital, Liverpool, UK; ⁴Liverpool NHS Foundation Trust, Liverpool, UK; ⁵Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, UK; ⁶MRC Integrative Epidemiology Unit, Population Health Science, University of Bristol, Bristol, UK; ⁷Centre for Academic Mental Health, University of Bristol, Bristol, UK and ⁸Avon and Wiltshire Mental Health Partnership NHS Trust, Bristol, UK

Abstract

Background. Depression is a common and serious mental illness that begins early in life. An association between cardiovascular disease (CVD) and subsequent depression is clear in adults. We examined associations between individual CVD risk factors and depression in young people.

Methods. We searched MEDLINE, EMBASE, and PsycINFO databases from inception to 1 January 2020. We extracted data from cohort studies assessing the longitudinal association between CVD risk factors [body mass index (BMI), smoking, systolic blood pressure (SBP), total cholesterol, high-density lipoprotein] and depression, measured using a validated tool in individuals with mean age of 24 years or younger. Random effect meta-analysis was used to combine effect estimates from individual studies, including odds ratio (OR) for depression and standardised mean difference for depressive symptoms.

Results. Based on meta-analysis of seven studies, comprising 15 753 participants, high BMI was associated with subsequent depression [pooled OR 1.61; 95% confidence interval (CI) 1.21-2.14; $I^2=31\%$]. Based on meta-analysis of eight studies, comprising 30 539 participants, smoking was associated with subsequent depression (pooled OR 1.73; 95% CI 1.36–2.20; $I^2=74\%$). Low, but not high, SBP was associated with an increased risk of depression (pooled OR 3.32; 95% CI 1.68–6.55; $I^2=0\%$), although this was based on a small pooled high-risk sample of 893 participants. Generalisability may be limited as most studies were based in North America or Europe.

Conclusions. Targeting childhood/adolescent smoking and obesity may be important for the prevention of both CVD and depression across the lifespan. Further research on other CVD risk factors including blood pressure and cholesterol in young people is required.

Introduction

Depression is a common and serious mental illness with a lifetime risk of 10–20% (Hasin et al., 2018). The majority of depression cases are established by age 24 (Kessler et al., 2010) and this condition is the leading cause of disability among children and young people (Polanczyk, Salum, Sugaya, Caye, & Rohde, 2015). Following an initial depressive episode, the risk of recurrence is 60% (Holtzheimer & Mayberg, 2011). Therefore, early-onset depression is associated with a longer period of risk for relapse as well as poor long-term outcomes (Holtzheimer & Mayberg, 2011; Kessler, 2012). A better understanding of the aetiology of depression in young people is required to develop effective strategies for prevention and treatment (Niarchou, Zammit, & Lewis, 2015).

Cardiovascular disease (CVD) is a leading cause of health-related disability worldwide (Roth et al., 2018). There is evidence for bidirectional associations between CVD and depression in adults (Hiles et al., 2015; Inouye et al., 2018; Khandaker et al., 2019; Smolderen et al., 2017). A substantial body of literature suggests that depression is a key risk factor for CVD in adults and that it may predict poor outcomes following a cardiac event (Barefoot & Schroll, 1996; Hiles et al., 2015; Inouye et al., 2018; Khandaker et al., 2019; Lippi, Montagnana, Favaloro, & Franchini, 2009; Van der Kooy et al., 2007). CVD is also associated with subsequent depression in adults (Choi, Kim, Marti, & Chen, 2014; Hare, Toukhsati, Johansson, & Jaarsma, 2014; Kendler, Gardner, Fiske, & Gatz, 2009; Lippi et al., 2009). However, studies



of CVD risk and subsequent depression in young people are relatively less common. A clearer understanding of the association between CVD risk factors and depression in young people is required. Early detection and management of CVD risk factors may reduce risks for both CVD and depression subsequently during the life-course.

The World Health Organization defines young people as individuals aged 24 years or younger (WHO Study Group of Young People, 1986). Existing studies of CVD risk and depression in young people have often focused on individual risk factors such as body mass index (BMI) or smoking. In the past decade, a number of systematic reviews have highlighted an association between obesity in young people and depression across the lifespan (Hoare, Skouteris, Fuller-Tyszkiewicz, Millar, & Allender, 2014; Mannan, Mamun, Doi, & Clavarino, 2016; Mühlig, Antel, Föcker, & Hebebrand, 2016; Sutaria, Devakumar, Yasuda, Das, & Saxena, 2019). However, none of these studies specifically examined depression risk in young people. A recent systematic review of individuals aged 14-35 reported that childhood obesity is associated with approximately 50% increased risk of depression (Sutaria et al., 2019). Another review reported that smoking in early life is associated with 73% increased risk of depression in young people (Chaiton, Cohen, O'Loughlin, & Rehm, 2009). According to the Framingham study, other established CVD risk factors for adults include systolic blood pressure (SBP), total cholesterol, and highdensity lipoprotein (HDL), in addition to smoking and BMI (Wilson et al., 1998; Wilson, Castelli, & Kannel, 1987). These CVD risk factors are all potentially modifiable and may be important in the aetiology and prevention of depression. CVD risk factors are increasingly being examined in young people; thus, a systematic review is required to summarise these findings.

We conducted a systematic review and meta-analysis of existing studies to quantify the longitudinal association of five key CVD risk factors (BMI, smoking, SBP, total cholesterol, and HDL) and depression in young people. These CVD risk factors were chosen for a number of reasons: (i) they are part of the Framingham Cardiovascular Risk Score for adults; (ii) they are potentially modifiable; and (iii) they remain relevant in the context of young people. Our outcome was depression (binary or continuous) assessed using a validated tool. We also performed a number of sensitivity analyses, for example by excluding studies that only looked at one gender or excluding studies based on quality assessment.

Methods

Search strategy and study selection

This study has been performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Details of the protocol were prospectively registered on PROSPERO (see https://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42020172460).

MEDLINE, EMBASE, and PsycINFO databases were searched to identify all relevant studies of the association between CVD risk factors and depression from database inception to 1 January 2020. The following keywords were used: '(cohort OR longitudinal OR prospective OR retrospective OR follow up stud*) AND depress* AND (adolescen* OR young person OR young people OR child* OR infant OR early adult OR youth* OR teen*) AND ((cardiovascular AND risk) OR total cholesterol OR high density lipoprotein OR hdl OR smok* OR bmi OR body mass index OR adiposity OR waist circumference OR body fat distribution OR skinfold

thickness OR lipid accumulation product OR systolic blood pressure OR systolic bp OR sbp)'. See online Supplementary materials for the full search strategy. No language restriction was applied. The electronic search was complemented by hand-searching the reference lists of included studies. All titles and abstracts were examined to retrieve potentially relevant studies. ABC, NFD, AGJ, DP, and RZA applied inclusion/exclusion criteria and selected the final studies for this review.

Selection criteria

We included studies that: (i) had a longitudinal population-based cohort design (prospective or retrospective); (ii) included participants with a mean age of 24 years old or younger at follow-up; (iii) had at least one of the five CVD risk factors (BMI, smoking, SBP, total cholesterol, HDL) as the exposure at baseline; (iv) used a validated tool to measure depression (binary outcome or symptom score) at follow-up; and (v) reported effect estimate(s) for the association between CVD risk and subsequent depression. Studies were excluded if they did not have an unexposed group for a particular risk factor (e.g. experimental smoking used as the comparison group rather than no smoking), had depression as the exposure and the CVD risk factor as the outcome, or measured depression comorbid with another mental illness such as bipolar disorder or anxiety.

Data extraction

Data extraction was performed independently by ABC, NFD and DP, and disagreements were resolved by consensus. For each included study, we extracted the following data: (i) details of the cohort (country, name/setting, design, sample size, and follow-up length); (ii) assessment of exposure and outcome; (iii) age and sex of the included participants; and (iv) results of analysis (number of participants exposed at baseline, number of participants with depression at follow-up, adjusted/unadjusted effect estimates). When studies reported various methods for assessing the exposure or repeated measures of the exposure, we used the most comprehensive measure. For example, one study measured BMI eight times from birth to age 12 years (Wang, Leung, & Schooling, 2014). We chose the age 7 measure as the earliest age where BMI may be an appropriate measure of central adiposity to maximise the length of follow-up. In cases where there was more than one published report from the same population, we included the study with the larger sample size (Bares, 2014; Duncan & Rees, 2005; Goodman & Capitman, 2000; Munafò, Hitsman, Rende, Metcalfe, & Niaura, 2008). Some studies reported results where depression at baseline was adjusted for as well as analysis where baseline depression cases were removed. In such cases, we only included results where baseline depression was excluded to minimise reverse causality.

Data synthesis and meta-analysis

We performed separate meta-analyses for BMI, smoking, and SBP. Results from studies were pooled using the inverse variance method meaning that studies with larger sample sizes were given greater weight (Higgins et al., 2021; Schwarzer, 2007). Results for other CVD risk factors were summarised using a narrative review. Meta-analyses were performed separately for studies that reported beta estimates (continuous depressive symptoms outcome) or odds ratio (OR) (binary depression outcome). We used random-effect meta-analysis, which is appropriate where there

is heterogeneity between the studies. Heterogeneity between studies was examined using the I^2 statistic.

We assessed publication bias by visual inspection of funnel plots, and by Egger's regression test for funnel plot asymmetry (mixed-effects meta-regression model). We considered a p value of <0.05 to indicate the existence of publication bias.

We assessed study quality using the Newcastle-Ottawa Scale for cohort studies (Stang, 2010). We repeated analyses with only good/fair quality studies. We also repeated analyses after excluding studies with only female or male participants. Meta-analyses were carried out using the *meta* package version 4.11 in R version 3.6.1 (Schwarzer, 2007).

Results

Electronic search identified 6616 studies. After removing duplicates, 4821 studies remained. After title and abstract screening, 197 (4.1%) potentially eligible studies were identified, of which 29 met the inclusion criteria and were included in the review (Albers & Biener, 2002; Bares, 2014; Beal, Negriff, Dorn, Pabst, & Schulenberg, 2014; Boutelle, Hannan, Fulkerson, Crow, & Stice, 2010; Chaiton, Cohen, Rehm, Abdulle, & O'Loughlin, 2015; Chang et al., 2017; Choi, Patten, Christian Gillin, Kaplan, & Pierce, 1997; Clark et al., 2007; Duncan & Rees, 2005; Eitle & Eitle, 2018; Frisco, Houle, & Lippert, 2013; Gage et al., 2015; Gomes et al., 2019; Goodman & Whitaker, 2002; Hammerton, Harold, Thapar, & Thapar, 2013; Hammerton, Thapar, & Thapar, 2014; Marmorstein, Iacono, & Legrand, 2014; Monshouwer et al., 2012; Perry et al., 2020; Piumatti, 2018; Pryor et al., 2016; Raffetti, Donato, Forsell, & Galanti, 2019; Ranjit et al., 2019a, b; Rhew et al., 2008; Roberts & Duong, 2013; Rubio, Kraemer, Farrell, & Day, 2008; Wang et al., 2014; Zhang, Woud, Becker, & Margraf, 2018). See Fig. 1 for study selection and Table 1 for characteristics of included studies.

All studies were prospective in design, except one which used a retrospective measure of depression at follow-up (Monshouwer et al., 2012). The majority of studies (55.2%) were rated as 'good' quality using the Newcastle-Ottawa Scale (online Supplementary Tables S1 and S2). Sex, age, parental education, race/ethnicity, baseline depression, and alcohol use were the most commonly used confounders (online Supplementary Table S3).

Based on data availability, meta-analysis for BMI included 13 studies (Boutelle et al., 2010; Chang et al., 2017; Clark et al., 2007; Eitle & Eitle, 2018; Frisco et al., 2013; Gomes et al., 2019; Goodman & Whitaker, 2002; Marmorstein et al., 2014; Monshouwer et al., 2012; Pryor et al., 2016; Rhew et al., 2008; Roberts & Duong, 2013; Zhang et al., 2018), and that for smoking included 11 studies (Albers & Biener, 2002; Bares, 2014; Beal et al., 2014; Chaiton et al., 2015; Choi et al., 1997; Clark et al., 2007; Duncan & Rees, 2005; Gage et al., 2015; Piumatti, 2018; Raffetti et al., 2019; Ranjit et al., 2019a, b; Rubio et al., 2008; Zhang et al., 2018) (online Supplementary Table S2). Meta-analysis for SBP included one study comprising two separate samples (Hammerton et al., 2013). We found no studies of total cholesterol or HDL that met our inclusion criteria.

Meta-analysis of adjusted effect estimates

Longitudinal association between high BMI at baseline and risk of depression at follow-up

Based on seven studies reporting an adjusted OR, comprising a total of 15 753 participants, the pooled OR for depression at

follow-up associated with high BMI (>25) at baseline was 1.61 (95% confidence interval (CI) 1.21–2.14) (Fig. 2). There was limited evidence of heterogeneity between studies ($I^2 = 31\%$; 95% CI 0–71%; Cochran's Q = 8.7; p = 0.19). Separate meta-analysis of unadjusted effect estimates showed lower pooled results (online Supplementary Fig. S1).

Longitudinal association between smoking at baseline and risk of depression at follow-up

Based on eight studies reporting an adjusted OR, comprising a total of 30 539 participants, the pooled OR for depression at follow-up associated with smoking at baseline was 1.73 (95% CI 1.36–2.20) (Fig. 3). There was evidence of heterogeneity between studies ($I^2 = 74\%$; 95% CI 52–86%; Cochran's Q = 35.3; p < 0.01). Separate meta-analysis of unadjusted effect estimates showed higher pooled results (online Supplementary Fig. S2).

Longitudinal association between SBP at baseline and risk of depression at follow-up

One study examined associations of both low and high SBP with depression in two separate samples comprising a total of 5111 participants. Meta-analysis of these studies suggest depression at follow-up is associated with low SBP at baseline (OR 3.32; 95% CI 1.68–6.55), but not with high SBP at baseline (OR 0.82; 95% CI 0.55–1.22) (Fig. 4). There was some evidence of heterogeneity for high SBP ($I^2 = 66\%$; 95% CI 0–92%; Cochran's Q = 3.0; p = 0.08) and little heterogeneity for low SBP ($I^2 = 0\%$; 95% CI 0–0%; Cochran's Q = 0.04; p = 0.84).

Longitudinal association between high BMI, smoking at baseline and depressive symptoms at follow-up

Based on five studies reporting an adjusted beta estimate, comprising a total of 11 516 participants, the standardised mean difference (SMD) for an increase in depressive symptoms at follow-up associated with high BMI at baseline was 0.05 (95% CI –0.08–0.18) (online Supplementary Fig. S3). There was evidence of heterogeneity between studies ($I^2 = 71\%$; 95% CI 34–88%; Cochran's Q = 17.5; p < 0.01).

Based on five studies reporting an adjusted beta estimate, comprising a total of 21 490 participants, the SMD for an increase in depressive symptoms at follow-up associated with smoking at baseline was 0.37 (95% CI 0.10–0.64) (online Supplementary Fig. S3). There was evidence of heterogeneity between studies ($I^2 = 89\%$; 95% CI 78–94%; Cochran's Q = 44.8; p < 0.01).

Results for sensitivity analysis

After excluding three studies with only female participants (Boutelle et al., 2010; Frisco et al., 2013; Zhang et al., 2018), the adjusted pooled OR for depression at follow-up for high BMI at baseline was 1.43 (95% CI 0.94–2.18) (Fig. 2). There was some heterogeneity between studies ($I^2 = 47\%$; 95% CI 0–83%; Cochran's Q = 5.7; p = 0.13).

After excluding four studies with only female or male participants (Choi et al., 1997; Duncan & Rees, 2005; Rubio et al., 2008; Zhang et al., 2018), the pooled adjusted OR for depression at follow-up for smoking at baseline was 1.23 (95% CI 1.02–1.49) (Fig. 3). There was little heterogeneity ($I^2 = 11\%$; 95% CI 0–86%; Cochran's Q = 3.4; p = 0.34).

After excluding one study based on quality assessment (Duncan & Rees, 2005), the pooled adjusted OR for depression




Fig. 1. PRISMA flow diagram for study selection.

at follow-up for smoking at baseline was 1.48 (95% CI 1.21–1.80) (online Supplementary Fig. S4). There was some heterogeneity ($I^2 = 38\%$; 95% CI 0–73%; Cochran's Q = 11.3; p = 0.12). For BMI, we did not exclude any studies based on quality assessment.

See online Supplementary Figs S5 and S6 for sensitivity analyses where the outcome of interest was depressive symptoms at follow-up.

Publication bias

Based on Egger's test and funnel plots, evidence for publication bias was not evident for studies reporting the adjusted association between high BMI and depression (Egger's test: p = 0.17; online Supplementary Fig. S7), smoking and depression (Egger's test: p = 0.80; online Supplementary Fig. S8), or high BMI and depressive

symptoms (Egger's test: p = 0.35) (online Supplementary Fig. S9). Evidence for publication bias was present for studies reporting the adjusted association between smoking and depressive symptoms (Egger's test: p = 0.01) (online Supplementary Fig. S10).

Discussion

Depression and CVD are associated with each other in mid to late adulthood. Although depression is known to arise commonly in young people, the timing of the association with CVD risk is unclear and common risk factors for the two conditions raise the prospect of joint prevention. To the best of our knowledge, this is the first systematic review to consider the association between various CVD risk factors and subsequent depression in young people. We report four key findings: (i) BMI and smoking

Table 1. Characteristics of studies included in systematic review

Study	Country	Female, No. (%)	Baseline age range/mean (years)	Minimum follow-up (years)	Assessment of exposure	Exposed at baseline (%) or mean (s.d.)	Assessment of depression	Depression at follow-up (%)	Mean (s.b.) depressive symptom score at follow-up
ВМІ									
Boutelle et al. (2010)	USA	495 (100) ^a	13.5	1	Obese v. healthy weight	10.7%	Structured interview (K-SADS)	1.2-3.5	1.3-1.4 (0.4)
Chang et al. (2017)	Taiwan	969 (51.2)	14.7	3	Obese <i>v.</i> healthy weight	3.5% (F); 3.4% (M)	Self-report (CES-DC)	-	1.8 (0.5) (F); 1.5 (0.4) (M)
Clark et al. (2007)	UK	776 (51.3)	11–14	2	Obese <i>v</i> . healthy weight	19.6%	Self-report (SMFQ)	15.1	-
Eitle and Eitle (2018)	USA	308 (47.0) (AI); 3403 (50.0) (W)	12-18	1	Obese v. not obese	19.0% (AI); 11.0% (W)	Self-report (CES-D)	-	12.4 (0.6) (AI); 10.1 (0.2) (W)
Frisco et al. (2013)	USA	5243 (100)	15.7	4	Consistently obese <i>v</i> . healthy weight	9.8%	Self-report (CES-D)	7.0	-
Gomes et al. (2019)	Brazil	NM	18	4	Obese v. not obese	6.7%	Structured interview (MINI-5)	2.9	-
Goodman and Whitaker (2002)	USA	4718 (48.6)	<20	1	Obese v. not obese	9.7%	Self-report (CES-D)	8.9	-
Hammerton et al. (2014) ^b	UK	EPAD: 165 (57.1);ALSPAC: 2533 (52.1)	EPAD: 9-17; ALSPAC: 11- 14	,	Standardised score	EPAD: 17.4%; ALSPAC: 23.7%	EPAD: Semi-structured interview (CAPA); ALSPAC: Structured interview (DAWBA)	EPAD: 8.3; ALSPAC: 1.6	EPAD: 2.2 (NM) (F); 1.6 (NM) (M);ALSPAC: NM
Marmorstein et al. (2014)	USA	NM	11-14	6	Obese v. not obese	14.7% (F); 10.7% (M)	Structured interview (SCID)	12.3 (F); 7.6 (M)	-
Monshouwer et al. (2012)	Netherlands	820 (53.3)	10-12	4	Obese <i>v.</i> healthy weight	14.1%	Structured interview (WHO CIDI-3)	5.6	-
Perry et al. (2020) ^b	UK	1655 (51.6)	9	9	Raw score	17.5 (2.5)	Self-report (CIS-R)	7.0	3.1 (NM)
Piumatti (2018) ^b	Italy	178 (78.1)	21.4	1	Raw score	21.1 (2.7)	Self-report (PHQ-2)	-	1.1 (0.6)
Pryor et al. (2016)	Canada	661 (54.1)	6–12	1	Early-onset <i>v</i> . never overweight	11.0%	Self-report (Kovacs CDI)	-	NM
Rhew et al. (2008)	USA	206 (46.2)	12	1	Overweight <i>v.</i> healthy weight	22.4% (F); 30.3% (M)	Structured interview (MFQ)	-	NM
Roberts and Duong (2013)	USA	2040 (48.9) ^a	11–17	1	Obese v. healthy weight	19.7%	Structured interview (DISC IV-Y)	1.7	-
Wang et al. (2014) ^b	Hong Kong	2793 (48.2)	7	7	Standardised score	-0.03 (1.1) (F); 0.3 (1.3) (M)	Self-report (PHQ-9)	=	4.0 (NM)
Zhang et al. (2018)	Germany	1196 (100)	21	1.4	Overweight <i>v</i> . healthy weight	7.2%	Structured interview (Diagnostic Interview for Mental Disorders Research Version)	6.5	-

1054

Table 1. (Continued.)

Study	Country	Female, No. (%)	Baseline age range/mean (years)	Minimum follow-up (years)	Assessment of exposure	Exposed at baseline (%) or mean (s.d.)	Assessment of depression	Depression at follow-up (%)	Mean (s.p.) depressive symptom score at follow-up
Smoking									
Albers and Biener (2002)	USA	259 (49.6) ^a	12-15	4	Ever v. never smoker	33.9%	Self-report (Kandel & Davies)	13.2	-
Bares (2014)	USA	2486 (53.0) ^a	16.6	5.5	Unit increase in cigarette use	31.3%	Self-report (CES-D)	-	4.9 (3.9)
Beal et al. (2014)	USA	262 (100)	11-20	2	Ever v. never smoker	2.9 (3.1)	Self-report (CDI/BDI)	-	46.3 (10.8)
Chaiton et al. (2015) ^b	Canada	416 (49.7)	12-13	5	Smoking initiation <i>v</i> . no initiation	47.4%	Self-report (Mellinger scale)	23.8	-
Choi et al. (1997)	USA	3215 (46.8)	12-18	4	Established <i>v.</i> never smoker	17.3%	Self-report (Kandel & Davies)	11.5	-
Clark et al. (2007)	UK	776 (51.3)	11-14	2	Tried/regular <i>v</i> . never smoker	35.2%	Self-report (SMFQ)	15.1	-
Duncan and Rees (2005)	USA	6748 (51.6)	11-21	1	Smoker v. non-smoker	25.8% (F); 25.3% (M)	Self-report (CES-D)	27.0 (F); 21.4 (M)	13.0 (8.7) (F);10.8 (7.5) (M)
Gage et al. (2015)	UK	NM	16	2	Unit increase in cigarette use	9.8%	Self-report (CIS-R)	7.2	-
Raffetti et al. (2019)	Sweden	1636 (51.2)	13-14	1	Current <i>v</i> . non-smoker	2.0%	Self-report (CES-DC)	8.3	15.7 (NM)
Ranjit et al. (2019 <i>a</i>) ^b	Finland	2358 (50.3) ^a	14	3	Regular <i>v</i> . never smoker	9.1%	Self-report (GBI)	5.1 (4.9)	-
Ranjit et al. (2019 <i>b</i>) ^b	Finland	2174 (51.8) ^a	17.5	5	Ever v. never smoker	25.3%	Self-report (GBI)	4.5 (4.7)	-
Rubio et al. (2008)	USA	278 (100)	23	0.25	Smoker v. non-smoker	72.0%	Self-report (CES-D)	85.0	-
Piumatti (2018)	Italy	178 (78.1)	21.4	1	Daily v. non-smokers	13.6%	Self-report (PHQ-2)	-	1.1 (0.6)
Zhang et al. (2018)	Germany	1196 (100)	21	1.4	Current <i>v</i> . non-smoker	23.6%	Structured interview (Diagnostic Interview for Mental Disorders Research Version)	6.5	-
Systolic blood pressure									
Hammerton et al. (2013)	UK	EPAD: 164 (58.4); ALSPAC: 2516 (52.1)	EPAD: 9-17; ALSPAC: 11- 14	,	Standardised score	EPAD: 117.3 (13.2); ALSPAC: 111.0 (9.5)	EPAD: Semi-structured interview (CAPA); ALSPAC: Structured interview (DAWBA)	EPAD: 8.5; ALSPAC: 1.6	-

s.b., standard deviation; F, female; M, male; Al, American Indian; W, White; EPAD, Early Prediction of Adolescent Depression study; ALSPAC, Avon Longitudinal Study of Parents and Children; NS, not significant; NM, not mentioned; K-SADS, Kiddie Schedule for Affective Disorders and Schizophrenia; SMFQ, Short Mood and Feelings Questionnaire; MINI-5, Mini International Neuropsychiatric Interview Version Five; DSM, Diagnostic and Statistical Manual of Mental Disorders; CES-D, Centre for Epidemiological Studies Depression; CAPA, Child and Adolescent Psychiatric Assessment (Child Version); DAWBA, Development and Wellbeing Assessment (Child Version); SCID, Structured Clinical Interview for DSM-III-R; CIS-R, Clinical Interview Schedule Revised; ICD-10, International Statistical Classification of Diseases and Related Health Problems Tenth Revision; DISC IV-Y, Diagnostic Interview Schedule for Children for direct administration to children or adolescents; WHO CIDI-3, World Health Organisation Composite International Diagnostic Interview Version Three; CDI, Children's Depression Inventory; PHQ, Patient Health Questionnaire; MFQ, Mood and Feelings Questionnaire; GBI, General Behaviour Inventory; BDI, Beck's Depression Inventory.

*Baseline sample only.**

^bNot included in meta-analysis. The four studies not included in BMI meta-analysis were excluded because they measured BMI as a continuous variable. The three studies not included in smoking meta-analysis were excluded because they did not report effect estimates comparable with the other studies.




Fig. 2. Meta-analysis of longitudinal association between high BMI at baseline and subsequent depression in young people.




Fig. 3. Meta-analysis of longitudinal association between smoking at baseline and subsequent depression in young people.

are the most well-studied risk factors for depression in this age group; (ii) both BMI and smoking at baseline are longitudinally associated with subsequent depression; (iii) smoking but not BMI is prospectively associated with depressive symptoms; and (iv) currently there is limited data on longitudinal associations of high SBP and cholesterol with subsequent depression in young people, which should be examined in future.

Our results suggest that obesity could be an important risk factor for depression in young people. The pooled OR of 1.61 for the

association of high BMI and depression is remarkably similar to previous studies in adults. Previous meta-analyses have reported ORs of 1.51 and 1.70 for the prospective association between childhood high BMI and adult depression (Luppino et al., 2010; Sutaria et al., 2019). Another meta-analysis reported that obese adolescents had an 40% increased risk of experiencing depression as adults (Mannan et al., 2016). A recent Mendelian randomisation study using data from 812 000 adult participants also found that fat mass could be a causal factor for depression




Fig. 4. Meta-analysis of longitudinal association between SBP at baseline and subsequent depression in young people. SBP, systolic blood pressure; EPAD, Early Prediction of Adolescent Depression study; ALSPAC, Avon Longitudinal Study of Parents and Children. *Subset of ALSPAC participants with mothers with recurrent depression. †EPAD participants have mother/father with recurrent depression.

(Speed, Jefsen, Børglum, Speed, & Østergaard, 2019). We did not find an association between high BMI and depressive symptoms score, indicating a possibly non-linear association between BMI and depression whereby association is restricted to only those with more severe symptoms.

Our findings also suggest that smoking could be a risk factor for depression in young people. The pooled OR of 1.73 for the association between smoking and depression is consistent with a previous meta-analysis in adults, which reported an OR of 1.62 (Luger, Suls, & Vander Weg, 2014). Similarly, a meta-analysis of nine cross-sectional and longitudinal studies found that adolescents exposed to second-hand smoking had increased odds of depression (Han, Liu, Gong, Ye, & Zhou, 2019). However, current evidence from observational cohort studies and genetic Mendelian randomisation studies reports mixed findings regarding the association between smoking and depression, with some reporting an association (Wootton et al., 2020) and others no association (Bjørngaard et al., 2013; Taylor et al., 2014). Therefore, residual confounding or reverse causality remain viable explanations for the observed association between smoking and depression. Further longitudinal studies and genetic Mendelian randomisation studies are required to investigate this issue.

A number of potential mechanisms may be involved in the association of high BMI and subsequent depression, including lowgrade systemic inflammation, hypothalamic-pituitary-adrenal axis (HPA) axis dysregulation, insulin resistance, and psychological distress. Inflammation is evident in around 25% of individuals with depression (Osimo, Baxter, Lewis, Jones, & Khandaker, 2019) and atypical depression is associated with inflammation and metabolic dysregulation (Lamers et al., 2013, 2020). Adipose tissue also contains abundant inflammatory cytokines that are involved in fat metabolism (Heredia, Gómez-Martínez, & Marcos, 2012). Similarly, the role of insulin in regulating adipocyte function contributes to the close link between insulin resistance and obesity (Kahn & Flier, 2000). Furthermore, melancholic depression, higher levels of abdominal fat, HPA axis hyperactivity, and cortisol dysregulation are inter-related (Incollingo Rodriguez et al., 2015; Lamers et al., 2013, 2020). Adverse childhood experiences are also one of the most robust risk factors for depression (Gardner, Thomas, & Erskine, 2019), and are associated with increased risk for obesity and metabolic dysregulation (Farr et al., 2015). Low-grade systemic inflammation may be important to the association between

smoking and depression (Berk et al., 2013). Studies are required to investigate the complex mechanisms that may underlie the association between high BMI and depression.

Low SBP, but not high SBP, appeared to be associated with risk for depression in young people. However, our meta-analysis was based on only two cohorts at high risk for depression and further study is required. Low SBP appears to be associated with depression in young people at high-risk of depression but not in the general population (Hammerton et al., 2013). Low SBP has also been associated with depression in cross-sectional and longitudinal studies of middle-aged and elderly adults (Hildrum et al., 2007; Huang, Su, Jiang, & Zhu, 2020). Conversely, higher SBP has been prospectively associated with fewer depressive symptoms in older adults with CVD risk factors (Herrmann-Lingen et al., 2018). In adult populations, the relationship between SBP and depression may be independent of a range of lifestyle factors, age, and sex (Hildrum et al., 2007; Huang et al., 2020).

The reason why low SBP has a potentially causal role in depression remains unclear. Neurons controlling blood pressure could be implicated in the association between SBP and depression. Neuropeptide Y, for example, reduces blood pressure and is involved in stress responses that have been linked to increased risk for depression, such as the HPA axis (Hildrum et al., 2007; Juruena, Bocharova, Agustini, & Young, 2018). Further research is required to understand the relationship between SBP and depression in young people, including potential mechanisms.

We found no studies assessing the association between either total cholesterol or HDL on risk for subsequent depression in young people. A meta-analysis of 30 cross-sectional studies reported that higher total cholesterol was associated with lower levels of depression in adults (Shin, Suls, & Martin, 2008). Evidence from adults indicates both higher and lower HDL to be associated with increased risk for depression in adults. In a meta-analysis of 16 cross-sectional studies, high HDL was related to higher levels of depression, especially in women (Shin et al., 2008). Conversely, a meta-analysis of 11 case-control studies reported that lower HDL levels may be associated with firstepisode major depressive disorder in adults (Wei et al., 2020). Given that abnormal HDL, LDL and triglyceride levels are increasingly common in adolescents [Centers for Disease Control and Prevention (CDC), 2010], effort should be made to study potential effects on mental health as well as physical health.

Strengths of this study include the systematic literature search which identified a large number of relevant studies comprising a total of 93 021 participants. We included studies considering the effect of various CVD risk factors on either binary or continuous measures of depression/depressive symptoms. We assessed the studies using the validated Newcastle-Ottawa Scale as well as conducting sensitivity analyses to examine the robustness of our findings. However, this study is not without limitations. First, the majority of studies came from North America and Europe, limiting the generalisability of the results to other parts of the world. The number of studies in each of the meta-analyses was also relatively small, which resulted in wide CIs for the pooled effect estimates, and reduced the statistical power to detect publication bias. There was a considerable amount of heterogeneity between studies, particularly studies of depressive symptoms. Sensitivity analyses revealed that sex explained heterogeneity in some of the meta-analyses. However, stratifying by sex decreased the sample size, and consequently, statistical power to detect an association. In future, studies with larger samples are required. Finally, the possibility of residual confounding by unidentified factors remains high so we are cautious in terms of any conclusion regarding causality. Although studies included in this meta-analysis controlled for various potential confounding effects, other factors may also explain these associations. Further research is needed to examine whether observed associations are likely to be causal. Since randomised controlled trials are neither feasible nor ethical for some of the exposures under investigation (e.g. smoking, obesity), genetic approaches to dealing with residual confounding, such as Mendelian randomisation, would be particularly useful.

In summary, we present evidence for a longitudinal association between CVD risk factors, namely high BMI and smoking, in childhood/adolescence and subsequent depression in young people. These risk factors could be important targets for the prevention of depression and CVD in young people and subsequently during the life course. Further study is needed to understand potential mechanisms for these associations as well as the relationship between other CVD risk factors, notably blood pressure and cholesterol and depression risk in young people.

Supplementary material. The supplementary material for this article can be found at https://doi.org/10.1017/S0033291721002488.

Financial support. ABC is supported by the National Institute for Health Research (NIHR) CLAHRC RCF (MRR73-1795-00000) (https://www.nihr.ac.uk/), NIHR ARC East of England, the MQ: Transforming Mental Health (Data Science Award; grant code: MQDS17/40) (https://www.mqmental-health.org/home/). GMK acknowledges funding support from the Wellcome Trust (grant code: 201486/Z/16/Z) (https://wellcome.org/), MQ as above, the Medical Research Council (grant code: MC_PC_17213 and MR/S037675/1) (https://mrc.ukri.org/), and the BMA Foundation (J Moulton grant 2019) (http://www.bmafoundationmr.org.uk/). PBJ acknowledges funding from MQ, NIHR ARC East of England and the Medical Research Council, as above, and from NIHR PGfAR 0616-20003. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Registration number: CRD42020172460.

Conflict of interest. None.

References

Albers, A. B., & Biener, L. (2002). The role of smoking and rebelliousness in the development of depressive symptoms among a cohort of Massachusetts adolescents. *Preventive Medicine*, 34(6), 625–631. doi: 10.1006/pmed.2002.1029. Barefoot, J. C., & Schroll, M. (1996). Symptoms of depression, acute myocardial infarction, and total mortality in a community sample. *Circulation*, 93 (11), 1976–1980. doi: 10.1161/01.CIR.93.11.1976.

- Bares, C. B. (2014). Gender, depressive symptoms and daily cigarette use. *Journal of Dual Diagnosis*, 10(4), 187–196. doi: 10.1080/15504263.2014.961852.
- Beal, S. J., Negriff, S., Dorn, L. D., Pabst, S., & Schulenberg, J. (2014). Longitudinal associations between smoking and depressive symptoms among adolescent girls. *Prevention Science*, 15(4), 506–515. doi: 10.1007/ s11121-013-0402-x.
- Berk, M., Williams, L. J., Jacka, F. N., O'Neil, A., Pasco, J. A., Moylan, S., ... Maes, M. (2013). So depression is an inflammatory disease, but where does the inflammation come from? *BMC Medicine*, 11(1), 200. doi: 10.1186/1741-7015-11-200.
- Bjørngaard, J. H., Gunnell, D., Elvestad, M. B., Davey Smith, G., Skorpen, F., Krokan, H., ... Romundstad, P. (2013). The causal role of smoking in anxiety and depression: A Mendelian randomization analysis of the HUNT study. *Psychological Medicine*, 43(4), 711–719. doi: 10.1017/S0033291712001274.
- Boutelle, K. N., Hannan, P., Fulkerson, J. A., Crow, S. J., & Stice, E. (2010). Obesity as a prospective predictor of depression in adolescent females. *Health Psychology*, 29(3), 293–298. doi: 10.1037/a0018645.
- Centers for Disease Control and Prevention (CDC) (2010). Prevalence of abnormal lipid levels among youths United States, 1999–2006. MMWR. Morbidity and Mortality Weekly Report, 59(2), 29–33.
- Chaiton, M. O., Cohen, J. E., O'Loughlin, J., & Rehm, J. (2009). A systematic review of longitudinal studies on the association between depression and smoking in adolescents. *BMC Public Health*, 9(1), 356. doi: 10.1186/ 1471-2458-9-356.
- Chaiton, M., Cohen, J. E., Rehm, J., Abdulle, M., & O'Loughlin, J. (2015).
 Confounders or intermediate variables? Testing mechanisms for the relationship between depression and smoking in a longitudinal cohort study.
 Addictive Behaviors, 42, 154–161. doi: 10.1016/j.addbeh.2014.11.026.
- Chang, L.-Y., Chang, H.-Y., Wu, W.-C., Lin, L. N., Wu, C.-C., & Yen, L.-L. (2017). Body mass index and depressive symptoms in adolescents in Taiwan: Testing mediation effects of peer victimization and sleep problems. International Journal of Obesity, 41(10), 1510–1517. doi: 10.1038/ijo.2017.111.
- Choi, N. G., Kim, J., Marti, C. N., & Chen, G. J. (2014). Late-life depression and cardiovascular disease burden: Examination of reciprocal relationship. The American Journal of Geriatric Psychiatry, 22(12), 1522–1529. doi: 10.1016/j.jagp.2014.04.004.
- Choi, W. S., Patten, C. A., Christian Gillin, J., Kaplan, R. M., & Pierce, J. P. (1997). Cigarette smoking predicts development of depressive symptoms among U.S. adolescents. *Annals of Behavioral Medicine*, 19(1), 42–50. doi: 10.1007/BF02883426.
- Clark, C., Haines, M. M., Head, J., Klineberg, E., Arephin, M., Viner, R., ... Stansfeld, S. A. (2007). Psychological symptoms and physical health and health behaviours in adolescents: A prospective 2-year study in East London. Addiction, 102(1), 126–135. doi: 10.1111/j.1360-0443.2006.01621.x.
- Duncan, B., & Rees, D. I. (2005). Effect of smoking on depressive symptomatology: A reexamination of data from the national longitudinal study of adolescent health. *American Journal of Epidemiology*, 162(5), 461–470. doi: 10.1093/aje/kwi219.
- Eitle, D., & Eitle, T. M. (2018). Obesity, overweightness, and depressive symptomology among American Indian youth. *Journal of Racial and Ethnic Health Disparities*, 5(6), 1305–1314. doi: 10.1007/s40615-018-0479-9.
- Farr, O. M., Ko, B.-J., Joung, K. E., Zaichenko, L., Usher, N., Tsoukas, M., ... Mantzoros, C. S. (2015). Posttraumatic stress disorder, alone or additively with early life adversity, is associated with obesity and cardiometabolic risk. Nutrition, Metabolism and Cardiovascular Diseases, 25(5), 479–488. doi: 10.1016/j.numecd.2015.01.007.
- Frisco, M. L., Houle, J. N., & Lippert, A. M. (2013). Weight change and depression among US young women during the transition to adulthood. *American Journal of Epidemiology*, 178(1), 22–30. doi: 10.1093/aje/kws462.
- Gage, S. H., Hickman, M., Heron, J., Munafò, M. R., Lewis, G., Macleod, J., & Zammit, S. (2015). Associations of Cannabis and cigarette use with depression and anxiety at age 18: Findings from the Avon longitudinal study of parents and children. *PLoS One*, 10(4), e0122896. doi: 10.1371/journal.pone.0122896.

- Gardner, M. J., Thomas, H. J., & Erskine, H. E. (2019). The association between five forms of child maltreatment and depressive and anxiety disorders: A systematic review and meta-analysis. *Child Abuse & Neglect*, 96, 104082. doi: 10.1016/j.chiabu.2019.104082.
- Gomes, A. P., Soares, A. L. G., Menezes, A. M. B., Assunção, M. C., Wehrmeister, F. C., Howe, L. D., & Gonçalves, H. (2019). Adiposity, depression and anxiety: Interrelationship and possible mediators. *Revista de Saúde Pública*, 53, 103. doi: 10.11606/S1518-8787.2019053001119.
- Goodman, E., & Capitman, J. (2000). Depressive symptoms and cigarette smoking among teens. *Pediatrics*, 106(4), 748–755. doi: 10.1542/peds.106.4.748.
- Goodman, E., & Whitaker, R. C. (2002). A prospective study of the role of depression in the development and persistence of adolescent obesity. *Pediatrics*, 110(3), 497–504.
- Hammerton, G., Harold, G., Thapar, A., & Thapar, A. (2013). Depression and blood pressure in high-risk children and adolescents: An investigation using two longitudinal cohorts. *BMJ Open*, 3(9), e003206. doi: 10.1136/ bmjopen-2013-003206.
- Hammerton, G., Thapar, A., & Thapar, A. K. (2014). Association between obesity and depressive disorder in adolescents at high risk for depression. *International Journal of Obesity*, 38(4), 513–519. doi: 10.1038/ijo.2013.133.
- Han, C., Liu, Y., Gong, X., Ye, X., & Zhou, J. (2019). Relationship between second-hand smoke exposure and depressive symptoms: A systematic review and dose-response meta-analysis. *International Journal of Environmental Research and Public Health*, 16(8), 1356. doi: 10.3390/ijerph16081356.
- Hare, D. L., Toukhsati, S. R., Johansson, P., & Jaarsma, T. (2014). Depression and cardiovascular disease: A clinical review. *European Heart Journal*, 35 (21), 1365–1372. doi: 10.1093/eurheartj/eht462.
- Hasin, D. S., Sarvet, A. L., Meyers, J. L., Saha, T. D., Ruan, W. J., Stohl, M., & Grant, B. F. (2018). Epidemiology of adult DSM-5 major depressive disorder and its specifiers in the United States. *JAMA Psychiatry*, 75(4), 336. doi: 10.1001/jamapsychiatry.2017.4602.
- Heredia, F. P. D., Gómez-Martínez, S., & Marcos, A. (2012). Obesity, inflammation and the immune system. *Proceedings of the Nutrition Society*, 71(2), 332–338. doi: 10.1017/S0029665112000092.
- Herrmann-Lingen, C., Meyer, T., Bosbach, A., Chavanon, M.-L., Hassoun, L., Edelmann, F., & Wachter, R. (2018). Cross-sectional and longitudinal associations of systolic blood pressure with quality of life and depressive mood in older adults with cardiovascular risk factors: Results from the observational DIAST-CHF study. *Psychosomatic Medicine*, 80(5), 468–474. doi: 10.1097/PSY.000000000000000591.
- Higgins, J., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M., & Welch, V. (2021). Cochrane Handbook for Systematic Reviews of Interventions version 6.2 (updated February 2021). Retrieved May 13, 2021, from Cochrane website: https://www.training.cochrane.org/handbook.
- Hildrum, B., Mykletun, A., Stordal, E., Bjelland, I., Dahl, A. A., & Holmen, J. (2007). Association of low blood pressure with anxiety and depression: The Nord-Trøndelag Health Study. *Journal of Epidemiology and Community Health*, 61(1), 53–58. doi: 10.1136/jech.2005.044966.
- Hiles, S. A., Baker, A. L., de Malmanche, T., McEvoy, M., Boyle, M., & Attia, J. (2015). The role of inflammatory markers in explaining the association between depression and cardiovascular hospitalisations. *Journal of Behavioral Medicine*, 38(4), 609–619. doi: 10.1007/s10865-015-9637-2.
- Hoare, E., Skouteris, H., Fuller-Tyszkiewicz, M., Millar, L., & Allender, S. (2014). Associations between obesogenic risk factors and depression among adolescents: A systematic review. *Obesity Reviews*, 15(1), 40–51. doi: 10.1111/obr.12069.
- Holtzheimer, P. E., & Mayberg, H. S. (2011). Stuck in a rut: Rethinking depression and its treatment. *Trends in Neurosciences*, 34(1), 1–9. doi: 10.1016/j.tins.2010.10.004.
- Huang, Y., Su, Y., Jiang, Y., & Zhu, M. (2020). Sex differences in the associations between blood pressure and anxiety and depression scores in a middle-aged and elderly population: The Irish longitudinal study on ageing (TILDA). *Journal of Affective Disorders*, 274, 118–125. doi: 10.1016/j.jad.2020.05.133.
- Incollingo Rodriguez, A. C., Epel, E. S., White, M. L., Standen, E. C., Seckl, J. R., & Tomiyama, A. J. (2015). Hypothalamic-pituitary-adrenal axis dysregulation

- and cortisol activity in obesity: A systematic review. *Psychoneuroendocrinology*, 62, 301–318. doi: 10.1016/j.psyneuen.2015.08.014.
- Inouye, M., Abraham, G., Nelson, C. P., Wood, A. M., Sweeting, M. J., Dudbridge, F., ... Samani, N. J. (2018). Genomic risk prediction of coronary artery disease in 480000 adults. *Journal of the American College of Cardiology*, 72(16), 1883–1893. doi: 10.1016/j.jacc.2018.07.079.
- Juruena, M. F., Bocharova, M., Agustini, B., & Young, A. H. (2018). Atypical depression and non-atypical depression: Is HPA axis function a biomarker? A systematic review. *Journal of Affective Disorders*, 233, 45–67. doi: 10.1016/j.jad.2017.09.052.
- Kahn, B. B., & Flier, J. S. (2000). Obesity and insulin resistance. The Journal of Clinical Investigation, 106(4), 473–481. doi: 10.1172/JCI10842.
- Kendler, K. S., Gardner, C. O., Fiske, A., & Gatz, M. (2009). Major depression and coronary artery disease in the Swedish Twin Registry. Archives of General Psychiatry, 66(8), 857–863. doi: 10.1001/archgenpsychiatry.2009.94.
- Kessler, R. C. (2012). The costs of depression. The Psychiatric Clinics of North America, 35(1), 1–14. doi: 10.1016/j.psc.2011.11.005.
- Kessler, R. C., McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., ... Williams, D. R. (2010). Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *The British Journal of Psychiatry*, 197(5), 378–385. doi: 10.1192/bjp.bp.110.080499.
- Khandaker, G. M., Zuber, V., Rees, J. M. B., Carvalho, L., Mason, A. M., Foley, C. N., ... Burgess, S. (2019). Shared mechanisms between coronary heart disease and depression: Findings from a large UK general population-based cohort. *Molecular Psychiatry*, 25, 1477–1486. doi: 10.1038/s41380-019-0395-3..
- Lamers, F., Milaneschi, Y., Vinkers, C. H., Schoevers, R. A., Giltay, E. J., & Penninx, B. W. J. H. (2020). Depression profilers and immuno-metabolic dysregulation: Longitudinal results from the NESDA study. *Brain, Behavior, and Immunity*, 88, 174–183. doi: 10.1016/j.bbi.2020.04.002.
- Lamers, F., Vogelzangs, N., Merikangas, K. R., de Jonge, P., Beekman, A. T. F., & Penninx, B. W. J. H. (2013). Evidence for a differential role of HPA-axis function, inflammation and metabolic syndrome in melancholic versus atypical depression. *Molecular Psychiatry*, 18(6), 692–699. doi: 10.1038/ mp.2012.144.
- Lippi, G., Montagnana, M., Favaloro, E. J., & Franchini, M. (2009). Mental depression and cardiovascular disease: A multifaceted, bidirectional association. Seminars in Thrombosis and Hemostasis, 35(3), 325–336. doi: 10.1055/s-0029-1222611.
- Luger, T. M., Suls, J., & Vander Weg, M. W. (2014). How robust is the association between smoking and depression in adults? A meta-analysis using linear mixed-effects models. *Addictive Behaviors*, 39(10), 1418–1429. doi: 10.1016/j.addbeh.2014.05.011.
- Luppino, F. S., Wit, L. M. D., Bouvy, P. F., Stijnen, T., Cuijpers, P., Penninx, B. W. J. H., & Zitman, F. G. (2010). Overweight, obesity, and depression: A systematic review and meta-analysis of longitudinal studies. *Archives of General Psychiatry*, 67(3), 220–229. doi: 10.1001/archgenpsychiatry.2010.2.
- Mannan, M., Mamun, A., Doi, S., & Clavarino, A. (2016). Prospective associations between depression and obesity for adolescent males and females: A systematic review and meta-analysis of longitudinal studies. *PLoS One*, 11(6), e0157240. doi: 10.1371/journal.pone.0157240.
- Marmorstein, N. R., Iacono, W. G., & Legrand, L. (2014). Obesity and depression in adolescence and beyond: Reciprocal risks. *International Journal of Obesity*, 38(7), 906–911. doi: 10.1038/ijo.2014.19.
- Monshouwer, K., Smit, F., Ruiter, M., Ormel, H., Verhulst, F., Vollebergh, W., & Oldehinkel, T. (2012). Identifying target groups for the prevention of depression in early adolescence: The TRAILS study. *Journal of Affective Disorders*, 138(3), 287–294. doi: 10.1016/j.jad.2012.01.026.
- Mühlig, Y., Antel, J., Föcker, M., & Hebebrand, J. (2016). Are bidirectional associations of obesity and depression already apparent in childhood and adolescence as based on high-quality studies? A systematic review. *Obesity Reviews*, 17(3), 235–249. doi: 10.1111/obr.12357.
- Munafò, M. R., Hitsman, B., Rende, R., Metcalfe, C., & Niaura, R. (2008). Effects of progression to cigarette smoking on depressed mood in adolescents: Evidence from the national longitudinal study of adolescent health. Addiction, 103(1), 162–171. doi: 10.1111/j.1360-0443.2007.02052.x.
- Niarchou, M., Zammit, S., & Lewis, G. (2015). The avon longitudinal study of parents and children (ALSPAC) birth cohort as a resource for studying

psychopathology in childhood and adolescence: A summary of findings for depression and psychosis. *Social Psychiatry and Psychiatric Epidemiology*, *50* (7), 1017–1027. doi: 10.1007/s00127-015-1072-8.

- Osimo, E. F., Baxter, L. J., Lewis, G., Jones, P. B., & Khandaker, G. M. (2019). Prevalence of low-grade inflammation in depression: A systematic review and meta-analysis of CRP levels. *Psychological Medicine*, 49(12), 1958– 1970. doi: 10.1017/S0033291719001454.
- Perry, B. I., Khandaker, G. M., Marwaha, S., Thompson, A., Zammit, S., Singh, S. P., & Upthegrove, R. (2020). Insulin resistance and obesity, and their association with depression in relatively young people: Findings from a large UK birth cohort. *Psychological Medicine*, 50(4), 556–565. doi: 10.1017/S0033291719000308.
- Piumatti, G. (2018). Motivation, health-related lifestyles and depression among university students: A longitudinal analysis. *Psychiatry Research*, 260, 412–417. doi: 10.1016/j.psychres.2017.12.009.
- Polanczyk, G. V., Salum, G. A., Sugaya, L. S., Caye, A., & Rohde, L. A. (2015). Annual research review: A meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *Journal of Child Psychology* and Psychiatry, and Allied Disciplines, 56(3), 345–365. doi: 10.1111/jcpp.12381.
- Pryor, L., Brendgen, M., Boivin, M., Dubois, L., Japel, C., Falissard, B., ... Côté, S. M. (2016). Overweight during childhood and internalizing symptoms in early adolescence: The mediating role of peer victimization and the desire to be thinner. *Journal of Affective Disorders*, 202, 203–209. doi: 10.1016/j.jad.2016.05.022.
- Raffetti, E., Donato, F., Forsell, Y., & Galanti, M. R. (2019). Longitudinal association between tobacco use and the onset of depressive symptoms among Swedish adolescents: The Kupol cohort study. *European Child & Adolescent Psychiatry*, 28(5), 695–704. doi: 10.1007/s00787-018-1237-6.
- Ranjit, A., Buchwald, J., Latvala, A., Heikkilä, K., Tuulio-Henriksson, A., Rose, R. J., ... Korhonen, T. (2019a). Predictive association of smoking with depressive symptoms: A longitudinal study of adolescent twins. Prevention Science, 20(7), 1021–1030. doi: 10.1007/s11121-019-01020-6.
- Ranjit, A., Korhonen, T., Buchwald, J., Heikkilä, K., Tuulio-Henriksson, A., Rose, R. J., ... Latvala, A. (2019b). Testing the reciprocal association between smoking and depressive symptoms from adolescence to adulthood: A longitudinal twin study. *Drug and Alcohol Dependence*, 200, 64–70. doi: 10.1016/j.drugalcdep.2019.03.012.
- Rhew, I. C., Richardson, L. P., Lymp, J., McTiernan, A., McCauley, E., & Stoep, A. V. (2008). Measurement matters in the association between early adolescent depressive symptoms and body mass index. *General Hospital Psychiatry*, 30(5), 458–466. doi: 10.1016/j.genhosppsych.2008.06.008.
- Roberts, R. E., & Duong, H. T. (2013). Obese youths are not more likely to become depressed, but depressed youths are more likely to become obese. *Psychological Medicine*, 43(10), 2143–2151. doi: 10.1017/S0033291712002991.
- Roth, G. A., Abate, D., Abate, K. H., Abay, S. M., Abbafati, C., Abbasi, N., ... Murray, C. J. L. (2018). Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: A systematic analysis for the Global Burden of Disease Study 2017. The Lancet, 392(10159), 1736–1788. doi: 10.1016/S0140-6736(18)32203-7.
- Rubio, D. M., Kraemer, K. L., Farrell, M. H., & Day, N. L. (2008). Factors associated with alcohol use, depression, and their co-occurrence during pregnancy. *Alcoholism: Clinical and Experimental Research*, 32(9), 1543–1551. doi: 10.1111/j.1530-0277.2008.00705.x.
- Schwarzer, G. (2007). Meta: An R package for meta-analysis. R News, 7, 40–45.

- Shin, J. Y., Suls, J., & Martin, R. (2008). Are cholesterol and depression inversely related? A meta-analysis of the association between two cardiac risk factors. *Annals of Behavioral Medicine*, 36(1), 33–43. doi: 10.1007/ s12160-008-9045-8.
- Smolderen, K. G., Spertus, J. A., Gosch, K., Dreyer, R. P., D'Onofrio, G., Lichtman, J. H., ... Krumholz, H. M. (2017). Depression treatment and health Status outcomes in young patients with acute myocardial infarction: Insights from the VIRGO study (variation in recovery: Role of gender on outcomes of young AMI patients). Circulation, 135(18), 1762–1764. doi: 10.1161/CIRCULATIONAHA.116.027042.
- Speed, M. S., Jefsen, O. H., Børglum, A. D., Speed, D., & Østergaard, S. D. (2019). Investigating the association between body fat and depression via mendelian randomization. *Translational Psychiatry*, 9(1), 1–9. doi: 10.1038/ s41398-019-0516-4.
- Stang, A. (2010). Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. European Journal of Epidemiology, 25(9), 603–605.
- Sutaria, S., Devakumar, D., Yasuda, S. S., Das, S., & Saxena, S. (2019). Is obesity associated with depression in children? Systematic review and meta-analysis. Archives of Disease in Childhood, 104(1), 64–74. doi: 10.1136/archdischild-2017-314608.
- Taylor, A. E., Fluharty, M. E., Bjørngaard, J. H., Gabrielsen, M. E., Skorpen, F., Marioni, R. E., ... Munafò, M. R. (2014). Investigating the possible causal association of smoking with depression and anxiety using Mendelian randomisation meta-analysis: The CARTA consortium. BMJ Open, 4(10), e006141. doi: 10.1136/bmjopen-2014-006141.
- Van der Kooy, K., van Hout, H., Marwijk, H., Marten, H., Stehouwer, C., & Beekman, A. (2007). Depression and the risk for cardiovascular diseases: Systematic review and meta analysis. *International Journal of Geriatric Psychiatry*, 22(7), 613–626. doi: 10.1002/gps.1723.
- Wang, H., Leung, G. M., & Schooling, C. M. (2014). Life course adiposity and adolescent depressive symptoms among Hong Kong adolescents. *Journal of Adolescent Health*, 55(3), 408–414. doi: 10.1016/j.jadohealth.2014.03.009.
- Wei, Y.-G., Cai, D.-B., Liu, J., Liu, R.-X., Wang, S.-B., Tang, Y.-Q., ... Wang, F. (2020). Cholesterol and triglyceride levels in first-episode patients with major depressive disorder: A meta-analysis of case-control studies. *Journal of Affective Disorders*, 266, 465–472. doi: 10.1016/j.jad.2020.01.114.
- WHO Study Group of Young People (1986). Young people's health a challenge for society: Report of a WHO study group on young people and 'health for all by the year 2000'. Geneva: World Health Organization. Retrieved from https://apps.who.int/iris/handle/10665/41720.
- Wilson, P. W. F., Castelli, W. P., & Kannel, W. B. (1987). Coronary risk prediction in adults (The Framingham Heart Study). The American Journal of Cardiology, 59(14), G91–G94. doi: 10.1016/0002-9149(87)90165-2.
- Wilson, P. W., D'Agostino, R. B., Levy, D., Belanger, A. M., Silbershatz, H., & Kannel, W. B. (1998). Prediction of coronary heart disease using risk factor categories. *Circulation*, 97(18), 1837–1847. doi: 10.1161/01.cir.97.18.1837.
- Wootton, R. E., Richmond, R. C., Stuijfzand, B. G., Lawn, R. B., Sallis, H. M., Taylor, G. M. J., ... Munafò, M. R. (2020). Evidence for causal effects of lifetime smoking on risk for depression and schizophrenia: A Mendelian randomisation study. *Psychological Medicine*, 50(14), 2435–2443. doi: 10.1017/ S0033291719002678.
- Zhang, X. C., Woud, M. L., Becker, E. S., & Margraf, J. (2018). Do health-related factors predict major depression? A longitudinal epidemiologic study. Clinical Psychology & Psychotherapy, 25(3), 378–387. doi: 10.1002/cpp.2171.