

CMS SUSY Searches at 13 TeV

LHC SEMINAR
9TH FEBRUARY 2016

PABLO MARTINEZ RUIZ DEL ARBOL

ON BEHALF OF THE CMS COLLABORATION



ETH Institute for
Particle Physics





Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The Standard Model: not quite everything

- The Standard Model of Particles even if extremely successful cannot explain everything
- The hierarchy problem → stability of the Higgs mass under quantum corrections
- The problem of dark matter → gravitational effects observed but not a “candidate”
- Impossibility to achieve Grand Unification → not converging couplings at Λ_{GUT}

A diagram showing two circular loops representing the annihilation of top quarks (t) and anti-top quarks (\bar{t}). The loops are labeled "H" at their vertices. An arrow points from this diagram to the following equation:


$$\Delta m_H^2 = \frac{-(|\lambda_f|^2)}{(8\pi)} [\Lambda_{UV}^2 + \dots]$$



Supersymmetry: elegant and gorgeous* Φ

ETH Institute for
Particle Physics


- › Supersymmetry imposes a new fundamental symmetry between fermions and bosons
- › The new super-partners can solve the aforementioned problems
 - › Δm_H compensated by boson loops → if gluino, stop, higgsino masses are at the \sim TeV scale
 - › Provides dark matter candidates → linear combinations of electroweak and Higgs SUSY partners
 - › Leads to unification → running couplings coincident at Λ_{GUT}



* Quoting Marcela Carena (ICHEP14, Plenary Talk)

Supersymmetry: is anybody there?


- No evidence of SUSY particles nor SM deviation have been found at the LHC
- Rare decays with high sensitivity to new particles through quantum corrections measured
 - $B_s \rightarrow \mu\mu$ in good agreement with the Standard Model prediction → ???
- After Run I we know there is a new 125 GeV scalar particle exists: the Higgs boson
 - This is important since it requires $\Delta m_H \sim 34$ GeV $m_{\text{Higgs}}^2 \approx m_Z^2 \left[1 + \frac{3m_t^2}{2\pi^2 m_Z^2} \log(m_{\text{stop}}/m_{\text{top}}) \right]$
 - This implies stops to have maximal mixing or to be very heavy → some tension with naturalness



Still in a valid range


Supersymmetry: Hiding from us?

- Very ambitious SUSY search programs have been carried out at the LHC during Run 1
- However SUSY might be hard to find ... it might be hiding
 - SUSY particles too heavy for 8 TeV? Cross sections very small at LHC, hardly produced?
 - Very **compressed spectra** leading to very difficult experimental signatures
 - **Stealth SUSY**: so long decays that spectrum of produced particles is very soft
 - **R-Parity Violation**: again complicated signatures with no momentum imbalance
- The 13 TeV data will provide insight on these searches → **we're back with a better tool**



SUSY experimental signatures

- › At the LHC we are looking for pair production of SUSY particles in the hard interaction
- › SUSY particles assumed to be heavy → long decay chains expected including SM particles
 - › As a result typical SUSY signatures contain usually many jets in the event
- › Also if **R-parity** is conserved the lightest neutralino (LSP) is stable and escapes detection
 - › Leaving a large amount of **Missing Transverse Energy** in the event



A “typical” SUSY event

High Jet multiplicity


High MET $\rightarrow |-\sum \vec{p}_T(\text{particle})|$

High HT $\rightarrow \sum |\vec{p}_T(\text{jets})|$

High MHT $\rightarrow |\sum \vec{p}_T(\text{jets})|$

SUSY Interpretations: Simplified Models




- Most of SUSY interpretations are performed using Simplified Models of Supersymmetry*
- This approach puts more emphasis on the experimental signature leaving aside the model details
- Hard interaction producing two SUSY particles (fully decoupled from other particles)
- A single decay chain is implemented producing a given and fixed topology
- However the masses of the SUSY particles are scanned usually in 2 dimensions
- Upper limits are set on the **cross-section x branching ratio** of the given process



* Interpretations in full models are also provided: mSUGRA, pMSSM, etc



SUSY searches at the LHC: Run 1

- A huge amount of searches seeking all kind of different topologies
- **Gluino mass** has been probed up to ~ 1300 GeV in different topologies



SUSY searches at the LHC: Run 1


- stop masses have been also probed up to ~ 750 GeV




- But also upper limits on light squarks, s-bottoms, neutralinos/charginos, s-leptons, RPV SUSY

CMS, a wonderful detector

The CMS detector at the LHC




The CMS detector at the LHC



LHC/CMS performance in 2015



CMS Integrated Luminosity, pp, 2015, $\sqrt{s} = 13 \text{ TeV}$

Data included from 2015-06-03 08:41 to 2015-11-03 06:25 UTC





Triggers at 13 TeV

- › Upgraded L1 trigger logic including better pile-up rejection and improved algorithms
- › HLT reconstruction also improved with a simplified version of the iterative tracking
- › Efficiency above **95%** for most trigger paths including MET, jets, single/double lepton
- › All these improvements allowed to keep trigger thresholds close to 8 TeV in most cases




Physics objects at 13 TeV: jets and sums

- Very good performance of jet reco and jet sum objects (MET, MHT) both at 50 and 25 ns
- Full chain of Jet Energy Corrections (L1L2L3Residual) extracted from collected data
- Detailed studies on MET resolution and response, dependency with pile-up, etc, etc
- Also new and improved b-tagging discriminators deployed for the 13 TeV data taken






Physics objects at 13 TeV: leptons-photons

- New and improved identification algorithms for muons, electrons and photons
- Combining cut-based and multivariate discriminators to improve ID efficiency
- Extended use of mini-isolation → isolation cone ΔR dependent on pT of the lepton




Nice performance for very boosted systems



Excellent physics performance


- › Di-muon resonances decaying into pairs of muons
- › Excellent muon reconstruction and resolution over a wide range of momentum



SUSY searches at 13 TeV

SUSY at 13 TeV in CMS

- › In the year 2015 a total of 2.2 fb^{-1} were collected by CMS at a center of mass of 13 TeV
- › Two competing effects: cross section increase due to CME and luminosity ratio
- › What can we expect at 13 TeV compared to the 8 TeV in terms of SUSY production?




Increased sensitivity is expected for gluino-gluino production and slightly for stop-stop production

https://twiki.cern.ch/twiki/bin/view/LHCPhysics/SUSYCrossSections#SUSY_Cross_Sections_using_8_TeV

SUSY searches at 13 TeV: overview


- › Full set of analysis covering different final states and using 2.2 fb^{-1} collected at 13 TeV
- › Most of them targeting the direct production of **gluino pairs** decaying into several topologies
- › All of them including a large variety of different signal regions to increase sensitivity





All-hadronic searches

Multijet + MET search

- Inclusive search binned in HT, MHT, #jets and #b-jets targeting gluino production





- Most important backgrounds coming from QCD events with instrumental MET
- Events with real MET like $Z(vv)+\text{jets}$, taus, $W+\text{jets}$ and top-pair production (with lost lepton)



Background prediction


- QCD estimated from a low $\Delta\phi_{\min}$ control region (min angle between jets and MHT vector)
 - In QCD events with instrumental MET usually MHT is aligned with one of the jets
 - This variable will be used by many other all-hadronic searches as it will be shown later
- $Z(v\bar{v}) + \text{jets} \rightarrow$ estimated from a $\gamma + \text{jets}$ sample subtracting the photon from the event
 - Correction factors extracted from data and MC to account for differences
- W + jets and top pair production with lost lepton is predicted from the lost lepton probability
 - Which is measured in a 1 lepton control sample and then applied to the signal region
- Hadronic tau leptons are finally estimated from $W \rightarrow \mu + \text{neutrino}$ control region

SUS-15-002




Multijet + MET search: results

- Good agreement seen between observation and Standard Model background predictions




Upper limits on Simplified Models

- Upper limits are set on the T1qqqq SMS pushing the limit on the gluino mass about ~ 100 GeV





8 TeV result





Upper limits on Simplified Models

- New Upper limit set on the T1bbbb SMS with this analysis reaching 1400 GeV in gluino mass




Upper limits on Simplified Models

- Upper limits are set on the T1tttt SMS pushing the limit on the gluino mass about ~ 300 GeV




All-hadronic analysis MT2

- › Inclusive search using the MT2 variable and targeting mainly gluino production




- › Leptons and photons are vetoed and signal regions are asked to contain more than 1 jet
- › Signal regions are requested to have $\Delta\Phi_{\min} > 0.3$ and then binned in HT, # of jets and b-jets and MT2
- › A new category is also included with only 1 jet and very low HT ($HT > 200$)
 - › This new category targets to improve the sensitivity of the analysis to models like T2qq or T2tt
 - › Particularly in those regions with very compressed spectra



SUS-15-003

MT2 and $\Delta\Phi_{\min}$

- › MT2 is a generalization of the transverse mass for a system with 2 invisible particles
 - › This variable has a cut-off at the mass of the produced particles
 - › In practice jets are merged into 2 “pseudo-jets” to apply the definition
 - › SM processes and more in particular multijet QCD events have very low values of MT2





$$M_{T2} = \min_{\vec{p}_T^{X(1)} + \vec{p}_T^{X(2)} = \vec{p}_T^{\text{miss}}} [\max(M_T^{(1)}, M_T^{(2)})]$$

SUS-15-003

- › The analysis also uses $\Delta\Phi_{\min}$ which is the minimum angle between a jet and the MET vector
 - › QCD events contain instrumental MET originating from jet mismeasurements
 - › Events where the MET is pointing in the same direction as a jet are likely to be QCD

Background prediction

- › The main backgrounds are QCD, Z($\nu\nu$)+jets, and top-pair production or W+jets with lost lepton
- › QCD is estimated by extrapolating the ratio $\#\Delta\Phi_{\min} > 0.3 / \#\Delta\Phi_{\min} < 0.3$ at low MT2 to high MT2
 - › The extrapolated ratio is multiplied to the yields in the QCD enriched region ($\Delta\Phi_{\min} < 0.3$) at high MT2




SUS-15-003

- › The Z($\nu\nu$) + Jets contribution is estimated from a γ +Jets control region and MC transfer factors
- › Ttbar and W+jets also estimated from a control region with 1 lepton and MC transfer factors

All-hadronic analysis MT2: Results

- Observations in the different signal regions are found to be compatible with SM predictions



Entries



SUS-15-003



Upper limits on Simplified Models

- Upper limits are set on the T1qqqq SMS pushing the limit on the gluino mass about ~ 300 GeV





Upper limits on Simplified Models

- Upper limits are set on the T1bbbb SMS pushing the limit on the gluino mass about ~ 300 GeV





Upper limits on Simplified Models

- Upper limits are set on the T1tttt SMS pushing the limit on the gluino mass about ~ 300 GeV



All-hadronic analysis using α -T

- › Pure hadronic search using the α -T and $\Delta\phi_{\min}^*$ variables to strongly reduce the amount of QCD
- › The analysis targets SUSY models with gluinos decaying into light quarks/bottoms and LSP





SUS-15-005

- › The analysis defines 3 different categories of signal regions attending to the two leading jets
 - › **Symmetric jet selection:** 2 leading jets with $\text{Pt} > 100 \text{ GeV}$
 - › **Asymmetric jet selection:** leading jet with $\text{Pt} > 100 \text{ GeV}$ and sub-leading $\text{Pt} > 40 \text{ GeV}$
 - › **Mono-jet category:** leading jet with $\text{Pt} > 100 \text{ GeV}$ and sub-leading $\text{Pt} < 40 \text{ GeV}$ (compressed spectra)
- › Additional binning according to HT, number of jets and number of b-jets
- › Varying α -T cuts for different HT bins and additional $\Delta\phi_{\min}^*$ and MHT/MET cuts

α -T and $\Delta\phi^*$ _{min} variables: QCD killers

- The α -T variable is defined as the transverse momentum of the second jet over mT
 - In events with multi-jet topology, jets are merged into a leading and sub-leading jet
 - The α -T variable gets values < 0.5 for QCD events (with a back-to-back jet topology)
- $\Delta\phi^*$ _{min} is defined as the min. angle between a jet and the MHT vector formed by the rest
 - Small values of this angle indicate a deficit of energy in the direction of the jets


SUS-15-005



Background prediction



- SM background are estimated from **control regions and Monte Carlo transfer factors**
 - Predictions are validated using additional control regions (validation regions)
- QCD from a QCD enriched control region where MHT/MET is inverted
- Other backgrounds with real MET:
 - W+Jets, DY+Jets ($Z \rightarrow \nu\nu + \text{Jets}$), top pair production
 - Extracted from Control Regions with:
 - $\gamma + \text{Jets}$, $\mu + \text{Jets}$ and $\mu\mu + \text{Jets}$
- Systematic studies performed on validation regions
- Yields in control/signal regions and transfer factors fitted
 - To get the final background prediction
 - Good agreement between observation/prediction

SUS-15-005





Upper limits on Simplified Models

- Upper limits are set on the T1qqqq SMS pushing the limit on the gluino mass about ~ 400 GeV




Upper limits on Simplified Models

- Upper limits are set on the T1bbbb SMS pushing the limit on the gluino mass about ~ 400 GeV



Razor analysis

- › Razor analysis is a combined 0 lepton and 1 lepton search using the **razor variables**
- › The analysis targets also the T1qqqq and T1bbbb SMS and the T1tttt (also in the 1 lepton)



- › Therefore three different categories with different base-line selections, all binned in #b-jets
- › **1 muon + multijet** → one muon with $\text{pt} > 25 \text{ GeV}$ and at least 4 jets with $\text{pt} > 40 \text{ GeV}$
- › **1 electron + multijet** → one electron with $\text{pt} > 25 \text{ GeV}$ and at least 4 jets with $\text{pt} > 40 \text{ GeV}$
- › **Multijet** → at least 4 jets with $\text{pt} > 40 \text{ GeV}$ and ≥ 2 jets with 80 GeV
- › In order to reduce the SM background a cut on the transverse mass of the lepton is performed
- › In the multijet case, jets are grouped in 2 megajets and cut $\Delta\Phi < 2.8$ (angle between megajets)
 - › Strongly reduces the amount of background

SUS-15-004



The Razor variables

- A shape analysis performed in the plane of two kinematic variables denoted MR and R^2
- These variables correspond to the mass and transverse energy flow of pair-produced particles
- In a decay: $\tilde{q} \rightarrow q + \text{LSP}$ MR peaks at $(M(\tilde{q}) - M(q))/M(\tilde{q})$ and R^2 is related to the MET

$$M_R \equiv \sqrt{(P_{j_1} + P_{j_2})^2 - (p_z^{j_1} + p_z^{j_2})^2} \quad M_T^R \equiv \sqrt{\frac{E_T^{\text{miss}}(p_T^{j_1} + p_T^{j_2}) - \vec{p}_T^{\text{miss}} \cdot (\vec{p}_T^{j_1} + \vec{p}_T^{j_2})}{2}} \quad R^2 \equiv \left(\frac{M_T^R}{M_R}\right)^2$$



- Many control (sidebands) and signal regions are defined in the MR and R^2 plane

SUS-15-004



Background estimation

- › Main backgrounds are QCD (instrumental MET) , top-pair production, W+jets and DY+jets
- › 2D analytical functions modeling MR/R2 are validated in control regions for every background
- › A total final fit is performed in the sidebands and extrapolated to the signal regions




SUS-15-004


Razor analysis: results

- Very good agreement found between the observation and the prediction in all signal regions

Multijet with 0 b-tag





Electron + multijet with 0 b-tag



SUS-15-004



Upper limits on Simplified Models

- First upper limits on the T1qqqq SMS reaching gluino masses up to ~ 1300 GeV





Upper limits on Simplified Models

- Upper limits are set on the T1bbbb SMS pushing the limit on the gluino mass about ~ 300 GeV




Upper limits on Simplified Models


- Upper limits are set on the T1tttt SMS pushing the limit on the gluino mass about ~ 400 GeV



T1qqqq topology (all-hadronic): summary



T1bbbb topology (all-hadronic): summary




Search with one lepton


1 Lepton sum of FatJet masses

- Search for SUSY with 1 lepton, jets and MET in a boosted regime and using two variables:
 - M_j = Sum of the **Fat-Jet** masses in the event
 - MT = transverse mass between the lepton and the MET vector
- Targeting gluino production decaying to boosted tops and LSP through off-shell stop

SUS-15-007




Fat-jet: jet with $R=1.2$ instead of the regular 0.4



- Baseline selection asking for $HT > 500$ GeV, $MET > 200$ GeV, $\#jets \geq 7$, $\#b\text{-jets} \geq 2$


FatJet masses: heavy objects

- › Large M_J is characteristic of decays of heavy objects producing very boosted particles (jets)
- › In SM events this can occur mainly because of accidental jet overlap and ISR jets





Background prediction: mT vs. MJ

- Background is mainly dominated by top pair production both with 1 and 2 leptons
- ABCD method implemented in the MJ and mT plane in bins of MET, #jets and #b-jets
- MJ and mT are approximately uncorrelated variables → differences corrected using MC



Upper limits on Simplified Model




- Good agreement is found between observations and background predictions
- Upper limit in the range 1500-1600 GeV for the gluino mass



Searches with two leptons



Same Sign Leptons

- Targeting production of Same Sign leptons in SUSY decay chains with gluinos and squarks



- Binned in regions of HT, MET, $M_{t\bar{t}}^{\min}$ and number of b-jets

SUS-15-008



3 lepton categories




High-High:
both $\text{pt} > 25 \text{ GeV}$

Low-High:
 $\text{pt} > 15 \text{ GeV}$ and $\text{pt} < 25 \text{ GeV}$
 $\text{pt} > 25 \text{ GeV}$

Low-Low:
 $\text{pt} > 15 \text{ GeV}$ and $\text{pt} < 25 \text{ GeV}$

Background prediction



- Standard model produces very few same sign lepton backgrounds:
 - Fake leptons (not necessarily non-prompt-leptons) → tight-to-loose method
 - Charge flipping for electrons → measured in DY Control Region
 - Rare SM backgrounds TTZ, TTW → from MC, WZ from data CR



SUS-15-008

Same Sign Leptons: results



› Good agreement is found between observation and SM prediction in all categories




SUS-15-008

Same Sign Leptons: limits on T1tttt

- Upper limits on gluino masses for the T1tttt model are pushed ~ 200 GeV





T1tttt topology: summary



Opposite-sign leptons: The EdgeZ analysis

- SUSY decay chains can lead to opposite sign same flavor leptons in the final state
- Decays involving an on-shell Z boson will produce an excess on the Z peak invariant mass
 - Search for an excess of events in the tails of the MET distribution (traditional SUSY-like)
- Off-shell Z boson or slepton decays lead to a characteristic “edge” shape
 - Search for this structure in the invariant mass distribution (bump hunting-like)

SUS-15-011





All possible mass endpoints calculated in CMS IN 2006/012, L.Pape, e.g. for a 3-body: $M_{ll}^{\max} = M_X - M_0$

- CMS observed a 2.6 sigma deviation in the off-Z region at 8 TeV
- ATLAS reported 3 sigma in the on-Z region at 8 TeV and 2.2 in the same region at 13 TeV

Opposite-sign leptons at 13 TeV

- Tha analysis has been repeated at 13 TeV only in its cut & count version
- The signal regions for the off-Z search have been kept unchanged with respect to 8 TeV
 - However new b-tagged exclusive regions were defined to explore the b-tag dependency
 - New signal regions covering the holes in the invariant mass have been also added




- The on-Z search extended by adding much higher granularity on jet multiplicity and MET
 - Signal A:** 2-3 jets and HT > 400 with and without b-tagging + binning in MET
 - Signal B:** ≥ 4 jets with and without b-tagging + binning in MET
 - A new region as similar as possible to the **ATLAS region** has been also defined
 - $\text{HT} + \text{pT(lep1)} + \text{pT(lep2)} > 600$, $\text{MET} > 225$ GeV, $\Delta\phi(\text{MET}, \text{jets}) > 0.4$

SUS-15-011


Background prediction & results

- › The main SM backgrounds: top-pair production (real MET) and DY+jets (instrumental MET)
- › Top-pair production → Extracted from $e\mu$ events after correcting for efficiency differences
- › DY + Jets → Measure instrumental MET shape in a $\gamma + \text{jets}$ control sample with a similar MET origin
- › Good agreement between prediction and observation for all signal regions



SUS-15-011



Edge Signal Regions



on-Z Signal Region




A closer look to the edge region

- Good agreement is observed in the classic edge region (also in the region with > 0 b-tags)
- 13 TeV data disfavors the signal hypothesis although it is on the limit of sensitivity**





A closer look to the ATLAS region

- CMS does not observe any excess in the so called “ATLAS” region



Upper limits on Simplified Models

- Upper limits on gluino masses for the T5ZZ model are pushed ~ 100 GeV



SUS-15-011

Summary & outlook

- › CMS collected 2.2 fb^{-1} of data in 2015 with a new center of mass energy of 13 TeV
- › A complete set of SUSY analysis covering different topologies have been deployed by CMS
- › Given the current amount of data most of them targeting mainly gluino production
- › A good agreement between observation and prediction has been observed in all cases
- › Upper limits on gluino mass has been pushed forward about $\sim 300 \text{ GeV}$ (for low LSP mass)
- › The “Edge-Z” analysis has been also repeated to crosscheck the previous results
 - › However 13 TeV results on the Edge region (2.6 sigma at 8 TeV) disfavor the signal hypothesis
 - › CMS does not see any excess in the region where ATLAS saw 3.0 and 2.2 sigma excesses
- › However this was only the first chunk of results with the first 2.2 fb^{-1} at 13 TeV
 - › Many more results will be coming by the time of Moriond
 - › And of course many more results will be coming with the new data of 2016

Stay tuned!

Thank you for the attention!

Questions???