

class14

AUTHOR

lisa chen A17082974

```
metadata <- read.csv("GSE37704_metadata.csv", row.names = 1)
counts <- read.csv("GSE37704_featurecounts.csv", row.names = 1)
```

```
head(counts)
```

	length	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370
ENSG00000186092	918	0	0	0	0	0
ENSG00000279928	718	0	0	0	0	0
ENSG00000279457	1982	23	28	29	29	28
ENSG00000278566	939	0	0	0	0	0
ENSG00000273547	939	0	0	0	0	0
ENSG00000187634	3214	124	123	205	207	212
	SRR493371					
ENSG00000186092	0					
ENSG00000279928	0					
ENSG00000279457	46					
ENSG00000278566	0					
ENSG00000273547	0					
ENSG00000187634	258					

```
head(metadata)
```

	condition
SRR493366	control_sirna
SRR493367	control_sirna
SRR493368	control_sirna
SRR493369	hoxa1_kd
SRR493370	hoxa1_kd
SRR493371	hoxa1_kd

Data exploration

Hmm... remember that we need the countData and colData files to match up so we will need to remove that odd first column in countData namely countData\$length.

Question

```
counts <- counts[, !names(counts) == "length"]
head(counts)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000186092	0	0	0	0	0	0
ENSG00000279928	0	0	0	0	0	0
ENSG00000279457	23	28	29	29	28	46
ENSG00000278566	0	0	0	0	0	0
ENSG00000273547	0	0	0	0	0	0
ENSG00000187634	124	123	205	207	212	258

Check:

```
all(colnames(counts) == metadata$id)
```

```
[1] TRUE
```

We need to remove all the zero count genes.

```
head(counts, 3)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000186092	0	0	0	0	0	0
ENSG00000279928	0	0	0	0	0	0
ENSG00000279457	23	28	29	29	28	46

Question:

To identify these zero count genes we can sum across the rows and check if the sum is more than zero.

```
to.keep.inds <- rowSums(counts) > 0
nonzero.counts <- counts[to.keep.inds,]
head(nonzero.counts)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000279457	23	28	29	29	28	46
ENSG00000187634	124	123	205	207	212	258

ENSG00000188976	1637	1831	2383	1226	1326	1504
ENSG00000187961	120	153	180	236	255	357
ENSG00000187583	24	48	65	44	48	64
ENSG00000187642	4	9	16	14	16	16

DESeq Setup and analysis

```
library(DESeq2)
```

```
dds <- DESeqDataSetFromMatrix(countData = nonzero.counts,  
                              colData = metadata,  
                              design=~condition)
```

Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in design formula are characters, converting to factors

```
dds <- DESeq(dds)
```

estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing

```
dds <- DESeq(dds)
```

using pre-existing size factors

estimating dispersions

found already estimated dispersions, replacing these

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing

Result extraction

```
res <- results(dds, contrast=c("condition", "hoxa1_kd", "control_sirna"))
```

Question

```
summary(res)
```

out of 15975 with nonzero total read count

adjusted p-value < 0.1

LFC > 0 (up) : 4349, 27%

LFC < 0 (down) : 4396, 28%

outliers [1] : 0, 0%

low counts [2] : 1237, 7.7%


(mean count < 0)

[1] see 'cooksCutoff' argument of ?results

[2] see 'independentFiltering' argument of ?results

Volcano plot

```
plot( res$log2FoldChange, -log(res$padj) )
```




Q. Improve this plot by completing the below code, which adds color and axis labels

```
# Make a color vector for all genes
mycols <- rep("gray", nrow(res) )

# Color red the genes with absolute fold change above 2
mycols[ abs(res$log2FoldChange) > 2 ] <- "red"

# Color blue those with adjusted p-value less than 0.01
# and absolute fold change more than 2
inds <- (res$padj < .01 ) & (abs(res$log2FoldChange) > 2 )
mycols[ inds ] <- "blue"

plot( res$log2FoldChange, -log(res$padj), col=mycols, xlab="Log2(FoldChange)", ylab="
```



Adding gene annotation

Question

```
library("AnnotationDbi")
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```

[1] "ACCNUM"	"ALIAS"	"ENSEMBL"	"ENSEMBLPROT"	"ENSEMBLTRANS"
[6] "ENTREZID"	"ENZYME"	"EVIDENCE"	"EVIDENCEALL"	"GENENAME"
[11] "GENETYPE"	"GO"	"GOALL"	"IPI"	"MAP"
[16] "OMIM"	"ONTOLOGY"	"ONTOLOGYALL"	"PATH"	"PFAM"

```
[21] "PMID"          "PROSITE"      "REFSEQ"       "SYMBOL"       "UCSCKG"
[26] "UNIPROT"
```

```
res$symbol = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="SYMBOL",
                    multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$entrez = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="ENTREZID",
                    multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$name = mapIds(org.Hs.eg.db,
                  keys=row.names(res),
                  keytype="ENSEMBL",
                  column="GENENAME",
                  multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
head(res, 10)
```

log2 fold change (MLE): condition hoxa1_kd vs control_sirna

Wald test p-value: condition hoxa1 kd vs control sirna

DataFrame with 10 rows and 9 columns

	baseMean	log2FoldChange	lfcSE	stat	pvalue
	<numeric>	<numeric>	<numeric>	<numeric>	<numeric>
ENSG00000279457	29.913579	0.1792571	0.3248216	0.551863	5.81042e-01
ENSG00000187634	183.229650	0.4264571	0.1402658	3.040350	2.36304e-03
ENSG00000188976	1651.188076	-0.6927205	0.0548465	-12.630158	1.43989e-36
ENSG00000187961	209.637938	0.7297556	0.1318599	5.534326	3.12428e-08
ENSG00000187583	47.255123	0.0405765	0.2718928	0.149237	8.81366e-01
ENSG00000187642	11.979750	0.5428105	0.5215599	1.040744	2.97994e-01
ENSG00000188290	108.922128	2.0570638	0.1969053	10.446970	1.51282e-25
ENSG00000187608	350.716868	0.2573837	0.1027266	2.505522	1.22271e-02

	padj	symbol	entrez	name
	<numeric>	<character>	<character>	<character>
ENSG00000188157	9128.439422	0.3899088	0.0467163	8.346304 7.04321e-17
ENSG00000237330	0.158192	0.7859552	4.0804729	0.192614 8.47261e-01
ENSG00000279457	6.86555e-01	NA	NA	NA
ENSG00000187634	5.15718e-03	SAMD11	148398	sterile alpha motif ..
ENSG00000188976	1.76549e-35	NOC2L	26155	NOC2 like nucleolar ..
ENSG00000187961	1.13413e-07	KLHL17	339451	kelch like family me..
ENSG00000187583	9.19031e-01	PLEKHN1	84069	pleckstrin homology ..
ENSG00000187642	4.03379e-01	PERM1	84808	PPARGC1 and ESRR ind..
ENSG00000188290	1.30538e-24	HES4	57801	hes family bHLH tran..
ENSG00000187608	2.37452e-02	ISG15	9636	ISG15 ubiquitin like..
ENSG00000188157	4.21963e-16	AGRN	375790	agrin
ENSG00000237330	NA	RNF223	401934	ring finger protein ..

Q. Question

```
res = res[order(res$pvalue),]
write.csv(res, file = "deseq_results.csv")
```

Section 2: Pathway analysis

```
library(gage)
library(gageData)
library(pathview)
```

```
data(kegg.sets.hs)
data(sigmet.idx.hs)

# Focus on signaling and metabolic pathways only
kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]

# Examine the first 3 pathways
head(kegg.sets.hs, 3)
```

```
$`hsa00232 Caffeine metabolism`
[1] "10" "1544" "1548" "1549" "1553" "7498" "9"
```

```
$`hsa00983 Drug metabolism – other enzymes`
[1] "10" "1066" "10720" "10941" "151531" "1548" "1549" "1551"
[9] "1553" "1576" "1577" "1806" "1807" "1890" "221223" "2990"
```



```
[17] "3251" "3614" "3615" "3704" "51733" "54490" "54575" "54576"
[25] "54577" "54578" "54579" "54600" "54657" "54658" "54659" "54963"
[33] "574537" "64816" "7083" "7084" "7172" "7363" "7364" "7365"
[41] "7366" "7367" "7371" "7372" "7378" "7498" "79799" "83549"
[49] "8824" "8833" "9" "978"
```

```
$`hsa00230 Purine metabolism`
```

```
[1] "100" "10201" "10606" "10621" "10622" "10623" "107" "10714"
[9] "108" "10846" "109" "111" "11128" "11164" "112" "113"
[17] "114" "115" "122481" "122622" "124583" "132" "158" "159"
[25] "1633" "171568" "1716" "196883" "203" "204" "205" "221823"
[33] "2272" "22978" "23649" "246721" "25885" "2618" "26289" "270"
[41] "271" "27115" "272" "2766" "2977" "2982" "2983" "2984"
[49] "2986" "2987" "29922" "3000" "30833" "30834" "318" "3251"
[57] "353" "3614" "3615" "3704" "377841" "471" "4830" "4831"
[65] "4832" "4833" "4860" "4881" "4882" "4907" "50484" "50940"
[73] "51082" "51251" "51292" "5136" "5137" "5138" "5139" "5140"
[81] "5141" "5142" "5143" "5144" "5145" "5146" "5147" "5148"
[89] "5149" "5150" "5151" "5152" "5153" "5158" "5167" "5169"
[97] "51728" "5198" "5236" "5313" "5315" "53343" "54107" "5422"
[105] "5424" "5425" "5426" "5427" "5430" "5431" "5432" "5433"
[113] "5434" "5435" "5436" "5437" "5438" "5439" "5440" "5441"
[121] "5471" "548644" "55276" "5557" "5558" "55703" "55811" "55821"
[129] "5631" "5634" "56655" "56953" "56985" "57804" "58497" "6240"
[137] "6241" "64425" "646625" "654364" "661" "7498" "8382" "84172"
[145] "84265" "84284" "84618" "8622" "8654" "87178" "8833" "9060"
[153] "9061" "93034" "953" "9533" "954" "955" "956" "957"
[161] "9583" "9615"
```

```
foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)
```

```
1266 54855 1465 51232 2034 2317
-2.422719 3.201955 -2.313738 -2.059631 -1.888019 -1.649792
```

```
# Get the results
keggres = gage(foldchanges, gsets=kegg.sets.hs)

attributes(keggres)
```

```
$names
[1] "greater" "less" "stats"
```

```
# Look at the first few down (less) pathways
head(keggres$less)
```

	p.geomean	stat.mean	p.val
hsa04110 Cell cycle	8.995727e-06	-4.378644	8.995727e-06
hsa03030 DNA replication	9.424076e-05	-3.951803	9.424076e-05
hsa03013 RNA transport	1.375901e-03	-3.028500	1.375901e-03
hsa03440 Homologous recombination	3.066756e-03	-2.852899	3.066756e-03
hsa04114 Oocyte meiosis	3.784520e-03	-2.698128	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	8.961413e-03	-2.405398	8.961413e-03

	q.val	set.size	exp1
hsa04110 Cell cycle	0.001448312	121	8.995727e-06
hsa03030 DNA replication	0.007586381	36	9.424076e-05
hsa03013 RNA transport	0.073840037	144	1.375901e-03
hsa03440 Homologous recombination	0.121861535	28	3.066756e-03
hsa04114 Oocyte meiosis	0.121861535	102	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	0.212222694	53	8.961413e-03

```
pathview(foldchanges, pathway.id="hsa04110")
```


Info: Writing image file hsa04110.pathview.pdf

```
## Focus on top 5 upregulated pathways here for demo purposes only
keggrespathways <- rownames(keggres$greater)[1:5]

# Extract the 8 character long IDs part of each string
keggresids = substr(keggrespathways, start=1, stop=8)
keggresids
```

```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

Draw plots

```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa04640.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa04630.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa00140.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa04142.pathview.png

Info: some node width is different from others, and hence adjusted!

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa04330.pathview.png

Q. Can you do the same procedure as above to plot the pathview figures for the top 5 down-regulated pathways?

```
## Focus on top 5 downregulated pathways here for demo purposes only
keggrespathways <- rownames(keggres$less)[1:5]

# Extract the 8 character long IDs part of each string
keggresids = substr(keggrespathways, start=1, stop=8)
keggresids
```

```
[1] "hsa04110" "hsa03030" "hsa03013" "hsa03440" "hsa04114"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa04110.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa03030.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa03013.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa03440.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lisabear/Desktop/BIMM143/class14

Info: Writing image file hsa04114.pathview.png

Section 3: Gene Ontology

```
data(go.sets.hs)
data(go.subs.hs)

# Focus on Biological Process subset of GO
gobpsets = go.sets.hs[go.subs.hs$BP]

gobpres = gage(foldchanges, gsets=gobpsets, same.dir=TRUE)

lapply(gobpres, head)
```

\$greater

	p.geomean	stat.mean	p.val
GO:0007156 homophilic cell adhesion	8.519724e-05	3.824205	8.519724e-05
GO:0002009 morphogenesis of an epithelium	1.396681e-04	3.653886	1.396681e-04
GO:0048729 tissue morphogenesis	1.432451e-04	3.643242	1.432451e-04
GO:0007610 behavior	1.925222e-04	3.565432	1.925222e-04
GO:0060562 epithelial tube morphogenesis	5.932837e-04	3.261376	5.932837e-04
GO:0035295 tube development	5.953254e-04	3.253665	5.953254e-04

	q.val	set.size	exp1
GO:0007156 homophilic cell adhesion	0.1952430	113	8.519724e-05
GO:0002009 morphogenesis of an epithelium	0.1952430	339	1.396681e-04
GO:0048729 tissue morphogenesis	0.1952430	424	1.432451e-04
GO:0007610 behavior	0.1968058	426	1.925222e-04
GO:0060562 epithelial tube morphogenesis	0.3566193	257	5.932837e-04
GO:0035295 tube development	0.3566193	391	5.953254e-04

\$less

	p.geomean	stat.mean	p.val
GO:0048285 organelle fission	1.536227e-15	-8.063910	1.536227e-15
GO:0000280 nuclear division	4.286961e-15	-7.939217	4.286961e-15
GO:0007067 mitosis	4.286961e-15	-7.939217	4.286961e-15
GO:0000087 M phase of mitotic cell cycle	1.169934e-14	-7.797496	1.169934e-14
GO:0007059 chromosome segregation	2.028624e-11	-6.878340	2.028624e-11
GO:0000236 mitotic prometaphase	1.729553e-10	-6.695966	1.729553e-10

	q.val	set.size	exp1
GO:0048285 organelle fission	5.843127e-12	376	1.536227e-15
GO:0000280 nuclear division	5.843127e-12	352	4.286961e-15
GO:0007067 mitosis	5.843127e-12	352	4.286961e-15
GO:0000087 M phase of mitotic cell cycle	1.195965e-11	362	1.169934e-14

G0:0007059 chromosome segregation	1.659009e-08	142	2.028624e-11
G0:0000236 mitotic prometaphase	1.178690e-07	84	1.729553e-10

\$stats

	stat.mean	exp1
G0:0007156 homophilic cell adhesion	3.824205	3.824205
G0:0002009 morphogenesis of an epithelium	3.653886	3.653886
G0:0048729 tissue morphogenesis	3.643242	3.643242
G0:0007610 behavior	3.565432	3.565432
G0:0060562 epithelial tube morphogenesis	3.261376	3.261376
G0:0035295 tube development	3.253665	3.253665

Section 4. Reactome Analysis

```
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

```
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE)
```

Q: What pathway has the most significant "Entities p-value"? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

The pathway with most significant p values is the M phase of the mitotic cell cycle as it has the lowest V value. The pathways don't match exactly but I do see some similarities. This could be due to multiple reasons like data input, specificity parameters, anotation differences when analyzing, etc.