

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/323150680>

ENVIRONMENTAL RISKS OF AN INSECTICIDE (DIMILIN® 25 WP) ON THE SHRIMP PALAEMON ADSPERSUS: BIOCHEMICAL COMPOSITION OF CUTICLE AND OXIDATIVE STRESS

Article in *Fresenius environmental bulletin* · February 2018

CITATIONS

5

READS

1,373

2 authors, including:



Noureddine Soltani

Badji Moktar-Annaba University

469 PUBLICATIONS 4,602 CITATIONS

SEE PROFILE

ENVIRONMENTAL RISKS OF AN INSECTICIDE (DIMILIN® 25 WP) ON THE SHRIMP *PALAEMON ADSPERSUS*: BIOCHEMICAL COMPOSITION OF CUTICLE AND OXIDATIVE STRESS

Hanene Lechekhab^{1,2}, Nouredine Soltani^{2,*}

¹Department of Biology, Faculty of Sciences, Chadli Benjdid University, 36000-El Tarf, Algeria

²Laboratory of Applied Animal Biology, Faculty of Sciences, Badji Mokhtar University, 23000-Annaba, Algeria

ABSTRACT

The present study aimed to assess, under laboratory conditions, the side effects of a trade formulation of diflubenzuron (Dimilin® 25 WP), a chitin synthesis inhibitor (CSI), on the biochemical composition of the cuticle and on the level of the biomarker of environmental stress glutathione (GSH), in a non-target species *Palaemon adspersus* (Rathke, 1837) (Crustacea, Decapoda). The insecticide was added to the shrimps' rearing water at two concentrations: 16 ng of active ingredient/L and 1 µg/L. The shrimps were exposed to Dimilin from stage A (newly molted shrimp) to Stage C (Intermolt), where mineralization and deposition of the cuticular layers are completed. The results show a significant decrease ($p < 0.05$) in the amounts of chitin in the treated batches as compared to controls for both assays. Moreover, there was a significant difference ($p < 0.05$) in the cuticular proteins between treated and control individuals. However, the compound had no significant effect on the amounts of calcium salts. Regarding GSH level, there was a significant inhibition ($p < 0.05$) in treated individuals as compared to controls. Our results show an induction of the detoxification system and confirm the mechanism of action of Dimilin on chitin biosynthesis, and indicate that this product can pose side-effects on non-target Arthropod organism like shrimps.

KEYWORDS:

Crustacea, *Palaemon adspersus*, Insecticides, Diflubenzuron, Cuticle, Chitin, Calcium salts, Glutathione.

INTRODUCTION

Human actions are one of the main factors affecting the quality of the environment and water [1]. Pollution of the aquatic ecosystem is due to human development and its side effects such as radioactive discharges, organic matter, heavy metals [2, 3], industrial and domestic discharges, and the intensive use of pesticides by farmers [4, 5]. Some pesticides

are applied directly to aquatic systems to reduce numbers of mosquito larvae (larvicides) and thereby reduce transmission of pathogens that mosquitoes vector to humans and wildlife [6]. Environmental imperatives [7] have pushed research towards the use of natural pesticides or biopesticides [8, 9, 10] and prompted the phytosanitary industry to develop more selective molecules with high metabolic and environmental stability [11, 12], acting on biochemical processes specific to the target organisms [13]. These new products are the insect growth disruptors (IGDs). These include the inhibitors of chitin synthesis which interferes in the formation of the cuticle [14, 15].

Dimilin (25% WP) is a trade formulation of diflubenzuron, a benzoylurea derivative and inhibitor of the synthesis of chitin [16, 17, 18, 19], one of the major compounds of the Arthropod cuticle. It is an approved insecticide in Algeria and is used on a large scale to fight forest defoliating insects. In addition, DFB was found a potent insecticide for controlling mosquito pullulations [20]. It was also Its leaching by rains contaminates the aquatic ecosystem including the sediments in which all stages of the life cycle of a crustacean take place. This prompted to study the degradation of the insecticide (DFB) in seawater [21].

In this study, we used the shrimp *Palaemon adspersus* (Rathke, 1837), a very abundant shrimp in the Northeastern region of Algeria and is of a relatively important interest to the local fishing industry. This shrimp was used as a bioindicator species in order to assess the quality of the water from the Mellah lagoon (Northeast, Algeria) [22] and as well as a model organism to test the effects of Novaluron, another chitin synthesis inhibitor (CSI), on chitin and cuticular proteins [23]. The Crustaceans from the gulf of Annaba have been the subject of several studies on the physiological [22-25] and toxicological aspects [23, 26, 27, 28, 29]. Within this context, the current study is a continuation of the previous research conducted by [23] who determined the amount of chitin and cuticular proteins of the same species when exposed to another molecule (Novaluron). This research adds to the latter in that it shows

the amount of calcium salts content of the cuticle and also measures the level of glutathione (GSH), a biomarker that plays a central role in intracellular antioxidant defense processes.

MATERIALS AND METHODS

Shrimp collection. *P. adspersus* (Decapoda, Palaemonidae) is a common shrimp in Norwegian coast and South of the Baltic Sea to the Mediterranean. This species is transparent, fairly uniform with yellow strips at the level of articulations and presents dark chromatophores on the ventral half of the rostrum. The shrimps (length: 8-10 cm; weight: 12-16 g) were collected from the lagoon of El Mellah (Northeast, Algeria) and transported to the laboratory. They were reared in aquariums (100 × 60 cm) filled with sea water under laboratory conditions (salinity 37 psu; temperature 24-27°C; photoperiod 14h light). They were daily fed with fresh mussels administered in the afternoon.

Shrimp datation. In crustaceans, the molt cycle, corresponding to the modifications that occur between two successive molts, is subdivided into five stages: A (early post-molt), B (late post-molt), C (intermolt) and D (pre-molt) and molting (E). Molt stages were determined according to [30] by microscopic examination of the uropod setae. Under these conditions, *P. adspersus* has a molt cycle of 20 ± 2 days, and the relative duration of each stage is 9% for A+B, 36% for C, and 59% for D.

Insecticide and treatment. Dimilin® (Wettable powder, 25% active ingredient, a.i.), a trade formulation of diflubenzuron, is an insecticide belonging to the benzoylphenylurea derivatives. It was kindly provided by Pr. G. Smagghe (Ghent University, Belgium). Dimilin was added to the rearing seawater at a final concentration of 1 µg active ingredient/L according to concentrations tested on several shrimp species [27, 28] and 16 ng/L corresponding

to the LC_{50} determined against fourth instar larvae of the most abundant mosquito species *Culex pipiens* [20]. Newly molted adult shrimps (0-8 h old) were continuously exposed to treatment up to intermolt. Control shrimps were reared in seawater alone.

Biochemical procedure. Shrimps were sampled at the beginning of the inter-molt and killed by freezing. The entire carapace was removed and the cuticle was separated from adhering tissue, washed and dried at 60°C to constant weight and decalcified by means of 10% trichloroacetic acid (TCA) according to [29]. The decalcified cuticle was extracted in 2 N NaOH at 110°C for 3-4 h. The colorless residues were washed in distilled water and dried at 60°C to constant weight. The weight loss was supposed to be due to removal of proteins, and the residue to be chitin.

Glutathione assay. The glutathione was determined according to the method of [32] previously described [33]. The flesh of the shrimp was homogenized in 1 ml EDTA solution (0.02M; pH 6) and subjected to a deproteinisation with sulfosalicylic acid (SSA) at 0.025%. Then the homogenates were maintained for 15 min on ice and centrifuged (3000g, 15 min). The mixture was vortexed, left for 15 minutes in an ice bath and centrifuged (1000 rpm/min for 5 min). The supernatant (0.5 mL) was taken and then 1 mL of Tris-EDTA buffer (0.02 M, pH 9.6) and 0.025 mL of DTNB (0.01 M) were added. The mixture was then left for 5 min at room temperature. The optical density was measured at a wavelength of 412 nm after 5 min. GSH levels are expressed in µM /mg proteins.

Protein assay. The protein assay was prepared according to the method of [34] described previously [27]. This uses the Coomassie brilliant blue (BBC) G 250 as a reagent and a solution of bovine serum albumin (BSA) as a standard. The absorbance was read by the spectrophotometer at a wavelength of 595 nm.



FIGURE 1
Palaemon. adspersus (Crustacea, Decapoda).




FIGURE 2
Molecular structure of diflubenzuron.

TABLE 1

Effect of Dimilin (1µg/L and 16ng/L) on cuticle chitin levels (%) of *P. adspersus* at intermolt stage in control and treated series (mean ± SD, n = 4-5).

Treatment	Control	DFB 16 ng/L	DFB 1 µg/L	p
Level (%)	58.46 ± 13.41 a	45.76 ± 8.47 b	30.68 ± 10.55 b	0.003

Values followed by a different letter are significantly different ($p < 0.05$).

TABLE 2

Effect of Dimilin (1µg/L and 16ng/L) on cuticle protein levels (%) of *P. adspersus* at intermolt stage in control and treated series (mean ± SD, n = 4-5).

Treatment	Control	DFB 16 ng/L	DFB 1 µg/L	p
Level (%)	41.54 ± 13.41 a	69.31 ± 10.55 b	69.17 ± 7.26 b	0.001

Values followed by a different letter are significantly different ($p < 0.05$).

TABLE 3

Effect of Dimilin (1µg/L and 16ng/L) on cuticle calcium salt levels (%) of *P. adspersus* at intermolt stage in control and treated series (mean ± SD, n = 4-5).

Treatment	Control	DFB 16 ng/L	DFB 1 µg/L	p
Level (%)	42.26 ± 6.21 a	49.40 ± 13.04 a	30.83 ± 7.26 a	0.065

Values followed by a different letter are significantly different ($p < 0.05$).

TABLE 4

Effect of Dimilin (1µg/L and 16ng/L) on GSH amounts (µM/mg protein) in the flesh of *P. adspersus* at intermolt stage in control and treated series (mean ± SD, n = 4-5).

Treatment	Control	DFB 16 ng/L	DFB 1 µg/L	P
Amounts	0.14 ± 0.02 a	0.13 ± 0.01a	0.09 ± 0.04 a	0.026

Values followed by a different letter are significantly different ($p < 0.05$).

Statistical analysis. Data are expressed as mean ± standard deviation (m ± SD) and were subjected to one-way analysis of variance (ANOVA) followed by Tukey test. All statistical analyses were performed using Minitab software (version 16, PA, State College, USA). With $p < 0.05$ considered as a statistically significant difference.

RESULTS

Effect on chitin levels. The amount of cuticular chitin noted in the control series was 58.46 ± 13.41%. The treatment with Dimilin led to a significant decrease ($p = 0.003$) in the chitin level (Table 1). The lowest level was observed in individuals exposed to a dose of 1 µg/L (30.68 ± 10.55%) compared with the dose of 16 ng/L (45.76 ± 8, 47%).

Effect on protein levels. Under normal conditions, the protein level at the intermolt is (41.54 ± 13.41%). In the treated series, there was a significant increase ($p \leq 0.001$) in the level of proteins for both doses: 69.31 ± 10.55% for the dose of 16 ng/L, and 69.17 ± 7.26% for the dose of 1 µg/L (Table 2).

Effect on calcium salt levels. The result for the calcium salt level in controls was 42.26 ± 6.21%. As for the treated series, it was noted that the level of calcium salts did not vary significantly ($p = 0.065$) compared to the levels in the control series (Table 3).

Effect on glutathione amounts. For the treated series, the GSH amounts recorded in stage C decreased significantly (0.13 ± 0.01 µM/mg proteins for the dose of 16 ng/L and 0.09 ± 0.04 for the dose of 1 µg/L) compared to the control series (0.14 ± 0.02 µM/mg proteins) (Table 4). The comparison of the mean values in the control and treated series shows that there was a significant difference in the GSH level ($p < 0.05$).

DISCUSSION

Crustaceans have an exoskeleton or cuticle which acts as a selective barrier between the internal and external environments [35, 36]. This non-extensible exoskeleton is periodically disposed of and replaced by a new cuticle during molting. The cuticle is a complex macromolecular structure. It is mainly composed of chitin fibers to which proteins bind.

These fibres are in turn associated to a protein matrix [37] and impregnated with limestone [38].

Regarding the effect of Dimilin on the amount of chitin at the intermolt, the compound was found to significantly reduce the level of chitin in the treated series as compared to controls. Our findings are consistent with those commonly reported and which assert that benzoylphenylurea derivatives interfere with the molting process by disrupting cuticular secretion via chitin synthesis [27, 39, 40, 41]. According to [42], diflubenzuron interferes with enzymes that contribute to the synthesis of chitin in crustacean species such as the benthic copepod *Tisbe battagliai*. Our results confirm previous studies using the same product on another shrimp species, *Penaeus kerathurus* [27]. It was also reported that Novaluron, another inhibitor of chitin synthesis, caused a decrease in the level of cuticular chitin of *P. adspersus* [23].

The proteins play a fundamental role within the organism of all living biological species [43]. Our results show a significant increase in their levels in the treated series compared to the control series. This is reflected in the composition of the cuticle, which consists of chitin which binds to proteins. Thus, there is a strong correlation between the two components [44]. Similar results have been reported by [26] and concluded that treatment with diflubenzuron causes an increase in the level of cuticular proteins in *P. kerathurus*. The same effect was observed in the treatment of *P. adspersus* with Novaluron [23].

Finally, regarding the effect of this xenobiotic on the calcium salt content, it is noted that Dimilin has no effect on the calcium salt levels. The same results were obtained in *P. kerathurus* following treatment with Dimilin [26]. Similarly, the work of [22] states that calcium salts in *P. adspersus* are not affected.

During the environmental stress caused by the presence of pollutants, biochemical responses are immediate in exposed organisms [45]. Little work has been done on the detoxification system in crustaceans. GSH is considered as one of the most important antioxidant agents involved in the protection of cell membranes against free radicals damage [46, 47, 48]. So in the treated series, our experiments show that Dimilin causes a decrease in GSH amounts measured at the intermolt (Stage C). In the work of [49], it was noted that the exposition of marbled crayfish (*Procambarus fallax f. virginalis*) to Cyperkill 25 EC (Pyrethroid insecticide) in concentration of 0.05 µg/L caused significant increase in GSH amounts ($p < 0.05$). Other research [50] shows that in shrimp *Palaemonetes pugio* and after 96-h exposure to phenothrin at five concentrations, the sublethal biomarkers glutathione were examined and there were no statistically significant differences in these levels in adults or larvae compared to controls.

CONCLUSION

Our experiments assessed the impacts of an inhibitor of chitin synthesis, Dimilin, on a non-target aquatic Arthropod *P. adspersus*. The obtained results confirm its mechanism of action on chitin biosynthesis. This compound can show secondary effects even at low doses on this edible shrimp species.

ACKNOWLEDGEMENTS

This research was supported by the National Fund for Scientific Research of Algeria (Laboratory of Applied Animal Biology) and the Ministry of Higher Education and Scientific Research of Algeria (CNEPRU project F01120140104 to Pr. N. Soltani).

REFERENCES

- [1] Ippolito, J.A., Novak, J.M., Busscher, W.J., Ahmedna, M., Rehrah, D. and Watt, D.W. (2012) wicthg rass biochar affects two Aridisols. *J. Environ. Qual.* 41, 1123–1130.
- [2] Onivogui, G., Balde, S., Bangoura, K. and Barry, M.K. (2015) Évaluation des risques de pollution en métaux lourds (Hg, Cd, Pb, Co, Ni, Zn) des eaux et des sédiments de l'estuaire du fleuve Konkouré (Rep. de Guinée). *Afrique Science: Revue Internationale des Sciences et Technologie.* 9(3), 36-44.
- [3] El Hassani, F., Boushaba, A., Raïs, N. and Benabidate, L. (2016) Etude de la Contamination par les Métaux Lourds des Eaux et Des Sédiments au Voisinage de la Mine De Tighza (Maroc Central Oriental). *European Scientific Journal.* 12(6), 176-184.
- [4] Delorme, R., Lerroux, P. and Gaillardon, P. (2002) Évolution des produits phytosanitaires à usage agricole: Les insecticides-acaricides. *Phytoma, La défense des Végétaux.* 548, 7- 13.
- [5] Mokbel, H., El Moll, A., Liaud, C. and Millet, M. (2015) Développement de méthodes d'analyses pour évaluer les niveaux de contamination atmosphérique d'OCP, PCB ET HAP. *Lebanese Science Journal.* 16(2), 47-64.
- [6] Lawler, S.P. (2017) Environmental safety review of methoprene and bacterially-derived pesticides commonly used for sustained mosquito control. *Ecotoxicology and Environmental Safety.* 139, 335-343.
- [7] Paoletti, M.G. and Pimentel, D. (2000) Environmental risks of pesticides versus genetic engineering for agricultural pest control. *J. Agric. Environ. Ethics.* 12(3), 279-303.

- [8] Maiza, A., Aribi, N., Smagghe, G., Kilani-Morakchi, S., Bendjedid, M. and Soltani, N. (2013) Sublethal effects on reproduction and biomarkers by spinosad and indoxacarb in cockroaches *Blattella germanica*. Bulletin. Insectol. 66(1), 11-20.
- [9] Tomé, H.V.V., Martins, J.C., Corrêa, A.S., Galdino, T.V.S., Picanço, M.C. and Guedes, R.N.C. (2013) Azadirachtin avoidance by larvae and adult females of the tomato leafminer *Tuta absoluta*. Crop Protection. 46, 63-69.
- [10] Cepeda-Palacios, R., Servín, R., Ramírez-Orduña, J.M., Ascencio, F., Dorchie, P. and Angulo-Valadez, C.E. (2014) *In vitro* and *in vivo* effects of neem tree (*Azadirachta* Références bibliographiques 70 indica A. Juss) products on larvae of the sheep nose bot fly (*Oestrus ovis* L. Diptera: Oestridae). Veterinary Parasitology. 200, 225-228.
- [11] Dhadialla, T.S., Carlson, G.R. and Lee, D.P. (1998) New insecticides with ecdysteroidal and juvenile hormone activity. Annu. Rev. Entomol. 43, 545-569.
- [12] Cassida, J. and Quistad, B. (1998) Golden age of insecticide research: past present or future? Annu. Rev. Entomol. 43, 1-16.
- [13] El-Sayed, K.A., Dunbar, D.C., Perry, T.L., Wilkins, S.P. and Hamann, M.T. (1997) Marine natural products as phototype insecticidal agents. J. Agri. Food chem. 45, 2735-2739.
- [14] Tunaz, H. and Uygün, N. (2004) Insect Growth Regulators for Insect Pest Control. Turk J. Agric For. 28, 377-387.
- [15] Sun, R., Liu, C., Zhang, H. and Wang, Q. (2015) Benzoylurea Chitin Synthesis Inhibitors. Journal of agricultural and food chemistry. 63(31), 6847-6865.
- [16] Soltani, N., Soltani-Mazouni, N., Quennedey, B. and Delachambre, J. (1996) Protein synthesis in developing ovaries of mealworm under *in vitro* condition: Effects of diflubenzuron. J. Stored Prod Res. 32(3), 205-212.
- [17] Gade, G., Hoffmann, R.H. and Spring, J.H. (1997) Hormonal regulation in insects: facts, gaps and future direction. Physiological reviews. 77(4), 963-1032.
- [18] Oberlander, H. and Silhacek, D.L. (1998) New perspectives on the mode of action of benzoylphenylurée insecticides. In: Ishaaya, I. and Degheele, D. (eds.) Insecticides with novel modes of action. Springer-Verlag, Berlin, Germany, 92-105.
- [19] Soltani, N., Rehimi, H., Beldi, H. and Bendal, I.F. (1999) Activité du triflumron sur *Culex pipiens* (Diptera: Culicidae) et impact sur deux espèces larvaires non visées. Ann. Soc. Entomol. Fr. 35, 502-508.
- [20] Rehimi, N. (2004) Etude de la reproduction chez *Culex pipiens pipiens*. Aspects: morphologique, éthologique et physiologique. Effets de quelques inhibiteurs du développement sur quelques paramètres biologiques. Thèse de doctorat d'état en Sciences Naturelles, mention Biologie Animale. Badji Mokhtar University of Annaba, Algeria. PhD Thesis, 195p.
- [21] Soltani, N. and Morsli, S.M. (2003) Quantification du Dimilin par Chromatographie Liquide à Haute Performance: Etude de sa Dégradation dans L'eau de Mer. Journal de Recherche Océanographique. 28, 118-120.
- [22] Benradia, H., Berghiche, H. and Soltani, N. (2016) Measure of environmental stress biomarkers in the shrimp *Palaemon adspersus* from the mellah lagoon (Algeria): Spatial and temporal variations. Fresen. Environ. Bull. 25, 2563-2566.
- [23] Berghiche, H., Benradia, H. and Soltani, N. (2016) Impact of insect growth disruptor, novaluron, on biochemical composition of cuticle from the shrimp *Palaemon adspersus*. Journal of Entomology and Zoology Studies. 4(2), 147-151.
- [24] Soltani, N. and Bezzazel, N. (2002) Profil des ecdystéroïdes hémolympatiques et corrélations avec la sécrétion cuticulaire au cours d'un cycle de mue chez la crevette, *Penaeus kerathurus*. J. Rech. Océanogr. 27(3-4), 226-231.
- [25] Derbal, F. and Soltani, N. (2008) Cycle cuticulaire et variations de la protéinémie et de la lipémie chez la crevette royale *Penaeus kerathurus* (Forsk., 1775) des côtes Est algériennes. Science & Technologie. 28, 80- 86.
- [26] Morsli, M.S., Soltani N. (2003) Effets d'un insecticide inhibiteur de la synthèse de la chitine, le diflubenzuron, sur la cuticule de la crevette *Penaeus kerathurus*. J. Rech. Océanogr. 28(12), 85-88.
- [27] Soltani, N., Lechekhab, H. and Smagghe, G. (2009) Impact of the insect growth regulator diflubenzuron on biochemical composition of cuticle of the shrimp *Penaeus kerathurus*. Communications in Agricultural and Applied Biological Sciences. 74(1), 137-141.
- [28] Morsli, S.M., Merad, I., Khebbab, M.E.H. and Soltani, N. (2015) Potential hazards of a chitin synthesis inhibitor diflubenzuron in the shrimp *Penaeus kerathurus*: biochemical composition of the hemolymph and muscle during the molt cycle. Advances in Environmental Biology. 9(3), 518-525.
- [29] Gheid, S., Nadji, S. and Khebbab, M.E.H. (2011) Taux des lipides et des protéines et composition en acides gras du tissu comestible des crustacés et des mollusques pêchés en Algérie: effet du halofenozide (RH-0345) sur la composition en acides gras de *Penaeus kerathurus* (Crustacé, Décapode). Synthèse. 22, 41-47.

- [30] Robertson, L., Bray, W., Leung-Trujillo, J. and Lawrence, A. (1987) Practical molt staging of *Penaeus setiferus* and *Penaeus stylirostris*. L. Journal of the World Aquaculture. Society. 18(3), 180-185.
- [31] Welinder, B.S. (1974) The Crustacean cuticle. I. Studies on the composition of the cuticle. Comp. Biochem. 47(A), 779-787.
- [32] Weckberker, G. and Cory, G. (1988) Ribonucleotide reductase activity and growth of glutathione-depleted mouse leukemia 1210 cells *in vitro*. Cancer Letters. 40, 257-264.
- [33] Zaidi, N. and Soltani, N. (2011) Environmental risks of two chitin synthesis inhibitors on *Gambusia affinis*: Chronic effects on growth and recovery of biological responses. Biological Control. 59, 106-113.
- [34] Bradford, M.M. (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. 72, 254-278.
- [35] Nagasawa, H. (2012) The crustacean cuticle: structure, composition and mineralization. Front Biosci (Elite Ed). 4, 71-200.
- [36] Prabu, K. and Natarajan, E. (2012) Bioprospecting of shells of crustaceans. International Journal of Pharmacy and Pharmaceutical Sciences. 4(4), 1-3.
- [37] Hepburn, H.R. (1985) Structure of the integument. In: Kerkut, G.A. and Gilbert, L.I. (Eds.) Comprehensive insect physiology biochemistry and pharmacology. Pergamon Press, Oxford. 3, 1-58.
- [38] Dennell, R. (1947) The occurrence and significance of phenolic hardening in the newly formed cuticle of Crustacea Decapoda. Proc. R. Soc. 134(B), 485-503.
- [39] Dhadialla, T.S., Retnakaran, A. and Smagghe, G. (2005) Insect growth and development disrupting insecticides. In: Gilbert, L.I., Kostas, I. and Gill, S. (eds.) Comprehensive Insect Molecular Science. Vol. 6. Pergamon Press, New York, NY, 55-116.
- [40] Dhadialla, T.S., Retnakaran, A. and Smagghe, G. (2010) Insect growth and development disrupting insecticides. In: Gilbert, L.I., Gill, S.S. (Eds.) Insect Control: Elsevier. 121-184.
- [41] Salokhe, S.G., Deshpande, S.G., Pakhle, K. and Maniyare, A.B. (2013) IGR, Lufenuron, alters chitin and total soluble protein content of *Aedes aegypti* larvae during development of *Aedes aegypti*. Curr Sci. 99(9), 1256-59.
- [42] Macken, A., Lillicrap, A. and Langford, K. (2015) Benzoylurea pesticides used as veterinary medicines in aquaculture: Risks and developmental effects on non-target crustaceans. Environmental Toxicology and Chemistry. 34(7), 1533-1542.
- [43] Mahler, H. and Cordes, E. (1968) Biological chemistry, Harper and Row. In: Maïza, A., Kilani, S., Farine, J.P., Smagghe, G., Aribi, N. and Soltani, N. (2004) Reproductive effects in German cockroaches by ecdysteroid agonist (RH-0345), juvenile hormone analog (methoprene and carbamate benfuracab). Comm. Appl. Biol. Gent University. 69(3), 257-266.
- [44] Lechekheb, H. and Soltani, N. (2009) Impact d'un inhibiteur de la synthèse de la chitine (dimilin®) sur la composition biochimique de la cuticule de *Penaeus kerathurus*. Bulletin de l'INSTM NS. 13, 96-100.
- [45] David, E., Tanguy, A., Riso, R., Quiniou, L., Laroche, L. and Morago, D. (2012) Responses of Pacific oyster *Crassostrea gigas* populations to abiotic stress environmentally contrasted estuaries along the Atlantic coast of France. Aquatic Toxicology. 109, 70-79.
- [46] Sies, H. and Akerboom, T.P.M. (1984) Glutathione disulfide (GSSG) efflux, from cells and tissues. Methods in Enzymology. 105, 445-451.
- [47] Martinez-Alvarez, R.M., Morales, A.E. and Sanz, A. (2005) Antioxidant defences in fish: biotic and abiotic factors. Reviews in Fish biology and Fisheries. 15, 75-88.
- [48] Lam, P.K.S. (2009) Use of biomarkers in environmental monitoring. Ocean and Coastal Management. 52, 348-354.
- [49] Lidova, J., Stara, A., Kouba, A. and Velisek, J. (2016) The effects of cypermethrin on oxidative stress and antioxidant biomarkers in marbled crayfish (*Procambarus fallax f. virginalis*). NeuroEndocrinol. Lett. 1, 53-59.
- [50] Key, P.B., Chung, K.W., Hoguet, J., Sapozhnikova, Y. and Delorenzo, M.E. (2011) Toxicity of the mosquito control insecticide phenothrin to three life stages of the grass shrimp (*Palaeomonetes pugio*). J. Environ Sci Health B. 46(5), 426-31.

Received: 18.10.2017

Accepted: 30.12.2017

CORRESPONDING AUTHOR

Nouredine Soltani

Laboratory of Applied Animal Biology,
Department of Biology,
Faculty of Sciences,
University Badji Mokhtar of Annaba,
23000 Annaba – Algeria

e-mail: noureddine.soltani@univ-annba.org