

# L25: Interpolation

Join the dots

Lucas A. J. Bastien

E7 Spring 2017, University of California at Berkeley

March 20, 2017

Version: release

# Announcements

**Lab 09 is due on March 24 at 12 pm (noon)**

## **Today:**

- ▶ Interpolation (chapter 14)
  - ▶ Introduction and motivation
  - ▶ Nearest neighbor interpolation
  - ▶ Linear interpolation
  - ▶ Lagrange polynomial
  - ▶ Cubic splines

## **Wednesday:**


- ▶ Series (chapter 15)
- ▶ Presentation of the final programming project

## **Friday:**

- ▶ Discussion

## Introduction to interpolation (well water contamination)

Consider the following (equally-spaced) houses and the corresponding water quality measurements (mock values, arbitrary units)




What is the water quality value in the fourth house from the left?

- (A) 50
- (B) 30
- (C) 26
- (D) 40
- (E) I am not sure

## Introduction to interpolation (well water contamination)

Consider the following (equally-spaced) houses and the corresponding water quality measurements (mock values, arbitrary units)



What is the water quality value in the fourth house from the left?

(A) 50


(B) 30

(C) 26


(D) 40

(E) I am not sure

# Introduction to interpolation (well water contamination)



# Introduction to interpolation (well water contamination)



# Introduction to interpolation


**Interpolation:** starting from discrete data, estimate values between data points, by using functions that go through all the data points

# Introduction to interpolation

**Interpolation:** starting from discrete data, estimate values between data points, by using functions that go through all the data points

## Linear regression

- ▶ Draw best-fit line going through a cloud of points
- ▶ Fitted line does **not always** go through all the data points






# Introduction to interpolation

**Interpolation:** starting from discrete data, estimate values between data points, by using functions that go through all the data points


## Linear regression

- ▶ Draw best-fit line going through a cloud of points
- ▶ Fitted line does **not always** go through all the data points




## Interpolation

- ▶ “Join the dots”, filling missing values between data points
- ▶ Interpolation line **does** go through all the data points




# Introduction to interpolation

Today, we learn about **four interpolation methods**: **nearest neighbor**, **linear interpolation**, **Lagrange polynomial**, and **cubic splines**




# Introduction to interpolation

Today, we learn about **four interpolation methods**: **nearest neighbor**, **linear interpolation**, **Lagrange polynomial**, and **cubic splines**




# Introduction to interpolation

Today, we learn about **four interpolation methods**: **nearest neighbor**, **linear interpolation**, **Lagrange polynomial**, and **cubic splines**




# Introduction to interpolation

Today, we learn about **four interpolation methods**: **nearest neighbor**, **linear interpolation**, **Lagrange polynomial**, and **cubic splines**



# Introduction to interpolation


Today, we learn about **four interpolation methods**: **nearest neighbor**, **linear interpolation**, **Lagrange polynomial**, and **cubic splines**



# Nearest neighbor interpolation

## Nearest neighbor interpolation:


Associate, to each value of  $x$ , the  $y_i$  value associated with the closest  $x_i$



# Linear interpolation

## Linear interpolation:

Link any two consecutive points with a straight line






# Linear interpolation

## Linear interpolation:


Link any two consecutive points with a straight line



# Linear interpolation

## Linear interpolation:

Link any two consecutive points with a straight line




$$y = y_i + \frac{y_{i+1} - y_i}{x_{i+1} - x_i}(x - x_i) \quad \text{for } x \in [x_i, x_{i+1}]$$

# Linear interpolation

## Linear interpolation:

Link any two consecutive points with a straight line




The equation varies between different pairs of points

# Lagrange polynomial

Consider a set of  $m$  data points  $(x_i, y_i), i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are different from each other

## Lagrange polynomial:

Polynomial of least degree that goes through all the points




# Lagrange polynomial

Consider a set of  $m$  data points  $(x_i, y_i)$ ,  $i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are different from each other

## Lagrange polynomial:

Polynomial of least degree that goes through all the points



Consider a set of  $m$  data points  $(x_i, y_i), i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are different from each other. How many different polynomials of degree  $m$  that go through all these data points can we find?

- (A) Zero
  - (B) One and only one
  - (C) An infinite number
  - (D) It depends on the data
-

# Lagrange polynomial

---

Consider a set of  $m$  data points  $(x_i, y_i), i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are different from each other. How many different polynomials of degree  $m$  that go through all these data points can we find?

- (A) Zero
  - (B) One and only one
  - (C) An infinite number
  - (D) It depends on the data
-

# Lagrange polynomial

---

Consider a set of  $m$  data points  $(x_i, y_i), i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are different from each other. We calculate the Lagrange polynomial for this data set. Which of the following statements are true?

- (A) The degree of the polynomial is  $m - 1$  or less
  - (B) The degree of the polynomial is  $m$
  - (C) The degree of the polynomial is  $m$  or more
  - (D) The degree of the polynomial is  $m + 1$  or more
  - (E) None of the above
-



# Lagrange polynomial

---


Consider a set of  $m$  data points  $(x_i, y_i), i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are different from each other. We calculate the Lagrange polynomial for this data set. Which of the following statements are true?

- (A) The degree of the polynomial is  $m - 1$  or less
  - (B) The degree of the polynomial is  $m$
  - (C) The degree of the polynomial is  $m$  or more
  - (D) The degree of the polynomial is  $m + 1$  or more
  - (E) None of the above
-

# Lagrange polynomial

## Lagrange polynomial $L$ :


Polynomial of least degree that goes through all the points



# Lagrange polynomial

## Lagrange polynomial $L$ :

Polynomial of least degree that goes through all the points




$$L(x) = \sum_{i=1}^m y_i l_i(x)$$

# Lagrange polynomial

## Lagrange polynomial $L$ :

Polynomial of least degree that goes through all the points



$$L(x) = \sum_{i=1}^m y_i l_i(x)$$


$l_i$ : basis polynomial, such that:

$$l_i(x_j) = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{if } j \neq i \end{cases}$$

# Lagrange polynomial

## Lagrange polynomial $L$ :

Polynomial of least degree that goes through all the points



$$L(x) = \sum_{i=1}^m y_i l_i(x)$$

$l_i$ : basis polynomial, such that:


$$l_i(x_j) = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{if } j \neq i \end{cases}$$

$$l_i(x) = \prod_{\substack{j=1 \\ j \neq i}}^m \frac{x - x_j}{x_i - x_j}$$

# Lagrange polynomial

$l_1$  is zero at all the  $x_i$ 's except for  $x_1$ , where it is equal to 1

$y_1 \times l_1$  is zero at all the  $x_i$ 's except for  $x_1$ , where it is equal to  $y_1$




$$y_1 l_1(x) = y_1 \prod_{\substack{j=1 \\ j \neq 1}}^m \frac{x - x_j}{x_1 - x_j}$$

# Lagrange polynomial

$l_2$  is zero at all the  $x_i$ 's except for  $x_2$ , where it is equal to 1

$y_2 \times l_2$  is zero at all the  $x_i$ 's except for  $x_2$ , where it is equal to  $y_2$




$$y_2 l_2(x) = y_2 \prod_{\substack{j=1 \\ j \neq 2}}^m \frac{x - x_j}{x_2 - x_j}$$

# Lagrange polynomial

$l_3$  is zero at all the  $x_i$ 's except for  $x_3$ , where it is equal to 1

$y_3 \times l_3$  is zero at all the  $x_i$ 's except for  $x_3$ , where it is equal to  $y_3$




$$y_3 l_3(x) = y_3 \prod_{\substack{j=1 \\ j \neq 3}}^m \frac{x - x_j}{x_3 - x_j}$$



# Lagrange polynomial

$l_4$  is zero at all the  $x_i$ 's except for  $x_4$ , where it is equal to 1

$y_4 \times l_4$  is zero at all the  $x_i$ 's except for  $x_4$ , where it is equal to  $y_4$




$$y_4 l_4(x) = y_4 \prod_{\substack{j=1 \\ j \neq 4}}^m \frac{x - x_j}{x_4 - x_j}$$

# Lagrange polynomial

$l_5$  is zero at all the  $x_i$ 's except for  $x_5$ , where it is equal to 1

$y_5 \times l_5$  is zero at all the  $x_i$ 's except for  $x_5$ , where it is equal to  $y_5$




$$y_5 l_5(x) = y_5 \prod_{\substack{j=1 \\ j \neq 5}}^m \frac{x - x_j}{x_5 - x_j}$$

# Lagrange polynomial

$l_6$  is zero at all the  $x_i$ 's except for  $x_6$ , where it is equal to 1

$y_6 \times l_6$  is zero at all the  $x_i$ 's except for  $x_6$ , where it is equal to  $y_6$




$$y_6 l_6(x) = y_6 \prod_{\substack{j=1 \\ j \neq 6}}^m \frac{x - x_j}{x_6 - x_j}$$

# Lagrange polynomial

$l_7$  is zero at all the  $x_i$ 's except for  $x_7$ , where it is equal to 1

$y_7 \times l_7$  is zero at all the  $x_i$ 's except for  $x_7$ , where it is equal to  $y_7$



$$y_7 l_7(x) = y_7 \prod_{\substack{j=1 \\ j \neq 7}}^m \frac{x - x_j}{x_7 - x_j}$$

# Lagrange polynomial

**Alternative approach:** let Matlab do the work

For  $m$  data points, we are looking for a polynomial of degree  $m - 1$  or smaller:

$$L(x) = a_{m-1}x^{m-1} + \dots + a_1x + a_0$$

# Lagrange polynomial

**Alternative approach:** let Matlab do the work

For  $m$  data points, we are looking for a polynomial of degree  $m - 1$  or smaller:

$$L(x) = a_{m-1}x^{m-1} + \dots + a_1x + a_0$$

that goes through all the data points *i.e.*

$$a_{m-1}x_1^{m-1} + \dots + a_1x_1 + a_0 = y_1$$

$$a_{m-1}x_2^{m-1} + \dots + a_1x_2 + a_0 = y_2$$

...

$$a_{m-1}x_m^{m-1} + \dots + a_1x_m + a_0 = y_m$$

# Lagrange polynomial

**Alternative approach:** let Matlab do the work

For  $m$  data points, we are looking for a polynomial of degree  $m - 1$  or smaller:

$$L(x) = a_{m-1}x^{m-1} + \dots + a_1x + a_0$$

that goes through all the data points *i.e.*

$$a_{m-1}x_1^{m-1} + \dots + a_1x_1 + a_0 = y_1$$

$$a_{m-1}x_2^{m-1} + \dots + a_1x_2 + a_0 = y_2$$


...

$$a_{m-1}x_m^{m-1} + \dots + a_1x_m + a_0 = y_m$$

We have a system of  $m$  linear algebraic equations with  $m$  unknowns (the coefficients  $a_0, a_1, \dots, a_{m-1}$ ). **We know how to solve such a system using Matlab!**

# Lagrange polynomials: not good with many data points


Lagrange polynomials should generally not be used on a data set that has many points, as the polynomial will likely feature **many “wiggles”**, sometimes of **high magnitude**. For example:






# Cubic splines: introduction

**General idea:** fit cubic polynomials (“splines”), one separate spline between each pair of consecutive points, and make the different splines connect smoothly




# Cubic splines: introduction

**General idea:** fit cubic polynomials (“splines”), one separate spline between each pair of consecutive points, and make the different splines connect smoothly



# Cubic splines: introduction

**General idea:** fit cubic polynomials (“splines”), one separate spline between each pair of consecutive points, and make the different splines connect smoothly



# Cubic splines: introduction

**General idea:** fit cubic polynomials (“splines”), one separate spline between each pair of consecutive points, and make the different splines connect smoothly

Consider a set of  $m$  data points  $(x_i, y_i), i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are in order and different from each other

Group of 2 points	Equation of the spline	Unknowns
$(x_1, y_1), (x_2, y_2)$	$a_1x^3 + b_1x^2 + c_1x + d_1$	$a_1, b_1, c_1, d_1$

# Cubic splines: introduction

**General idea:** fit cubic polynomials (“splines”), one separate spline between each pair of consecutive points, and make the different splines connect smoothly

Consider a set of  $m$  data points  $(x_i, y_i)$ ,  $i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are in order and different from each other

Group of 2 points	Equation of the spline	Unknowns
$(x_1, y_1), (x_2, y_2)$	$a_1x^3 + b_1x^2 + c_1x + d_1$	$a_1, b_1, c_1, d_1$
$(x_2, y_2), (x_3, y_3)$	$a_2x^3 + b_2x^2 + c_2x + d_2$	$a_2, b_2, c_2, d_2$

# Cubic splines: introduction

**General idea:** fit cubic polynomials (“splines”), one separate spline between each pair of consecutive points, and make the different splines connect smoothly

Consider a set of  $m$  data points  $(x_i, y_i)$ ,  $i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are in order and different from each other

Group of 2 points	Equation of the spline	Unknowns
$(x_1, y_1), (x_2, y_2)$	$a_1x^3 + b_1x^2 + c_1x + d_1$	$a_1, b_1, c_1, d_1$
$(x_2, y_2), (x_3, y_3)$	$a_2x^3 + b_2x^2 + c_2x + d_2$	$a_2, b_2, c_2, d_2$
$\dots$	$\dots$	$\dots$

# Cubic splines: introduction

**General idea:** fit cubic polynomials (“splines”), one separate spline between each pair of consecutive points, and make the different splines connect smoothly

Consider a set of  $m$  data points  $(x_i, y_i)$ ,  $i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are in order and different from each other

Group of 2 points	Equation of the spline	Unknowns
$(x_1, y_1), (x_2, y_2)$	$a_1x^3 + b_1x^2 + c_1x + d_1$	$a_1, b_1, c_1, d_1$
$(x_2, y_2), (x_3, y_3)$	$a_2x^3 + b_2x^2 + c_2x + d_2$	$a_2, b_2, c_2, d_2$
$\dots$	$\dots$	$\dots$

**Number of splines:**  $m - 1$

**Number of unknown coefficients:**  $4(m - 1)$

# Cubic splines: introduction

**General idea:** fit cubic polynomials (“splines”), one separate spline between each pair of consecutive points, and make the different splines connect smoothly

Consider a set of  $m$  data points  $(x_i, y_i)$ ,  $i = \{1, 2, \dots, m\}$ , such that all the  $x_i$ 's are in order and different from each other

Group of 2 points	Equation of the spline	Unknowns
$(x_1, y_1), (x_2, y_2)$	$a_1x^3 + b_1x^2 + c_1x + d_1$	$a_1, b_1, c_1, d_1$
$(x_2, y_2), (x_3, y_3)$	$a_2x^3 + b_2x^2 + c_2x + d_2$	$a_2, b_2, c_2, d_2$
$\dots$	$\dots$	$\dots$

**Number of splines:**  $m - 1$

**Number of unknown coefficients:**  $4(m - 1)$

We need to write  $4(m - 1)$  independent equations



## Cubic splines: first set of equations

Obtain the first set of equations by **making each spline go through its respective two data points**:

## Cubic splines: first set of equations

Obtain the first set of equations by **making each spline go through its respective two data points**:

First spline:

$$a_1x_1^3 + b_1x_1^2 + c_1x_1 + d_1 = y_1$$

$$a_1x_2^3 + b_1x_2^2 + c_1x_2 + d_1 = y_2$$

## Cubic splines: first set of equations

Obtain the first set of equations by **making each spline go through its respective two data points**:

First spline:

$$a_1x_1^3 + b_1x_1^2 + c_1x_1 + d_1 = y_1$$

$$a_1x_2^3 + b_1x_2^2 + c_1x_2 + d_1 = y_2$$

Second spline:

$$a_2x_2^3 + b_2x_2^2 + c_2x_2 + d_2 = y_2$$

$$a_2x_3^3 + b_2x_3^2 + c_2x_3 + d_2 = y_3$$

And so on...

## Cubic splines: first set of equations

Obtain the first set of equations by **making each spline go through its respective two data points**:

First spline:

$$a_1x_1^3 + b_1x_1^2 + c_1x_1 + d_1 = y_1$$

$$a_1x_2^3 + b_1x_2^2 + c_1x_2 + d_1 = y_2$$

Second spline:

$$a_2x_2^3 + b_2x_2^2 + c_2x_2 + d_2 = y_2$$

$$a_2x_3^3 + b_2x_3^2 + c_2x_3 + d_2 = y_3$$

And so on...

We can write  $2(m - 1)$  such equations

## Cubic splines: second set of equations

Obtain the second set of equations by **making the splines' first derivatives match at the connection points**:

## Cubic splines: second set of equations

Obtain the second set of equations by **making the splines' first derivatives match at the connection points**:

Connection between the first and second splines ( $x = x_2$ ):

$$3a_1x_2^2 + 2b_1x_2 + c_1 = 3a_2x_2^2 + 2b_2x_2 + c_2$$

## Cubic splines: second set of equations

Obtain the second set of equations by **making the splines' first derivatives match at the connection points**:

Connection between the first and second splines ( $x = x_2$ ):

$$3a_1x_2^2 + 2b_1x_2 + c_1 = 3a_2x_2^2 + 2b_2x_2 + c_2$$

Connection between the second and third splines ( $x = x_3$ ):

$$3a_2x_3^2 + 2b_2x_3 + c_2 = 3a_3x_3^2 + 2b_3x_3 + c_3$$

And so on...

## Cubic splines: second set of equations

Obtain the second set of equations by **making the splines' first derivatives match at the connection points**:

Connection between the first and second splines ( $x = x_2$ ):

$$3a_1x_2^2 + 2b_1x_2 + c_1 = 3a_2x_2^2 + 2b_2x_2 + c_2$$

Connection between the second and third splines ( $x = x_3$ ):

$$3a_2x_3^2 + 2b_2x_3 + c_2 = 3a_3x_3^2 + 2b_3x_3 + c_3$$

And so on...

We can write  $(m - 2)$  such equations  
We have written  $2(m - 1) + (m - 2)$  equations so far



## Cubic splines: third set of equations

Obtain the third set of equations by **making the splines' second derivatives match at the connection points**:

## Cubic splines: third set of equations

Obtain the third set of equations by **making the splines' second derivatives match at the connection points**:

Connection between the first and second splines ( $x = x_2$ ):

$$6a_1x_2 + 2b_1 = 6a_2x_2 + 2b_2$$

## Cubic splines: third set of equations

Obtain the third set of equations by **making the splines' second derivatives match at the connection points**:

Connection between the first and second splines ( $x = x_2$ ):

$$6a_1x_2 + 2b_1 = 6a_2x_2 + 2b_2$$

Connection between the second and third splines ( $x = x_3$ ):

$$6a_2x_3 + 2b_2 = 6a_3x_3 + 2b_3$$

And so on...

## Cubic splines: third set of equations

Obtain the third set of equations by **making the splines' second derivatives match at the connection points**:

Connection between the first and second splines ( $x = x_2$ ):

$$6a_1x_2 + 2b_1 = 6a_2x_2 + 2b_2$$

Connection between the second and third splines ( $x = x_3$ ):

$$6a_2x_3 + 2b_2 = 6a_3x_3 + 2b_3$$

And so on...

We can write  $(m - 2)$  such equations  
We have written  $2(m - 1) + 2(m - 2)$  equations so far

## Cubic splines: last set of equations, and solving the system

We need 2 more equations. These two equations depend on the application. Often, one sets the second derivative of the corresponding splines to be zero at  $x = x_1$  (first point) and  $x = x_m$  (last point):

$$6a_1x_1 + 2b_1 = 0$$

$$6a_{m-1}x_m + 2b_{m-1} = 0$$

## Cubic splines: last set of equations, and solving the system

We need 2 more equations. These two equations depend on the application. Often, one sets the second derivative of the corresponding splines to be zero at  $x = x_1$  (first point) and  $x = x_m$  (last point):

$$6a_1x_1 + 2b_1 = 0$$

$$6a_{m-1}x_m + 2b_{m-1} = 0$$

We have written  $2(m-1) + 2(m-2) + 2 = 4(m-1)$  equations  
We have  $4(m-1)$  unknown coefficients to calculate

## Cubic splines: last set of equations, and solving the system

We need 2 more equations. These two equations depend on the application. Often, one sets the second derivative of the corresponding splines to be zero at  $x = x_1$  (first point) and  $x = x_m$  (last point):

$$6a_1x_1 + 2b_1 = 0$$

$$6a_{m-1}x_m + 2b_{m-1} = 0$$

We have written  $2(m-1) + 2(m-2) + 2 = 4(m-1)$  equations  
We have  $4(m-1)$  unknown coefficients to calculate

Write the system of algebraic linear equations in matrix form, set up the corresponding matrices in Matlab, and “let Matlab do the rest”

# Lagrange polynomial and cubic splines are different

Cubic splines are generally preferable over the Lagrange polynomial when the data set has many points (less “wiggly”). For example:

