

the length of the orbit is

$$L = \int_0^{2\pi} \sqrt{r^2 + (dr/d\theta)^2} d\theta = a(1 - e^2) \int_0^{2\pi} \frac{\sqrt{1 + e^2 + 2e \cos \theta}}{(1 + e \cos \theta)^2} d\theta \approx 3.6 \times 10^8 \text{ km}$$

This seems reasonable, since Mercury's orbit is nearly circular, and the circumference of a circle of radius a is $2\pi a \approx 3.6 \times 10^8 \text{ km}$.

TRUE-FALSE QUIZ

1. False. Consider the curve defined by $x = f(t) = (t - 1)^3$ and $y = g(t) = (t - 1)^2$. Then $g'(t) = 2(t - 1)$, so $g'(1) = 0$, but its graph has a vertical tangent when $t = 1$. Note: The statement is true if $f'(1) \neq 0$ when $g'(1) = 0$.
2. False. If $x = f(t)$ and $y = g(t)$ are twice differentiable, then $\frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{\frac{d}{dt} \left(\frac{dy}{dx} \right)}{\frac{dx}{dt}}$.
3. False. For example, if $f(t) = \cos t$ and $g(t) = \sin t$ for $0 \leq t \leq 4\pi$, then the curve is a circle of radius 1, hence its length is 2π , but $\int_0^{4\pi} \sqrt{[f'(t)]^2 + [g'(t)]^2} dt = \int_0^{4\pi} \sqrt{(-\sin t)^2 + (\cos t)^2} dt = \int_0^{4\pi} 1 dt = 4\pi$, since as t increases from 0 to 4π , the circle is traversed twice.
4. False. If $(r, \theta) = (1, \pi)$, then $(x, y) = (-1, 0)$, so $\tan^{-1}(y/x) = \tan^{-1} 0 = 0 \neq \theta$. The statement is true for points in quadrants I and IV.
5. True. The curve $r = 1 - \sin 2\theta$ is unchanged if we rotate it through 180° about O because $1 - \sin 2(\theta + \pi) = 1 - \sin(2\theta + 2\pi) = 1 - \sin 2\theta$. So it's unchanged if we replace r by $-r$. (See the discussion after Example 8 in Section 10.3.) In other words, it's the same curve as $r = -(1 - \sin 2\theta) = \sin 2\theta - 1$.
6. True. The polar equation $r = 2$, the Cartesian equation $x^2 + y^2 = 4$, and the parametric equations $x = 2 \sin 3t$, $y = 2 \cos 3t$ [$0 \leq t \leq 2\pi$] all describe the circle of radius 2 centered at the origin.
7. False. The first pair of equations gives the portion of the parabola $y = x^2$ with $x \geq 0$, whereas the second pair of equations traces out the whole parabola $y = x^2$.
8. True. $y^2 = 2y + 3x \Leftrightarrow (y - 1)^2 = 3x + 1 = 3(x + \frac{1}{3}) = 4(\frac{3}{4})(x + \frac{1}{3})$, which is the equation of a parabola with vertex $(-\frac{1}{3}, 1)$ and focus $(-\frac{1}{3} + \frac{3}{4}, 1)$, opening to the right.
9. True. By rotating and translating the parabola, we can assume it has an equation of the form $y = cx^2$, where $c > 0$. The tangent at the point (a, ca^2) is the line $y - ca^2 = 2ca(x - a)$; i.e., $y = 2cax - ca^2$. This tangent meets the parabola at the points (x, cx^2) where $cx^2 = 2cax - ca^2$. This equation is equivalent to $x^2 = 2ax - a^2$ [since $c > 0$]. But $x^2 = 2ax - a^2 \Leftrightarrow x^2 - 2ax + a^2 = 0 \Leftrightarrow (x - a)^2 = 0 \Leftrightarrow x = a \Leftrightarrow (x, cx^2) = (a, ca^2)$. This shows that each tangent meets the parabola at exactly one point.

10. True. Consider a hyperbola with focus at the origin, oriented so that its polar equation is $r = \frac{ed}{1 + e \cos \theta}$, where $e > 1$.

The directrix is $x = d$, but along the hyperbola we have $x = r \cos \theta = \frac{ed \cos \theta}{1 + e \cos \theta} = d \left(\frac{e \cos \theta}{1 + e \cos \theta} \right) \neq d$.

10 Review


EXERCISES

1. $x = t^2 + 4t$, $y = 2 - t$, $-4 \leq t \leq 1$. $t = 2 - y$, so

$$x = (2 - y)^2 + 4(2 - y) = 4 - 4y + y^2 + 8 - 4y = y^2 - 8y + 12 \Leftrightarrow$$


$$x + 4 = y^2 - 8y + 16 = (y - 4)^2. \text{ This is part of a parabola with vertex}$$

$(-4, 4)$, opening to the right.




2. $x = 1 + e^{2t}$, $y = e^t$.

$$x = 1 + e^{2t} = 1 + (e^t)^2 = 1 + y^2, y > 0.$$



3. $y = \sec \theta = \frac{1}{\cos \theta} = \frac{1}{x}$. Since $0 \leq \theta \leq \pi/2$, $0 < x \leq 1$ and $y \geq 1$.


This is part of the hyperbola $y = 1/x$.



4. $x = 2 \cos \theta$, $y = 1 + \sin \theta$, $\cos^2 \theta + \sin^2 \theta = 1 \Rightarrow$

$$\left(\frac{x}{2}\right)^2 + (y - 1)^2 = 1 \Rightarrow \frac{x^2}{4} + (y - 1)^2 = 1. \text{ This is an ellipse,}$$

centered at $(0, 1)$, with semimajor axis of length 2 and semiminor axis of length 1.



5. Three different sets of parametric equations for the curve $y = \sqrt{x}$ are



(i) $x = t$, $y = \sqrt{t}$

(ii) $x = t^4$, $y = t^2$

(iii) $x = \tan^2 t$, $y = \tan t$, $0 \leq t < \pi/2$

There are many other sets of equations that also give this curve.


6. For $t < -1$, $x > 0$ and $y < 0$ with x decreasing and y increasing. When $t = -1$, $(x, y) = (0, 0)$. When $-1 < t < 0$, we have $-1 < x < 0$ and $0 < y < 1/2$. When $t = 0$, $(x, y) = (-1, 0)$. When $0 < t < 1$, $-1 < x < 0$ and $-\frac{1}{2} < y < 0$. When $t = 1$, $(x, y) = (0, 0)$ again. When $t > 1$, both x and y are positive and increasing.





The Cartesian coordinates are $x = 4 \cos \frac{2\pi}{3} = 4(-\frac{1}{2}) = -2$ and $y = 4 \sin \frac{2\pi}{3} = 4(\frac{\sqrt{3}}{2}) = 2\sqrt{3}$, that is, the point $(-2, 2\sqrt{3})$.

- (b) Given $x = -3$ and $y = 3$, we have $r = \sqrt{(-3)^2 + 3^2} = \sqrt{18} = 3\sqrt{2}$. Also, $\tan \theta = \frac{y}{x} \Rightarrow \tan \theta = \frac{3}{-3}$, and since $(-3, 3)$ is in the second quadrant, $\theta = \frac{3\pi}{4}$. Thus, one set of polar coordinates for $(-3, 3)$ is $(3\sqrt{2}, \frac{3\pi}{4})$, and two others are $(3\sqrt{2}, \frac{11\pi}{4})$ and $(-3\sqrt{2}, \frac{7\pi}{4})$.



8. $1 \leq r < 2$, $\frac{\pi}{6} \leq \theta \leq \frac{5\pi}{6}$





9. $r = 1 + \sin \theta$. This cardioid is symmetric about the $\theta = \pi/2$ axis.





10. $r = \sin 4\theta$. This is an eight-leaved rose.





11. $r = \cos 3\theta$. This is a three-leaved rose. The curve is traced twice.





12. $r = 3 + \cos 3\theta$. The curve is symmetric about the horizontal axis.




13. $r = 1 + \cos 2\theta$. The curve is symmetric about the pole and both the horizontal and vertical axes.




14. $r = 2 \cos(\theta/2)$. The curve is symmetric about the pole and both the horizontal and vertical axes.



15. $r = \frac{3}{1 + 2 \sin \theta} \Rightarrow e = 2 > 1$, so the conic is a hyperbola. $de = 3 \Rightarrow d = \frac{3}{2}$ and the form “ $+2 \sin \theta$ ” imply that the directrix is above the focus at the origin and has equation $y = \frac{3}{2}$. The vertices are $(1, \frac{\pi}{2})$ and $(-3, \frac{3\pi}{2})$.



16. $r = \frac{3}{2 - 2 \cos \theta} \cdot \frac{1/2}{1/2} = \frac{3/2}{1 - \cos \theta} \Rightarrow e = 1$, so the conic is a parabola. $de = \frac{3}{2} \Rightarrow d = \frac{3}{2}$ and the form “ $-2 \cos \theta$ ” imply that the directrix is to the left of the focus at the origin and has equation $x = -\frac{3}{2}$. The vertex is $(\frac{3}{4}, \pi)$.





$$17. x + y = 2 \Leftrightarrow r \cos \theta + r \sin \theta = 2 \Leftrightarrow r(\cos \theta + \sin \theta) = 2 \Leftrightarrow r = \frac{2}{\cos \theta + \sin \theta}$$

$$18. x^2 + y^2 = 2 \Rightarrow r^2 = 2 \Rightarrow r = \sqrt{2}. [r = -\sqrt{2} \text{ gives the same curve.}]$$

$$19. r = (\sin \theta)/\theta. \text{ As } \theta \rightarrow \pm\infty, r \rightarrow 0.$$


As $\theta \rightarrow 0$, $r \rightarrow 1$. In the first figure, there are an infinite number of x -intercepts at $x = \pi n$, n a nonzero integer. These correspond to pole points in the second figure.



$$20. r = \frac{2}{4 - 3 \cos \theta} = \frac{1/2}{1 - \frac{3}{4} \cos \theta} \Rightarrow e = \frac{3}{4} \text{ and } d = \frac{2}{3}. \text{ The equation of}$$

the directrix is $x = -\frac{2}{3} \Rightarrow r = -2/(3 \cos \theta)$. To obtain the equation of the rotated ellipse, we replace θ in the original equation with $\theta - \frac{2\pi}{3}$,

$$\text{and get } r = \frac{2}{4 - 3 \cos(\theta - \frac{2\pi}{3})}.$$



$$21. x = \ln t, y = 1 + t^2; t = 1. \frac{dy}{dt} = 2t \text{ and } \frac{dx}{dt} = \frac{1}{t}, \text{ so } \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2t}{1/t} = 2t^2.$$

When $t = 1$, $(x, y) = (0, 2)$ and $dy/dx = 2$.

$$22. x = t^3 + 6t + 1, y = 2t - t^2; t = -1. \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2 - 2t}{3t^2 + 6}. \text{ When } t = -1, (x, y) = (-6, -3) \text{ and } \frac{dy}{dx} = \frac{4}{9}.$$

$$23. r = e^{-\theta} \Rightarrow y = r \sin \theta = e^{-\theta} \sin \theta \text{ and } x = r \cos \theta = e^{-\theta} \cos \theta \Rightarrow$$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\frac{dr}{d\theta} \sin \theta + r \cos \theta}{\frac{dr}{d\theta} \cos \theta - r \sin \theta} = \frac{-e^{-\theta} \sin \theta + e^{-\theta} \cos \theta}{-e^{-\theta} \cos \theta - e^{-\theta} \sin \theta} \cdot \frac{-e^{\theta}}{-e^{\theta}} = \frac{\sin \theta - \cos \theta}{\cos \theta + \sin \theta}.$$

$$\text{When } \theta = \pi, \frac{dy}{dx} = \frac{0 - (-1)}{-1 + 0} = \frac{1}{-1} = -1.$$

$$24. r = 3 + \cos 3\theta \Rightarrow \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\frac{dr}{d\theta} \sin \theta + r \cos \theta}{\frac{dr}{d\theta} \cos \theta - r \sin \theta} = \frac{-3 \sin 3\theta \sin \theta + (3 + \cos 3\theta) \cos \theta}{-3 \sin 3\theta \cos \theta - (3 + \cos 3\theta) \sin \theta}.$$

$$\text{When } \theta = \pi/2, \frac{dy}{dx} = \frac{(-3)(-1)(1) + (3 + 0) \cdot 0}{(-3)(-1)(0) - (3 + 0) \cdot 1} = \frac{3}{-3} = -1.$$

$$25. x = t + \sin t, y = t - \cos t \Rightarrow \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{1 + \sin t}{1 + \cos t} \Rightarrow$$

$$\frac{d^2y}{dx^2} = \frac{\frac{d}{dt} \left(\frac{dy}{dx} \right)}{dx/dt} = \frac{\frac{(1 + \cos t) \cos t - (1 + \sin t)(-\sin t)}{(1 + \cos t)^2}}{1 + \cos t} = \frac{\cos t + \cos^2 t + \sin t + \sin^2 t}{(1 + \cos t)^3} = \frac{1 + \cos t + \sin t}{(1 + \cos t)^3}$$


$$26. x = 1 + t^2, y = t - t^3. \frac{dy}{dt} = 1 - 3t^2 \text{ and } \frac{dx}{dt} = 2t, \text{ so } \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{1 - 3t^2}{2t} = \frac{1}{2}t^{-1} - \frac{3}{2}t.$$

$$\frac{d^2y}{dx^2} = \frac{d(dy/dx)/dt}{dx/dt} = \frac{-\frac{1}{2}t^{-2} - \frac{3}{2}}{2t} = -\frac{1}{4}t^{-3} - \frac{3}{4}t^{-1} = -\frac{1}{4t^3}(1 + 3t^2) = -\frac{3t^2 + 1}{4t^3}.$$

27. We graph the curve $x = t^3 - 3t$, $y = t^2 + t + 1$ for $-2.2 \leq t \leq 1.2$.

By zooming in or using a cursor, we find that the lowest point is about $(1.4, 0.75)$. To find the exact values, we find the t -value at which

$$dy/dt = 2t + 1 = 0 \Leftrightarrow t = -\frac{1}{2} \Leftrightarrow (x, y) = \left(\frac{11}{8}, \frac{3}{4}\right).$$



28. We estimate the coordinates of the point of intersection to be $(-2, 3)$. In fact this is exact, since both $t = -2$ and $t = 1$ give the point $(-2, 3)$. So the area enclosed by the loop is

$$\begin{aligned} \int_{t=-2}^{t=1} y \, dx &= \int_{-2}^1 (t^2 + t + 1)(3t^2 - 3) \, dt = \int_{-2}^1 (3t^4 + 3t^3 - 3t - 3) \, dt \\ &= \left[\frac{3}{5}t^5 + \frac{3}{4}t^4 - \frac{3}{2}t^2 - 3t \right]_{-2}^1 = \left(\frac{3}{5} + \frac{3}{4} - \frac{3}{2} - 3 \right) - \left(-\frac{96}{5} + 12 - 6 - (-6) \right) = \frac{81}{20} \end{aligned}$$

29. $x = 2a \cos t - a \cos 2t \Rightarrow \frac{dx}{dt} = -2a \sin t + 2a \sin 2t = 2a \sin t(2 \cos t - 1) = 0 \Leftrightarrow$


$$\sin t = 0 \text{ or } \cos t = \frac{1}{2} \Rightarrow t = 0, \frac{\pi}{3}, \pi, \text{ or } \frac{5\pi}{3}.$$

$$y = 2a \sin t - a \sin 2t \Rightarrow \frac{dy}{dt} = 2a \cos t - 2a \cos 2t = 2a(1 + \cos t - 2 \cos^2 t) = 2a(1 - \cos t)(1 + 2 \cos t) = 0 \Rightarrow$$

$$t = 0, \frac{2\pi}{3}, \text{ or } \frac{4\pi}{3}.$$

Thus the graph has vertical tangents where $t = \frac{\pi}{3}, \pi$ and $\frac{5\pi}{3}$, and horizontal tangents where $t = \frac{2\pi}{3}$ and $\frac{4\pi}{3}$. To determine what the slope is where $t = 0$, we use l'Hospital's Rule to evaluate $\lim_{t \rightarrow 0} \frac{dy/dt}{dx/dt} = 0$, so there is a horizontal tangent there.

t	x	y
0	a	0
$\frac{\pi}{3}$	$\frac{3}{2}a$	$\frac{\sqrt{3}}{2}a$
$\frac{2\pi}{3}$	$-\frac{1}{2}a$	$\frac{3\sqrt{3}}{2}a$
π	$-3a$	0
$\frac{4\pi}{3}$	$-\frac{1}{2}a$	$-\frac{3\sqrt{3}}{2}a$
$\frac{5\pi}{3}$	$\frac{3}{2}a$	$-\frac{\sqrt{3}}{2}a$



30. From Exercise 29, $x = 2a \cos t - a \cos 2t$, $y = 2a \sin t - a \sin 2t \Rightarrow$

$$\begin{aligned} A &= 2 \int_{\pi}^0 (2a \sin t - a \sin 2t)(-2a \sin t + 2a \sin 2t) \, dt = 4a^2 \int_0^{\pi} (2 \sin^2 t + \sin^2 2t - 3 \sin t \sin 2t) \, dt \\ &= 4a^2 \int_0^{\pi} \left[(1 - \cos 2t) + \frac{1}{2}(1 - \cos 4t) - 6 \sin^2 t \cos t \right] \, dt = 4a^2 \left[t - \frac{1}{2} \sin 2t + \frac{1}{2} t - \frac{1}{8} \sin 4t - 2 \sin^3 t \right]_0^{\pi} \\ &= 4a^2 \left(\frac{3}{2} \right) \pi = 6\pi a^2 \end{aligned}$$


31. The curve $r^2 = 9 \cos 5\theta$ has 10 “petals.” For instance, for $-\frac{\pi}{10} \leq \theta \leq \frac{\pi}{10}$, there are two petals, one with $r > 0$ and one with $r < 0$.

$$A = 10 \int_{-\pi/10}^{\pi/10} \frac{1}{2} r^2 \, d\theta = 5 \int_{-\pi/10}^{\pi/10} 9 \cos 5\theta \, d\theta = 5 \cdot 9 \cdot 2 \int_0^{\pi/10} \cos 5\theta \, d\theta = 18 [\sin 5\theta]_0^{\pi/10} = 18$$

32. $r = 1 - 3 \sin \theta$. The inner loop is traced out as θ goes from $\alpha = \sin^{-1}(\frac{1}{3})$ to $\pi - \alpha$, so

$$\begin{aligned} A &= \int_{\alpha}^{\pi-\alpha} \frac{1}{2} r^2 \, d\theta = \int_{\alpha}^{\pi-\alpha} (1 - 3 \sin \theta)^2 \, d\theta = \int_{\alpha}^{\pi-\alpha} \left[1 - 6 \sin \theta + \frac{9}{2}(1 - \cos 2\theta) \right] \, d\theta \\ &= \left[\frac{11}{2} \theta + 6 \cos \theta - \frac{9}{4} \sin 2\theta \right]_{\alpha}^{\pi-\alpha} = \frac{11}{4} \pi - \frac{11}{2} \sin^{-1} \left(\frac{1}{3} \right) - 3\sqrt{2} \end{aligned}$$


33. The curves intersect when $4 \cos \theta = 2 \Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \pm \frac{\pi}{3}$
for $-\pi \leq \theta \leq \pi$. The points of intersection are $(2, \frac{\pi}{3})$ and $(2, -\frac{\pi}{3})$.




34. The two curves clearly both contain the pole. For other points of intersection, $\cot \theta = 2 \cos(\theta + 2n\pi)$ or $-2 \cos(\theta + \pi + 2n\pi)$, both of which reduce to $\cot \theta = 2 \cos \theta \Leftrightarrow \cos \theta = 2 \sin \theta \cos \theta \Leftrightarrow \cos \theta(1 - 2 \sin \theta) = 0 \Rightarrow \cos \theta = 0$ or $\sin \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}$ or $\frac{3\pi}{2} \Rightarrow$ intersection points are $(0, \frac{\pi}{2})$, $(\sqrt{3}, \frac{\pi}{6})$, and $(\sqrt{3}, \frac{11\pi}{6})$.

35. The curves intersect where $2 \sin \theta = \sin \theta + \cos \theta \Rightarrow \sin \theta = \cos \theta \Rightarrow \theta = \frac{\pi}{4}$, and also at the origin (at which $\theta = \frac{3\pi}{4}$ on the second curve).

$$\begin{aligned} A &= \int_0^{\pi/4} \frac{1}{2} (2 \sin \theta)^2 d\theta + \int_{\pi/4}^{3\pi/4} \frac{1}{2} (\sin \theta + \cos \theta)^2 d\theta \\ &= \int_0^{\pi/4} (1 - \cos 2\theta) d\theta + \frac{1}{2} \int_{\pi/4}^{3\pi/4} (1 + \sin 2\theta) d\theta \\ &= \left[\theta - \frac{1}{2} \sin 2\theta \right]_0^{\pi/4} + \left[\frac{1}{2} \theta - \frac{1}{4} \cos 2\theta \right]_{\pi/4}^{3\pi/4} = \frac{1}{2}(\pi - 1) \end{aligned}$$



36. $A = 2 \int_{-\pi/2}^{\pi/6} \frac{1}{2} [(2 + \cos 2\theta)^2 - (2 + \sin \theta)^2] d\theta$
 $= \int_{-\pi/2}^{\pi/6} [4 \cos 2\theta + \cos^2 2\theta - 4 \sin \theta - \sin^2 \theta] d\theta$
 $= [2 \sin 2\theta + \frac{1}{2} \theta + \frac{1}{8} \sin 4\theta + 4 \cos \theta - \frac{1}{2} \theta + \frac{1}{4} \sin 2\theta]_{-\pi/2}^{\pi/6}$
 $= \frac{51}{16} \sqrt{3}$



37. $x = 3t^2$, $y = 2t^3$.

$$\begin{aligned} L &= \int_0^2 \sqrt{(dx/dt)^2 + (dy/dt)^2} dt = \int_0^2 \sqrt{(6t)^2 + (6t^2)^2} dt = \int_0^2 \sqrt{36t^2 + 36t^4} dt = \int_0^2 \sqrt{36t^2} \sqrt{1 + t^2} dt \\ &= \int_0^2 6|t| \sqrt{1 + t^2} dt = 6 \int_0^2 t \sqrt{1 + t^2} dt = 6 \int_1^5 u^{1/2} (\frac{1}{2} du) \quad [u = 1 + t^2, du = 2t dt] \\ &= 6 \cdot \frac{1}{2} \cdot \frac{2}{3} [u^{3/2}]_1^5 = 2(5^{3/2} - 1) = 2(5\sqrt{5} - 1) \end{aligned}$$

38. $x = 2 + 3t$, $y = \cosh 3t \Rightarrow (dx/dt)^2 + (dy/dt)^2 = 3^2 + (3 \sinh 3t)^2 = 9(1 + \sinh^2 3t) = 9 \cosh^2 3t$, so
 $L = \int_0^1 \sqrt{9 \cosh^2 3t} dt = \int_0^1 |3 \cosh 3t| dt = \int_0^1 3 \cosh 3t dt = [\sinh 3t]_0^1 = \sinh 3 - \sinh 0 = \sinh 3$.

39. $L = \int_{\pi}^{2\pi} \sqrt{r^2 + (dr/d\theta)^2} d\theta = \int_{\pi}^{2\pi} \sqrt{(1/\theta)^2 + (-1/\theta^2)^2} d\theta = \int_{\pi}^{2\pi} \frac{\sqrt{\theta^2 + 1}}{\theta^2} d\theta$
 $\stackrel{24}{=} \left[-\frac{\sqrt{\theta^2 + 1}}{\theta} + \ln(\theta + \sqrt{\theta^2 + 1}) \right]_{\pi}^{2\pi} = \frac{\sqrt{\pi^2 + 1}}{\pi} - \frac{\sqrt{4\pi^2 + 1}}{2\pi} + \ln\left(\frac{2\pi + \sqrt{4\pi^2 + 1}}{\pi + \sqrt{\pi^2 + 1}}\right)$
 $= \frac{2\sqrt{\pi^2 + 1} - \sqrt{4\pi^2 + 1}}{2\pi} + \ln\left(\frac{2\pi + \sqrt{4\pi^2 + 1}}{\pi + \sqrt{\pi^2 + 1}}\right)$

$$40. L = \int_0^\pi \sqrt{r^2 + (dr/d\theta)^2} d\theta = \int_0^\pi \sqrt{\sin^6(\frac{1}{3}\theta) + \sin^4(\frac{1}{3}\theta) \cos^2(\frac{1}{3}\theta)} d\theta$$

$$= \int_0^\pi \sin^2(\frac{1}{3}\theta) d\theta = [\frac{1}{2}(\theta - \frac{3}{2} \sin(\frac{2}{3}\theta))]_0^\pi = \frac{1}{2}\pi - \frac{3}{8}\sqrt{3}$$

$$41. x = 4\sqrt{t}, y = \frac{t^3}{3} + \frac{1}{2t^2}, 1 \leq t \leq 4 \Rightarrow$$

$$S = \int_1^4 2\pi y \sqrt{(dx/dt)^2 + (dy/dt)^2} dt = \int_1^4 2\pi (\frac{1}{3}t^3 + \frac{1}{2}t^{-2}) \sqrt{(2/\sqrt{t})^2 + (t^2 - t^{-3})^2} dt$$


$$= 2\pi \int_1^4 (\frac{1}{3}t^3 + \frac{1}{2}t^{-2}) \sqrt{(t^2 + t^{-3})^2} dt = 2\pi \int_1^4 (\frac{1}{3}t^5 + \frac{5}{6} + \frac{1}{2}t^{-5}) dt = 2\pi [\frac{1}{18}t^6 + \frac{5}{6}t - \frac{1}{8}t^{-4}]_1^4 = \frac{471,295}{1024}\pi$$

$$42. x = 2 + 3t, y = \cosh 3t \Rightarrow (dx/dt)^2 + (dy/dt)^2 = 3^2 + (3 \sinh 3t)^2 = 9(1 + \sinh^2 3t) = 9 \cosh^2 3t, \text{ so}$$







$$S = \int_0^1 2\pi y ds = \int_0^1 2\pi \cosh 3t \sqrt{9 \cosh^2 3t} dt = \int_0^1 2\pi \cosh 3t |3 \cosh 3t| dt = \int_0^1 2\pi \cosh 3t \cdot 3 \cosh 3t dt$$

$$= 6\pi \int_0^1 \cosh^2 3t dt = 6\pi \int_0^1 \frac{1}{2}(1 + \cosh 6t) dt = 3\pi [t + \frac{1}{6} \sinh 6t]_0^1 = 3\pi(1 + \frac{1}{6} \sinh 6) = 3\pi + \frac{\pi}{2} \sinh 6$$

43. For all c except -1 , the curve is asymptotic to the line $x = 1$. For $c < -1$, the curve bulges to the right near $y = 0$. As c increases, the bulge becomes smaller, until at $c = -1$ the curve is the straight line $x = 1$. As c continues to increase, the curve bulges to the left, until at $c = 0$ there is a cusp at the origin. For $c > 0$, there is a loop to the left of the origin, whose size and roundness increase as c increases. Note that the x -intercept of the curve is always $-c$.




44. For a close to 0, the graph of $r^a = |\sin 2\theta|$ consists of four thin petals. As a increases, the petals get wider, until as $a \rightarrow \infty$, each petal occupies almost its entire quarter-circle.

 $a = 0.01$  $a = 0.1$  $a = 1$  $a = 5$  $a = 10$  $a = 25$

45. $\frac{x^2}{9} + \frac{y^2}{8} = 1$ is an ellipse with center $(0, 0)$.

$$a = 3, b = 2\sqrt{2}, c = 1 \Rightarrow$$

foci $(\pm 1, 0)$, vertices $(\pm 3, 0)$.




46. $4x^2 - y^2 = 16 \Leftrightarrow \frac{x^2}{4} - \frac{y^2}{16} = 1$ is a hyperbola

with center $(0, 0)$, vertices $(\pm 2, 0)$, $a = 2$, $b = 4$,

$c = \sqrt{16 + 4} = 2\sqrt{5}$, foci $(\pm 2\sqrt{5}, 0)$ and

asymptotes $y = \pm 2x$.




47. $6y^2 + x - 36y + 55 = 0 \Leftrightarrow$

$$6(y^2 - 6y + 9) = -(x + 1) \Leftrightarrow$$

$(y - 3)^2 = -\frac{1}{6}(x + 1)$, a parabola with vertex $(-1, 3)$,

opening to the left, $p = -\frac{1}{24} \Rightarrow$ focus $(-\frac{25}{24}, 3)$ and

directrix $x = -\frac{23}{24}$.



48. $25x^2 + 4y^2 + 50x - 16y = 59 \Leftrightarrow$


$$25(x + 1)^2 + 4(y - 2)^2 = 100 \Leftrightarrow$$

$\frac{1}{4}(x + 1)^2 + \frac{1}{25}(y - 2)^2 = 1$ is an ellipse centered at

$(-1, 2)$ with foci on the line $x = -1$, vertices $(-1, 7)$

and $(-1, -3)$; $a = 5$, $b = 2 \Rightarrow c = \sqrt{21} \Rightarrow$

foci $(-1, 2 \pm \sqrt{21})$.



49. The ellipse with foci $(\pm 4, 0)$ and vertices $(\pm 5, 0)$ has center $(0, 0)$ and a horizontal major axis, with $a = 5$ and $c = 4$,

so $b^2 = a^2 - c^2 = 5^2 - 4^2 = 9$. An equation is $\frac{x^2}{25} + \frac{y^2}{9} = 1$.

50. The distance from the focus $(2, 1)$ to the directrix $x = -4$ is $2 - (-4) = 6$, so the distance from the focus to the vertex is $\frac{1}{2}(6) = 3$ and the vertex is $(-1, 1)$. Since the focus is to the right of the vertex, $p = 3$. An equation is

$$(y - 1)^2 = 4 \cdot 3[x - (-1)], \text{ or } (y - 1)^2 = 12(x + 1).$$

51. The center of a hyperbola with foci $(0, \pm 4)$ is $(0, 0)$, so $c = 4$ and an equation is $\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$.

The asymptote $y = 3x$ has slope 3, so $\frac{a}{b} = \frac{3}{1} \Rightarrow a = 3b$ and $a^2 + b^2 = c^2 \Rightarrow (3b)^2 + b^2 = 4^2 \Rightarrow$

$10b^2 = 16 \Rightarrow b^2 = \frac{8}{5}$ and so $a^2 = 16 - \frac{8}{5} = \frac{72}{5}$. Thus, an equation is $\frac{y^2}{72/5} - \frac{x^2}{8/5} = 1$, or $\frac{5y^2}{72} - \frac{5x^2}{8} = 1$.

52. Center is $(3, 0)$, and $a = \frac{8}{2} = 4$, $c = 2 \Leftrightarrow b = \sqrt{4^2 - 2^2} = \sqrt{12} \Rightarrow$

an equation of the ellipse is $\frac{(x - 3)^2}{12} + \frac{y^2}{16} = 1$.

53. $x^2 + y = 100 \Leftrightarrow x^2 = -(y - 100)$ has its vertex at $(0, 100)$, so one of the vertices of the ellipse is $(0, 100)$. Another form of the equation of a parabola is $x^2 = 4p(y - 100)$ so $4p(y - 100) = -(y - 100) \Rightarrow 4p = -1 \Rightarrow p = -\frac{1}{4}$. Therefore the shared focus is found at $(0, \frac{399}{4})$ so $2c = \frac{399}{4} - 0 \Rightarrow c = \frac{399}{8}$ and the center of the ellipse is $(0, \frac{399}{8})$. So $a = 100 - \frac{399}{8} = \frac{401}{8}$ and $b^2 = a^2 - c^2 = \frac{401^2 - 399^2}{8^2} = 25$. So the equation of the ellipse is $\frac{x^2}{b^2} + \frac{(y - \frac{399}{8})^2}{a^2} = 1 \Rightarrow \frac{x^2}{25} + \frac{(y - \frac{399}{8})^2}{(\frac{401}{8})^2} = 1$, or $\frac{x^2}{25} + \frac{(8y - 399)^2}{160,801} = 1$.
54. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \Rightarrow \frac{2x}{a^2} + \frac{2y}{b^2} \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{b^2}{a^2} \frac{x}{y}$. Therefore $\frac{dy}{dx} = m \Leftrightarrow y = -\frac{b^2}{a^2} \frac{x}{m}$. Combining this condition with $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, we find that $x = \pm \frac{a^2 m}{\sqrt{a^2 m^2 + b^2}}$. In other words, the two points on the ellipse where the tangent has slope m are $(\pm \frac{a^2 m}{\sqrt{a^2 m^2 + b^2}}, \mp \frac{b^2}{\sqrt{a^2 m^2 + b^2}})$. The tangent lines at these points have the equations $y \pm \frac{b^2}{\sqrt{a^2 m^2 + b^2}} = m(x \mp \frac{a^2 m}{\sqrt{a^2 m^2 + b^2}})$ or $y = mx \mp \frac{a^2 m^2}{\sqrt{a^2 m^2 + b^2}} \mp \frac{b^2}{\sqrt{a^2 m^2 + b^2}} = mx \mp \sqrt{a^2 m^2 + b^2}$.
55. Directrix $x = 4 \Rightarrow d = 4$, so $e = \frac{1}{3} \Rightarrow r = \frac{ed}{1 + e \cos \theta} = \frac{4}{3 + \cos \theta}$.
56. See the end of the proof of Theorem 10.6.1. If $e > 1$, then $1 - e^2 < 0$ and Equations 10.6.4 become $a^2 = \frac{e^2 d^2}{(e^2 - 1)^2}$ and $b^2 = \frac{e^2 d^2}{e^2 - 1}$, so $\frac{b^2}{a^2} = e^2 - 1$. The asymptotes $y = \pm \frac{b}{a} x$ have slopes $\pm \frac{b}{a} = \pm \sqrt{e^2 - 1}$, so the angles they make with the polar axis are $\pm \tan^{-1}[\sqrt{e^2 - 1}] = \cos^{-1}(\pm 1/e)$.
57. In polar coordinates, an equation for the circle is $r = 2a \sin \theta$. Thus, the coordinates of Q are $x = r \cos \theta = 2a \sin \theta \cos \theta$ and $y = r \sin \theta = 2a \sin^2 \theta$. The coordinates of R are $x = 2a \cot \theta$ and $y = 2a$. Since P is the midpoint of QR , we use the midpoint formula to get $x = a(\sin \theta \cos \theta + \cot \theta)$ and $y = a(1 + \sin^2 \theta)$.
58. (a) If (a, b) lies on the curve, then there is some parameter value t_1 such that $\frac{3t_1}{1 + t_1^3} = a$ and $\frac{3t_1^2}{1 + t_1^3} = b$. If $t_1 = 0$, the point is $(0, 0)$, which lies on the line $y = x$. If $t_1 \neq 0$, then the point corresponding to $t = \frac{1}{t_1}$ is given by $x = \frac{3(1/t_1)}{1 + (1/t_1)^3} = \frac{3t_1^2}{t_1^3 + 1} = b$, $y = \frac{3(1/t_1)^2}{1 + (1/t_1)^3} = \frac{3t_1}{t_1^3 + 1} = a$. So (b, a) also lies on the curve. [Another way to see this is to do part (e) first; the result is immediate.] The curve intersects the line $y = x$ when $\frac{3t}{1 + t^3} = \frac{3t^2}{1 + t^3} \Rightarrow t = t^2 \Rightarrow t = 0$ or 1 , so the points are $(0, 0)$ and $(\frac{3}{2}, \frac{3}{2})$.

- (b) $\frac{dy}{dt} = \frac{(1+t^3)(6t) - 3t^2(3t^2)}{(1+t^3)^2} = \frac{6t - 3t^4}{(1+t^3)^2} = 0$ when $6t - 3t^4 = 3t(2 - t^3) = 0 \Rightarrow t = 0$ or $t = \sqrt[3]{2}$, so there are horizontal tangents at $(0, 0)$ and $(\sqrt[3]{2}, \sqrt[3]{4})$. Using the symmetry from part (a), we see that there are vertical tangents at $(0, 0)$ and $(\sqrt[3]{4}, \sqrt[3]{2})$.

- (c) Notice that as $t \rightarrow -1^+$, we have $x \rightarrow -\infty$ and $y \rightarrow \infty$. As $t \rightarrow -1^-$, we have $x \rightarrow \infty$ and $y \rightarrow -\infty$. Also

$$y - (-x - 1) = y + x + 1 = \frac{3t + 3t^2 + (1+t^3)}{1+t^3} = \frac{(t+1)^3}{t^2 - t + 1} \rightarrow 0 \text{ as } t \rightarrow -1. \text{ So } y = -x - 1 \text{ is a slant asymptote.}$$


- (d) $\frac{dx}{dt} = \frac{(1+t^3)(3) - 3t(3t^2)}{(1+t^3)^2} = \frac{3-6t^3}{(1+t^3)^2}$ and from part (b) we have $\frac{dy}{dt} = \frac{6t-3t^4}{(1+t^3)^2}$. So $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{t(2-t^3)}{1-2t^3}$.

$$\text{Also } \frac{d^2y}{dx^2} = \frac{\frac{d}{dt}\left(\frac{dy}{dx}\right)}{dx/dt} = \frac{2(1+t^3)^4}{3(1-2t^3)^3} > 0 \Leftrightarrow t < \frac{1}{\sqrt[3]{2}}.$$

So the curve is concave upward there and has a minimum point at $(0, 0)$

and a maximum point at $(\sqrt[3]{2}, \sqrt[3]{4})$. Using this together with the

information from parts (a), (b), and (c), we sketch the curve.



- (e) $x^3 + y^3 = \left(\frac{3t}{1+t^3}\right)^3 + \left(\frac{3t^2}{1+t^3}\right)^3 = \frac{27t^3 + 27t^6}{(1+t^3)^3} = \frac{27t^3(1+t^3)}{(1+t^3)^3} = \frac{27t^3}{(1+t^3)^2}$

$$\text{and } 3xy = 3\left(\frac{3t}{1+t^3}\right)\left(\frac{3t^2}{1+t^3}\right) = \frac{27t^3}{(1+t^3)^2}, \text{ so } x^3 + y^3 = 3xy.$$

- (f) We start with the equation from part (e) and substitute $x = r \cos \theta$, $y = r \sin \theta$. Then $x^3 + y^3 = 3xy \Rightarrow$

$$r^3 \cos^3 \theta + r^3 \sin^3 \theta = 3r^2 \cos \theta \sin \theta. \text{ For } r \neq 0, \text{ this gives } r = \frac{3 \cos \theta \sin \theta}{\cos^3 \theta + \sin^3 \theta}. \text{ Dividing numerator and denominator}$$

$$\text{by } \cos^3 \theta, \text{ we obtain } r = \frac{3\left(\frac{1}{\cos \theta}\right) \frac{\sin \theta}{\cos \theta}}{1 + \frac{\sin^3 \theta}{\cos^3 \theta}} = \frac{3 \sec \theta \tan \theta}{1 + \tan^3 \theta}.$$

- (g) The loop corresponds to $\theta \in (0, \frac{\pi}{2})$, so its area is

$$\begin{aligned} A &= \int_0^{\pi/2} \frac{r^2}{2} d\theta = \frac{1}{2} \int_0^{\pi/2} \left(\frac{3 \sec \theta \tan \theta}{1 + \tan^3 \theta} \right)^2 d\theta = \frac{9}{2} \int_0^{\pi/2} \frac{\sec^2 \theta \tan^2 \theta}{(1 + \tan^3 \theta)^2} d\theta = \frac{9}{2} \int_0^{\infty} \frac{u^2 du}{(1 + u^3)^2} \quad [\text{let } u = \tan \theta] \\ &= \lim_{b \rightarrow \infty} \frac{9}{2} \left[-\frac{1}{3} (1 + u^3)^{-1} \right]_0^b = \frac{3}{2} \end{aligned}$$

- (h) By symmetry, the area between the folium and the line $y = -x - 1$ is equal to the enclosed area in the third quadrant, plus twice the enclosed area in the fourth quadrant. The area in the third quadrant is $\frac{1}{2}$, and since $y = -x - 1 \Rightarrow$

$$r \sin \theta = -r \cos \theta - 1 \Rightarrow r = -\frac{1}{\sin \theta + \cos \theta}, \text{ the area in the fourth quadrant is}$$

$$\frac{1}{2} \int_{-\pi/2}^{-\pi/4} \left[\left(-\frac{1}{\sin \theta + \cos \theta} \right)^2 - \left(\frac{3 \sec \theta \tan \theta}{1 + \tan^3 \theta} \right)^2 \right] d\theta \stackrel{\text{CAS}}{=} \frac{1}{2}. \text{ Therefore, the total area is } \frac{1}{2} + 2\left(\frac{1}{2}\right) = \frac{3}{2}.$$