George M. Bergman Bechtel Auditorium

Final Examination

14 May, 2001 8:00-11:00 AM

- 1. (60 points, 6 points apiece) Find the following. If an expression is undefined, say so.
- (a) The area of the region of the plane described in polar coordinates by the conditions $0 \le \theta \le 1$, $0 \le r \le 1 + e^{\theta}$.
- (b) A unit vector perpendicular to both $\langle 1, 2, 3 \rangle$ and $\langle 4, 5, 6 \rangle$.
- The length of the curve given by $x = t^2$, $y = (t^3/3) t$, where $-1 \le t \le 1$.
- $\frac{d}{dt} f(g(t^2), g(t^3))$, where f is a differentiable function of two variables and g is a differentiable function of one variable. The answer should be expressed in terms of f, g, and their derivatives and/or partial derivatives.
 - (e) $\int_0^1 \int_{1-x}^{1+x} xy \, dy \, dx$.
 - (f) An expression for $\iiint_E f(x, y, z) dV$, as an iterated integral, where $E = \{(x, y, z) \mid x^2 + y^2 + z^2 \le 100\}$, and f is a continuous function. (Do not change coordinates.)
 - (g) $\int_C \mathbf{F} \cdot d\mathbf{r}$, where \mathbf{F} is the vector field $\langle 1, 2, y \rangle$ and C is the curve given by $\mathbf{r}(t) = \langle t^2, t^3, t^5 \rangle$ for $0 \le t \le 10$.
 - (h) An expression for $\iint_{D_1} f(x, y) dx dy$ as an integral over D_2 , where D_1 is a region of the x-y-plane, and D_2 is a region of the u-v-plane which is mapped in a one-to-one fashion onto D_1 by the transformation x = uv, $y = u^2/v$; and where f is a continuous function on D_1 .
 - (i) $\int_C xy^{-1}dx$, where C is the segment of the curve $y = x^3$ between x = -1 and x = 1
 - (j) $\iint_{S} \mathbf{F} \cdot d\mathbf{S}$, where \mathbf{F} is the constant vector field $\langle 2, 1, -1 \rangle$, and S is the parallelogram with vertices (1,0,0), (0,2,0), (0,0,3) and (1,-2,3), and upward orientation.

- 2. (12 points; 4 points each) In each part, give the definition asked for. Note that you are **not** asked to give examples or other related information.
- (a) Define the partial derivative $f_1(a, b)$ of a function f at a point (a, b) of its domain. (There are many other symbols for this, e.g., $(\partial f/\partial x)(a, b)$, but we are asking for a definition, not alternative notation. A brief general statement of how one would find this partial derivative would be an acceptable answer.)
- (b) What does it mean to say that a function f of two variables is differentiable?
- (c) If f is a function of two variables, and \mathbf{u} a unit vector in the plane, what is meant by the directional derivative $D_{\mathbf{u}} f$?
- (12 points) (a) (6 points) Find constants a and b such that the vector field $(3x^2y \sin y 2x^2\cos y, ax^3y\cos y + bx^3\sin y)$ is the gradient of a function f(x, y). (You do not have to find the function f.)
 - (b) (6 points) Suppose a and b are as in part (a), so that \mathbf{F} is the gradient of a function. Prove that $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$ for every curve C beginning at (0,0) and ending at (0,1). (Suggestion: first show this fact for one particular curve. You do not have to have done part (a) to do part (b).) 另一种方法是?
- (8 points) Let S_1 be the hemisphere $z = \sqrt{(1-x^2-y^2)}$ and S_2 the hemisphere $z = -\sqrt{(1-x^2-y^2)}$, both with upward orientation, and let E be the solid ball $\{(x,y,z) \mid x^2+y^2+z^2 \le 1\}$. If F is any vector field whose component functions have continuous partial derivatives on an open set containing E, write an equation expressing the triple integral over E of the divergence of F in terms of surface integrals over S_1 and S_2 . 表达式是?
 - (8 points) Let \mathbb{F} be the vector field $\langle xy, yz^2, zx^3 \rangle + \nabla e^{x+\sin y}$, and let S be the part of the surface $z = xy^2(1-x-y)^3$ lying above the triangle with vertices (0,0), (1,0), (0,1), with upward orientation. Compute $\iint_S \text{curl } \mathbb{F} \cdot d\mathbb{S}$. (Suggestion: Treat the two summands of \mathbb{F} separately.)