


Principal Component Analysis


Question 1

Consider the following 2D dataset:



Which of the following figures correspond to possible values that PCA may return for $u^{(1)}$ (the first eigenvector / first principal component)? Check all that apply (you may have to check more than one figure).



The maximal variance is along the $y = x$ line, so this option is correct.



The maximal variance is along the $y = x$ line, so the negative vector along that line is correct for the first principal component.



Question 2

Which of the following is a reasonable way to select the number of principal components k ?

(Recall that n is the dimensionality of the input data and m is the number of input examples.)

☐ Choose the value of k that minimizes the approximation error $\frac{1}{m} \sum_{i=1}^m ||x^{(i)} - x_{\text{approx}}^{(i)}||^2$.

☒ **Choose k to be the smallest value so that at least 99% of the variance is retained.**

This is correct, as it maintains the structure of the data while maximally reducing its dimension.

☐ Choose k to be the smallest value so that at least 1% of the variance is retained.

☐ Choose k to be 99% of n (i.e., $k = 0.99 * n$, rounded to the nearest integer).

Question 3

Suppose someone tells you that they ran PCA in such a way that "95% of the variance was retained." What is an equivalent statement to this?

- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2} \geq 0.05$
- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}\|^2} \geq 0.95$
- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2} \geq 0.95$
- ☒ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2} \leq 0.05$
-

Question 4

Which of the following statements are true? Check all that apply.

- ☒ **Given an input $x \in \mathbb{R}^n$, PCA compresses it to a lower-dimensional $z \in \mathbb{R}^k$.**
- PCA compresses it to a lower dimensional vector by projecting it onto the learned principal components.
- ☐ Feature scaling is not useful for PCA, since the eigenvector calculation (such as using Octave's `svd(Sigma)` routine) takes care of this automatically.
- ☐ PCA can be used only to reduce the dimensionality of data by 1 (such as 3D to 2D, or 2D to 1D).
- ☒ **If the input features are on very different scales, it is a good idea to perform feature scaling before applying PCA.**
- Feature scaling prevents one feature dimension from becoming a strong principal component only because of the large magnitude of the feature values (as opposed to large variance on that dimension).
-

Question 5

Which of the following are recommended applications of PCA? Select all that apply.

- ☒ **Data compression: Reduce the dimension of your data, so that it takes up less memory / disk space.**
- If memory or disk space is limited, PCA allows you to save space in exchange for losing a little of the data's information. This can be a reasonable tradeoff.
- ☒ **Data visualization: Reduce data to 2D (or 3D) so that it can be plotted.**
- This is a good use of PCA, as it can give you intuition about your data that would otherwise be impossible to see.
- ☐ Preventing overfitting: Reduce the number of features (in a supervised learning problem), so that there are fewer parameters to learn.

☐ To get more features to feed into a learning algorithm.