

Math 3GR3 - Abstract Algebra

Sang Woo Park

November 9, 2017

Course Outline

- Office hours: Monday 9:30-10:20 and Wednesday 2:30-3:20 (HH 419)

Contents

1	Set theory	3
1.1	Reveiew	3
1.2	Equivalence relation	4
1.3	Well ordering principle and division algorithm	7
2	Groups and rings	12
2.1	Group theory	12
2.2	Subgroups	17
3	Special groups	19
3.1	Cyclic groups	19
3.2	Permutation groups	25
3.3	Alternating groups	30
3.4	Group of rigid motions	31
3.5	Lagrange's Theorem	33
4	Fermat's little theorem	38
4.1	Fermat's little theorem	38
5	Isomorphisms	39
5.1	Isomorphisms	39
5.2	Cyclic groups	39
5.3	Cayley's Theorem	41
5.4	Direct Products	43
6	Factor groups	46
6.1	Factor groups	46
6.2	Simple groups	50

7	Homomorphism	52
7.1	Homomorphism	52
7.2	Kernels	54
8	Rings	60
8.1	Introduction	60
8.2	Subring	62

1 Set theory

1.1 Reveiw

Definition 1.1. *Set is a collection of distinct objects.*

Here are some properties of a set:

- $\{\text{apple}, 2, \{3\}\}$ is a set.
- If x is in A , we write $x \in A$. If not, we write $x \notin A$.
- \emptyset is an empty set.
- Note that order or repeated elements are not important: $\{1, 2, 3\} = \{3, 1, 2\}$ and $\{1, 1, 1, 2, 2, 3\} = \{1, 2, 3\}$.

Definition 1.2. *Let A and B be sets. B is a subset of A if for all $x \in B$, $x \in A$ and we write $B \subseteq A$. B is a proper subset of A if B is a subset of A but $B \neq A$ and we write $B \subset A$.*

Theorem 1.1. *A and B are equal if and only if $B \subseteq A$ and $A \subseteq B$.*

Example 1.1.1.

- \mathbb{N} is a set of natural numbers: $\{0, 1, 2, 3, \dots\}$.
- \mathbb{Z} is a set of integers: $\{\dots, -2, -1, 0, 1, 2, \dots\}$.
- \mathbb{Q} is a set of rational numbers.
- \mathbb{R} is a set of real numbers.
- \mathbb{C} is a set of complex numbers.

Definition 1.3. *Universal set U contains all elements.*

Let A and B be sets. Then, we can define the following:

Definition 1.4 (Intersection). $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.

Definition 1.5 (Union). $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

Definition 1.6 (Complement). $A' = \{x \mid x \in U \text{ and } x \notin A\}$.

Definition 1.7 (Set difference). $A - B = \{x \mid x \in A \text{ but } x \notin B\}$.

Definition 1.8 (Cartesian product). $A \times B = \{(a, b) \mid a \in A, b \in B\}$.

Example 1.1.2. Let $A = \{0, 1\}$ and $B = \{\text{dog}, \text{cat}\}$. Then,

$$A \times B = \{(0, \text{dog}), (0, \text{cat}), (1, \text{dog}), (1, \text{cat})\}$$

Theorem 1.2 (DeMorgan's Laws). *Let A and B be sets. Then,*

- $(A \cup B)' = A' \cap B'$.
- $(A \cap B)' = A' \cup B'$.

Proof. To show that $(A \cap B)' = A' \cup B'$, we want to show that $(A \cap B)' \subseteq A' \cup B'$ and $A' \cup B' \subseteq (A \cap B)'$.

First, let $x \in (A \cap B)'$. Then, $x \notin (A \cap B)$. So either $x \notin A$ or $x \notin B$. If $x \notin A$, then $x \in A'$. Since $A' \subset A' \cup B'$, $x \in A' \cup B'$. If $x \in B'$, then $x \in B' \subset A' \cup B'$. Therefore, $x \in A' \cup B'$.

Now, we want to prove the opposite direction. Take $x \in A' \cup B'$. So $x \in A'$ or $x \in B'$. Thus, $x \notin A$ or $x \notin B$. In either case, $x \notin (A \cap B)$. Therefore, $x \in (A \cap B)'$. \square

1.2 Equivalence relation

Definition 1.9. Let A and B be sets. Then, a relation is any subset $S \subseteq A \times B$

Example 1.2.1. Let $A = \{0, 1\}$ and $B = \{\text{dog}, \text{cat}\}$. Then,

$$S = \{(0, \text{dog}), (1, \text{cat})\} \subseteq A \times B$$

Functions can give you relations:

Example 1.2.2. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = x^2$. Then, the following is a relation:

$$\{(x, f(x)) \mid x \in \mathbb{R}\} \subseteq \mathbb{R} \times \mathbb{R}$$

Example 1.2.3. Let X be a set of all McMaster students. Then,

$$R = \{(x, y) \mid x \text{ has same height as } y\} \subseteq X \times X$$

Definition 1.10. Let X be a set. An equivalence relation on X is a set $R \subseteq X \times X$ such that

- $(x, x) \in R$ for all $x \in X$ (reflexive)
- If $(x, y) \in R$ and $(y, x) \in R$ (symmetric)
- If $(x, y) \in R$ and $(y, z) \in R$, then $(x, z) \in R$ (transitive)

Example 1.2.4. Example 1.2.1 is not an equivalence relation since $A \neq B$.

Example 1.2.5. Example 1.2.2 is not an equivalence relation since $(2, 2) \notin \{(x, x^2) \mid x \in \mathbb{R}\}$.

Example 1.2.6. Example 1.2.3 is an equivalence relation.

- (reflexive) For any student $x \in X$, x has the same height as x , so $(x, x) \in R$.
- (symmetric) Suppose $(x, y) \in R$ so x and y have the same height. But y and x have the same height so $(y, x) \in R$.

- (transitive) if $(x, y) \in R$ and $(y, z) \in R$, then x and y have the same height and y and z have the same height. So x and z have the same height, i.e. $x, z \in R$.

Remark. Sometimes, we write $x \sim y$ to mean $(x, y) \in R$.

Example 1.2.7. Prove that the following is an equivalence relation

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{Z} \times \mathbb{Z}$$

Proof.

- (reflective) For any $x \in \mathbb{Z}$, $x = x$ and $(x, x) \in R$.
- (symmetric) If $x \sim y$ then $x = y$ so $y = x$, and $y \sim x$.
- (transitive) If $x \sim y$ and $y \sim z$, then $x = y = z$, so $x \sim z$.

□

Definition 1.11. Fix a positive integer $n > 0$. We say r is congruent to s modulo n if n divides $r - s$, i.e. $(r - s) = nl$ for some integer l . We write

$$r \equiv s \pmod{n}$$

Example 1.2.8. Let $n = 7$. Then, $22 \equiv 8 \pmod{7}$ since 7 divides $22 - 8$. However, $22 \not\equiv 10 \pmod{7}$ since 7 does not divide $22 - 10 = 12$.

Example 1.2.9. Congruent definition is an equivalence relation on \mathbb{Z} :

$$R = \{(r, s) \mid r \equiv s \pmod{n}\} \subseteq \mathbb{Z} \times \mathbb{Z}$$

Proof.

- (reflexive) For all $r \in \mathbb{Z}$, n divides $r - r = 0$. So $r \equiv r \pmod{n}$ for all r . So $(r, r) \in R$.
- (symmetric) Suppose $(r, s) \in R$ so $r - s = nl$ for some l . We multiply both sides by (-1) to obtain

$$(s - r) = (-1)(r - s) = (-1)(nl) = n(-l).$$

So n divides $s - r$ and $(s, r) \in R$.

- (transitive) If $(r, s) \in R$ and $(s, t) \in R$, then $r - s = nl$ and $s - t = nk$. But then

$$(r - t) = (r - s) + (s - t) = nl + nk = n(l + k),$$

so $(r, t) \in R$.

□

Definition 1.12. If R is an equivalence relation on X , and $x \in X$, the equivalence class of x is

$$[x] = \{y \mid (x, y) \in R\}$$

Example 1.2.10. Consider

$$R = \{(x, y) \mid x \text{ and } y \text{ have the same height}\}.$$

Then,

$$[\text{Abby}] = \{\text{all people who have same height as Abby}\}.$$

Example 1.2.11. Consider

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{Z} \times \mathbb{Z}.$$

Then,

$$[42] = \{42\}.$$

Example 1.2.12. Consider

$$R = \{(r, s) \mid r \equiv s \pmod{5}\} \subseteq \mathbb{Z} \times \mathbb{Z}.$$

Then,

$$[3] = \{\dots, -7, -2, 3, 8, 13, 18, \dots\}.$$

Definition 1.13. A partition P of set X is a collection of sets, X_0, X_1, X_2, \dots such that

$$X = \bigcup_i X_i$$

and $X_i \cap X_j = \emptyset$ for all $i \neq j$.

Example 1.2.13. In Example 1.2.12, we have

$$\mathbb{Z} = [0] \cup [1] \cup [2] \cup [3] \cup [4]$$

Theorem 1.3. If R is an equivalence relation on X , then the distinct equivalence classes form a partition of X .

Proof. For any $x \in X$, $x \sim x$ so $x \in [x]$. Thus,

$$X = \bigcup_{x \in X} [x].$$

Given $x, y \in X$, we want to show that $[x] = [y]$ or $[x] \cap [y] = \emptyset$. Suppose that $[x] \cap [y] \neq \emptyset$. Let $z \in [x] \cap [y]$. So $x \sim z$ and $y \sim z$. Let $a \in [x]$. Then, $x \sim a$ so $a \sim x$, and $x \sim z$ and $z \sim y$. So $a \sim y$. Thus $y \sim a$, and thus $a \in [y]$. So $[x] \subseteq [y]$.

Same argument shows $[y] \subseteq [x]$. So have $[x] \cap [y] = \emptyset$ or $[x] = [y]$. So considering only distinct classes, we have a partition:

$$X = [x_0] \cup [x_1] \cup \dots,$$

□

1.3 Well ordering principle and division algorithm

Theorem 1.4. (First principle of mathematical induction) *Set $S(n)$ be a statement about integer $n \in \mathbb{N}$ and suppose $S(n)$ is true for some $n_0 \geq 1$. If for all integers $k \geq 0$, if $S(k)$ is true implies $S(k+1)$ is true, then $S(n)$ is true for all $n \geq n_0$.*

Theorem 1.5 (Second principle of mathematical induction). *Let $S(n)$ be a statement for integers $n \in \mathbb{N}$ and assume $S(n_0)$ is true. If $S(n_0), S(n_0 + 1), \dots, S(k)$ imply that $S(k+1)$ is true, then $S(n)$ is true for all $n \geq n_0$.*

Definition 1.14 (Well ordering property). *Every nonempty set of positive integers has a smallest element.*

Remark. Well ordering property becomes false once you include negative values.

Lemma 1.1. *Principle of mathematical induction implies 1 is the smallest integer.*

Theorem 1.6. *Principle of mathematical induction implies well ordering property.*

Proof. Let S be a nonempty set of positive integers. If $1 \in S$, then by above lemma, the set S has a smallest element. Assume that if S is a set that contains $1 \leq k \leq n$, then S satisfies the well ordering property. Let S be any set that contains an integer $1 \leq k \leq n+1$. If S does not contain any elements smaller than $n+1$, $n+1$ is the smallest element. If S does contain an integer $k < n+1$, then by induction step, we have already shown that S has well ordering property. By induction, all S satisfy well ordering property. \square

Remark. Induction and well ordering property are equivalent.

Recall long division. If we divide 304 with 14, we get $304 = 14(21) + 10$. Here, we call 304 a dividend, 14 a divisor, 21 a quotient, and 10 a remainder. Now, we want to know whether this process stops and whether the answer is unique:

Theorem 1.7 (Division algorithm). *Let A and B be integers with $b > 0$. Then, there exists unique integers q and r such that*

$$a = bq + r \text{ with } 0 \leq r < b$$

Proof. To prove that the above theorem is true, we have to show (1) existence and (2) uniqueness.

First, let $S = \{a - bk \mid a - bk \geq 0\}$. If $0 \in S$, then there is a k such that $a - bk = 0 \iff a = bk$. Then, we can let $q = k$ and $r = 0$. If $0 \notin S$, we want to use the well ordering principle. We need to check that $S \neq \emptyset$.

- If $a < 0$, then $a - ba = a(1 - b) > 0$, since $b > 0$. So $S \neq \emptyset$.
- If $a = 0$, then $0 - b(-1) > 0$, so $S \neq \emptyset$.

- If $a > 0$, then $a - b(0) > 0$, so $S \neq \emptyset$.

By the well ordering property, there exists a smallest element say r in S , i.e. there is a q such that $a - bq = r$.

We claim that we also have $0 \leq r < b$. If $r \geq b$,

$$r - b = (a - bq) - b = a - b(q + 1) \geq 0.$$

So $r - b \in S$ and $r - b$ is smaller than r , the smallest element of S . So we must have $0 \leq r < b$.

Now, suppose there was q, r, q', r' such that

$$\begin{cases} a = bq + r, 0 \leq r < b \\ a = bq' + r', 0 \leq r' < b \end{cases}$$

So $bq + r = bq' + r' \implies bq - bq' = r' - r$. Note that

$$-b < -r < r' - r < r' < b.$$

Thus,

$$-b < bq - bq' < b.$$

If we divide both sides by b , we get $-1 < q - q' < 1$. So we find that $q - q' = 0$. \square

Definition 1.15. a divides b if there exists m such that $b = am$. We write $a|b$.

Example 1.3.1. $3|12$ since $12 = 3 \cdot 4$.

Definition 1.16. d is a common divisor of a and b if $d|a$ and $d|b$.

Example 1.3.2. 2 is a common divisor of 12 and 18.

Definition 1.17. d is the greatest common divisor of a and b if (1) d is a common divisor of a and b and (2) if $d'|a$ and $d'|b$, then $d'|d$. We write $d = \gcd(a, b)$.

Example 1.3.3. $6 = \gcd(12, 18)$.

Definition 1.18. a and b are relatively prime if $\gcd(a, b) = 1$.

Remark. For any integer b , $b|0$ since $0 = b \cdot 0$. Furthermore, $\gcd(b, 0) = |b|$.

Theorem 1.8. Let a and b be non-zero integers. Then, there exists r and s such that $\gcd(a, b) = ra + sb$.

Example 1.3.4. $6 = \gcd(12, 18) = 12(-1) + 18 \cdot 1$

Proof. Let $S = \{am + bn \mid m, n \in \mathbb{Z}, am + bn > 0\}$. If $a < 0$, then $a(-1) + b(0) > 0$, so $S \neq \emptyset$. If $a > 0$, then $a(1) + b(0) > 0$ so $S \neq \emptyset$. By the well ordering property, there exists a smallest element in S , say d . So $d = am + bn$ for some $m + n$.

Now, we want to prove that $d = \gcd(a, b)$. First, by the division algorithm, there exists q and r such that $a = dq + r$ with $0 \leq r < d$. If $r > 0$, then,

$$\begin{aligned} r &= a - dq = a - (am + bn)q \\ &= a - amq - bnq \\ &= a(1 - mq) + b(-nq) > 0. \end{aligned}$$

Then $r \in S$ and $r < d$ but d is the smallest element of S . So $r = 0$, i.e. $a = dq + 0$. So $d|a$. Same proof shows $d|b$.

Now, suppose that $d'|a$ and $d'|b$. So $a = d'a'$ and $b = d'b'$. But then

$$\begin{aligned} d &= am + bn \\ &= d'a'm + d'b'n \\ &= d'(a'm + b'n) \end{aligned}$$

So $d'|d$. Hence, $\gcd(a, b) = d$. □

Remark. If $\gcd(a, b) = 1$, then $1 = as + br$ for some s and r .

Lemma 1.2. Suppose a, b, q and r such that $a = bq + r$. Then, $\gcd(a, b) = \gcd(b, r)$.

Proof. Let $d = \gcd(a, b)$ and $e = \gcd(b, r)$. Now, $d|a$ and $d|b$, so $a = da'$ and $b = db'$. Since $r = a - bq$, we have $r = da' - db'q = d(a' - b'q)$. So $d|r$ and $d|b$, so $d \leq \gcd(b, r) = e$.

Now, $e|b$ and $e|r$. So $b = eb^*$ and $r = er^*$. So $a = bq + r = eb^*q + er^* = e(b^*q + r^*)$. So $e|a$ and $e|b$. So $e \leq d$. Hence $d \leq e \leq d$, i.e. $e = d$. □

Now, we introduce the *Euclidean algorithm* to find the greatest common divisors of two integers: To compute $\gcd(a, b)$, repeatedly apply division algorithm:

$$\begin{aligned} a &= bq_1 + r_1 \\ b &= r_1q_1 + r_2 \\ r_1 &= r_2q_2 + r_3 \\ &\vdots \\ r_{n-2} &= r_{n-1}q_{n-1} + r_n \\ r_{n-1} &= r_nq_n + 0 \end{aligned}$$

Then, the last non-zero remainder, r_n is the greatest common divisor.

Remark. This algorithm is guaranteed to stop because r_n is a monotonically decreasing sequence, i.e. $b > r_1 > r_2 > r_3 > \dots \geq 0$. At some point, we must reach $r_{n+1} = 0$ for some n .

Example 1.3.5. We want to find $\gcd(234, 96)$. Note $234 = 96 \cdot 2 + 42$. Note that $\gcd(234, 96) = \gcd(96, 42)$. Then, since $96 = 42 \cdot 2 + 12$, we have $\gcd(96, 42) = \gcd(42, 12)$. Likewise, we can continue to obtain $\gcd(234, 96) = 6$.

Remark. We can reverse this algorithm to find s and t such that $\gcd(a, b) = sa + bt$. Notice that

$$\begin{aligned} 234 &= 96(2) + 42 \\ 96 &= 42(2) + 12 \\ 42 &= 12(3) + 6 \\ 42 &= 234 + 96(-2) \quad 12 = 96 + 42(-2) \\ 6 &= 42 + 12(-3) \end{aligned}$$

So

$$\begin{aligned} 6 &= 42 + [96 + 42(-2)](-3) \\ &= 42(7) + 96(-3) \end{aligned}$$

Then,

$$\begin{aligned} 6 &= [234 + 96(-2)](7) + 96(-3) \\ &= (234)(7) + 96(-3) + 96(-3) \\ &= 234(7) + 96(-17) \end{aligned}$$

Definition 1.19. A positive integer $p > 1$ is prime if its only divisions are 1 and p . Otherwise, a number is composite.

Example 1.3.6. 7 is a prime.

Lemma 1.3. Let a and b be integers and p a prime. If $p|ab$, then $p|a$ or $p|b$. This statement is false when p is not a prime.

Proof. If $p \nmid a$, we want to show that $p|b$. If $p \nmid a$, then $\gcd(a, p) = 1$. So there exists s and t such that $1 = as + pt$. Then, we have $b = abs + pbt$. Since $p|ab$, we have $ab = pk$. So,

$$b = pks + pbt = p(ks + bt).$$

Therefore, $p|b$. □

Theorem 1.9 (Fundamental theorem of arithmetic). Let $n > 1$ be any integer.

$$n = p_1 p_2 \cdots p_k,$$

where p_i is a prime (not necessarily distinct). Furthermore, this decomposition is unique in the following sense. If $n = q_1 \cdots q_l$ is another production of primes, then $k = l$ and after relabelling, $p_i = q_i$.

Proof. (Existence) Let

$$S = \{a \in \mathbb{Z} \mid a > 1 \text{ and } a \text{ does not have a primary decomposition}\}.$$

If $S \neq \emptyset$, then by the well ordering principle, there is a smallest $a \in S$. Note a is not a prime because if a is prime then $a = a$ is a factorization. So a is

composite and $a = bc$ with $1 < b, c < a$. However, $b, c \notin S$ so they have a factorization:

$$\begin{aligned} b &= p_1 \cdots p_l \\ c &= q_1 \cdots q_k \end{aligned}$$

But then $a = p_1 \cdots p_l q_1 \cdots q_k$. So $a \notin S$, This is a contradiction and $S = \emptyset$.
(Uniqueness). Suppose

$$n = p_1 \cdots p_k = q_1 \cdots q_l$$

Since $p_1 | n$, $p_1 | q_1 \cdots q_l$. So $p_1 | q_i$ for some i by the Lemma. Since q_i is prime and $p_1 > 1$, then $p_1 = q_i$. Then, we do a relabelling so that q_i is q_1 . So we have

$$\begin{aligned} p_1 p_2 \cdots p_k &= q_1 q_2 \cdots q_l \\ \implies p_2 \cdots p_k &= q_2 \cdots q_l \end{aligned}$$

We repeat the process. If $k > 1$, we would end with

$$p_{l+1} p_{l+2} \cdots p_k = 1.$$

Likewise, we would end with a similar equation if $k < l$. Both cases are impossible because $p_i, q_i > 1$. So $k = l$ and $p_i = q_i$ for all i . \square

Theorem 1.10. *There exists an infinite number of primes.*

Proof. Suppose only primes are p_1, p_2, \dots, p_n . Let

$$P = p_1 p_2 \cdots p_n + 1.$$

Since $P > p_1, \dots, p_n$, P is not a prime. So P is a composite number by FTA, some p_i must divide P . Since $P - p_1 p_2 \cdots p_n = 1$, then $p_i | 1$, yielding contradiction. So there must be infinite number of primes. \square

Example 1.3.7. Prove that if $\gcd(a, b) = 1$ and $a | bc$, then $a | c$.

Proof. Because $\gcd(a, b) = 1$, there exists integers s and t such that $as + bt = 1$. This follows from theorem 2.10. If we multiply both sides by c , we get

$$acs + bct = c$$

Since $a | bc$, $bc = ak$ for some integer k . After substitution, we have

$$c = acs + akt.$$

But this means

$$c = a(cs + kt).$$

So $a | c$, as desired. \square

2 Groups and rings

2.1 Group theory

Before we begin, we're going to look at sets with *extra structure*.

Example 2.1.1 (Integer equivalence classes). Let $n = 6$. Consider the distinct equivalence classes modulo 6:

$$R = \{(a, b) \mid a \equiv b \pmod{6}\} \subseteq \mathbb{Z} \times \mathbb{Z}$$

Then,

$$[0] = \{\dots, -6, 0, 6, \dots\}$$

$$[1] = \{\dots, -5, 1, 7, \dots\}$$

$$[2] = \{\dots, -4, 2, 8, \dots\}$$

$$[3] = \{\dots, -3, 3, 9, \dots\}$$

$$[4] = \{\dots, -2, 4, 10, \dots\}$$

$$[5] = \{\dots, -1, 5, 11, \dots\}$$

We denote the six distinct equivalence classes by

$$\mathbb{Z}_6 = \{[0], [1], [2], [3], [4], [5]\}.$$

Usually, we write

$$\mathbb{Z}_6 = \{0, 1, 2, 3, 4, 5\}.$$

In general, for any $n > 1$, let

$$\mathbb{Z}_n = \{0, 1, 2, \dots, n-1\}.$$

Then, we can add and multiply elements of \mathbb{Z}_n :

$$a + b = (a + b) \pmod{n}$$

$$ab = (ab) \pmod{n}$$

In fact, for any $a \in \mathbb{Z}$ and $n > 1$, if $a = nq + r$ with $0 \leq r < n$, then $[a] = [r]$. Equivalently, $a \equiv r \pmod{n}$ and $a \equiv r$ in \mathbb{Z}_n .

We can look at some other properties of addition and multiplication in \mathbb{Z}_n :

- Addition and multiplication commute
- Addition and multiplication are associative
- There are additive and multiplicative identities
- For every element in \mathbb{Z}_n , there exists an additive inverse.
- Multiplication is distributive over addition
- If $\gcd(a, n) = 1$, then there exists an integer b such that $ab \equiv 1 \pmod{n}$.

Consider a square cut in the plane. We can flip it, rotate it, and but not stretch it, and then put it back in the original spot. Then, we have 8 operations.

Let R_0 be rotating 0° , R_{90} rotating 90° , R_{180} rotating 180° , and R_{270} rotating 270° . Then, H will be a flip on the horizontal axis, V on the vertical axis, D_1 on the main-diagonal, and D_2 on the anti-diagonal. Note that you can perform one operation, then followed by another, and end back up with another known operation. For example H, R_{270} is equivalent to D_1 . Note that order is important.

We want to think of these as functions, i.e., each function maps a square to itself. Let

$$D_4 = \{R_0, R_{90}, R_{180}, R_{270}, V, H, D_1, D_2\}.$$

We call is a dihedral group and it has the following properties:

- Operations of composition is closed.
- R_0 is an identity element.
- Each element $A \in D_4$ has an inverse, i.e., we can reverse it to R_0 .
- The operation is associative.

In fact, D_4 forms a group and those are the four properties that all groups must have.

Now, we want to formally define a group.

Definition 2.1. *Given any set G , a binary operation \circ is any function*

$$\circ : G \times G \rightarrow G$$

that maps a pair $(a, b) \in G \times G$ to an element $a \circ b$.

Example 2.1.2. $+$ on \mathbb{Z} is a binary operation

$$+ : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$$

. Likewise, multiplication is also a binary operation.

Example 2.1.3. Composition of functions on D_4 is a binary operation:

$$\circ_{D_4} : D_4 \times D_4 \rightarrow D_4$$

Definition 2.2. *A group (G, \circ) is a set G with a binary operation \circ such that*

- *(associative) $a \circ (b \circ c) = (a \circ b) \circ c$.*
- *(identity) there exists an $e \in G$ such that $a \circ e = e \circ a = a$ for all $a \in G$.*
- *(inverse) for all $a \in G$ exists $a^{-1} \in G$ such that $a \circ a^{-1} = a^{-1} \circ a = e$.*

Definition 2.3. If a group G satisfies commutativity,

$$a \circ b = b \circ a, \forall a, b \in G,$$

then G is called abelian.

Example 2.1.4. D_4 is a group where the binary operation is composition of functions. D_4 is not abelian since

$$D_1 \circ H \neq H \circ D_1$$

Example 2.1.5. Consider

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}.$$

There are two operations on \mathbb{Z} : addition and multiplication. \mathbb{Z} with addition is an abelian group with identity 0. However, \mathbb{Z} with multiplication is not a group because it doesn't have an inverse.

Example 2.1.6. Rationals, real numbers, and complex numbers are all groups with operation of $+$.

Example 2.1.7 (Trivial group). $G = \{e\}$.

Example 2.1.8. Fix $n > 1$. Then, $\mathbb{Z}_n = \{0, 1, 2, \dots, n-1\}$ is a group under addition. However, it's not a group under multiplication.

Example 2.1.9. \mathbb{R} is not a group under multiplication. It satisfies associativity and existence of identity but 0 does not have a multiplicative inverse. However,

$$\mathbb{R}^* = \mathbb{R} \setminus \{0\}$$

is a group under multiplication. Likewise, $\mathbb{Q}^* = \mathbb{Q} \setminus \{0\}$ and $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ are groups under multiplication.

Example 2.1.10. Let $n > 1$ and

$$u(n) = \{a \mid 1 \leq a \leq n-1, \gcd(a, n) = 1\}.$$

For example,

$$\begin{aligned} u(3) &= \{1, 2\} & u(5) &= \{1, 2, 3, 4\} \\ u(4) &= \{1, 3\} & u(8) &= \{1, 3, 5, 7\} \end{aligned}$$

For all $n > 1$, $u(n)$ is a group under multiplication modulo n .

Example 2.1.11. Consider

$$M_2(\mathbb{R}) = \{\text{all } 2 \times 2 \text{ matrices with entries in } \mathbb{R}\}.$$

This set is a group under addition.

Example 2.1.12. All vector spaces are groups under addition.

Example 2.1.13 (General linear group).

$$GL_2(\mathbb{R}) = \{\text{all } 2 \times 2 \text{ matrices that are invertible}\} \\ = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid ad - cb \neq 0 \right\}$$

This is a group under matrix multiplication.

We want to make new groups from existing groups. Let G and H be groups and that let \square and $*$ denote their binary operations. Then,

$$G \times H = \{(g, h) \mid g \in G, h \in H\}.$$

This is also a group where

$$(g_1, h_1) \circ (g_2, h_2) = (g_1 \square g_2, h_1 * h_2).$$

Example 2.1.14. Consider

$$G = \mathbb{Z}_3 = \{0, 1, 2\}, H = \mathbb{R}^* = \mathbb{R} \setminus \{0\}$$

Then,

$$(2, 4) \circ (2, 6) = (2 + 2, 4 \times 6) = (1, 24) \in G \times H.$$

In this case, the identity of $\mathbb{Z}_3 \times \mathbb{R}^*$ is $(0, 1)$.

Definition 2.4. The order of G refers to number of elements in G and is denoted by $|G|$. G is finite if $|G| < \infty$. Otherwise, it is infinite.

There are many different binary operations used to define groups. Normally, we will use the multiplicative notation. The only exception is when we are proving something about an additive group.

From now on, we will be using the following notations:

$$a^n = \begin{cases} a \cdot a \cdots a & (\text{n times}) \text{ if } n > 0 \\ 1 & n = 0 \\ (a^{-1} \cdots a^{-1}) & n < 0 \end{cases}$$

$$na = \begin{cases} a + a + \cdots + a & (\text{n times}) \text{ if } n > 0 \\ 0 & n = 0 \\ (-a) + (-a) + \cdots + (-a) & n < 0 \end{cases}$$

Theorem 2.1. For every group G , identity is unique.

Proof. Suppose e and e' are identities of G . So for any $a \in G$, (1) $ae = a$ and (2) $e'a = a$. If $a = e'$, (1) implies $e'e = e'$. If $a = e$, (2) implies $e'e = e$. So

$$e' = e'e = e,$$

and $e' = e$. □

Theorem 2.2. *If $g \in G$, then inverse of g is unique.*

Proof. Suppose that g' and g'' are inverses of g . So $g'g = gg' = e$ and $g''g = gg'' = e$. So

$$gg' = gg'' = e.$$

If we multiply both sides by g' ,

$$\begin{aligned} g'(gg') &= g'(gg'') \\ \implies (g'g)g' &= (g'g)g'' \\ \implies eg' &= g' = g'' = eg''. \end{aligned}$$

□

Theorem 2.3 (Socks-shoes property). $(ab)^{-1} = b^{-1}a^{-1}$.

Proof. By definition, $(ab)^{-1}$ is the inverse of (ab) , i.e.,

$$(ab)(ab)^{-1} = e.$$

But we also have

$$\begin{aligned} (ab)(b^{-1}a^{-1}) &= a(bb^{-1})a^{-1} \\ &= aea^{-1} \\ &= aa^{-1} \\ &= e. \end{aligned}$$

So $b^{-1}a^{-1}$ is also an inverse of (ab) . Since inverses are unique, we have

$$(ab)^{-1} = b^{-1}a^{-1}.$$

□

Theorem 2.4. *If G , cancellation works, i.e. if $ab = bc$, then $a = c$.*

Proof. Suppose that $ab = ac$. Then, $a^{-1} \in G$. So we multiply both sides by a^{-1} on the left

$$a^{-1}(ab) = a^{-1}(ac).$$

So $b = c$.

□

Remark. As a consequence, each row and column in a Cayley table (group operation table) has a distinct element. In other words, if $ab_i = ab_j$ then $b_i = b_j$

Theorem 2.5. *For any $a, b \in G$, there exists unique x and y such that $ax = b$ and $ya = b$.*

Proof. One solution is $x = a^{-1}b$ since

$$a(a^{-1}b) = (aa^{-1})b = b.$$

This is unique because if $ax_1 = b = ax_2$, by cancellation $x_1 = x_2$.

□

2.2 Subgroups

Definition 2.5. A subset H of a group G is a group if it is a group under the same operation of G .

Example 2.2.1. If $G \neq \{e\}$, the G has at least two subgroups:

- $\{e\} \subseteq G$,
- G itself.

These are trivial groups but we want $\{e\} \subset H \subset G$.

Example 2.2.2. Consider $G = \mathbb{Z}$. Then,

$$E = \{n \in G \mid n \text{ is even}\} = \{-4, -2, 0, 2, 4\}$$

is a subgroup because

- because it is closed under addition.
- $0 \in E$.
- addition is associative.
- for any $a \in E$, $-a \in E$ so every element in E has an inverse.

Example 2.2.3. The set of odd integers is not a subgroup because it is not closed under addition and 0 is not an element.

Example 2.2.4. $m\mathbb{Z} = \{mn \mid n \in \mathbb{Z}\}$ is a subgroup.

Example 2.2.5. Consider D_4 . Let $H = \{R_0, R_{90}, R_{180}, R_{270}\}$. Note D_4 is not abelian but H is.

Example 2.2.6. Consider $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$, a group under multiplication. Then,

$$H = \{1, -1, i, -i\}$$

is a finite subgroup of \mathbb{C}^* :

	1	i	-1	$-i$
1	1	i	-1	$-i$
i	i	-1	$-i$	1
-1	-1	$-i$	1	i
$-i$	$-i$	1	i	-1

Example 2.2.7. Show that if $a^2 = e$ for all $a \in G$ then G is abelian.

Proof. Given any $a, b \in G$, we want to show $ab = ba$. Given that $aa = e$, since inverses are unique, $a = a^{-1}$. Now, consider $(ab)^2$. Since $ab \in G$,

$$(ab^2) = (ab)(ab) = e.$$

Now, we multiply $(ab)(ab) = e$ on the left by a and on the right by b :

$$\begin{aligned} a(ab)(ab)b &= aeb \\ (aa)(ba)(bb) &= ab \\ ba &= ab \end{aligned}$$

So G is abelian. □

Theorem 2.6. *A subset H of a group G is a subgroup if*

- $e \in H$.
- $\forall g_1, g_2 \in H, g_1 \circ g_2 \in H$.
- $\forall g \in H, g^{-1} \in H$

Proof. 1 implies that H has an identity, 2 implies that H is closed under operation. 3 implies that every $g \in H$ has an inverse. So we only need to check the associative property.

Let $a, b, c \in H$. Now, $a, b, c \in G$, so

$$(ab)c = a(bc)$$

holds in G . But since the operation is closed, (ab) and (bc) are in H , so

$$(ab)c = a(bc)$$

also holds in H . □

Definition 2.6 (Center of a group). *For any group G , the center of G is defined as*

$$Z(G) = \{a \in G \mid ag = ga, \forall g \in G\}.$$

Example 2.2.8. If G is abelian, $G = Z(G)$. If $G = D_3$, $Z(D_4) = \{R_0, R_{180}\}$. For all G , $e \in Z(G)$.

Theorem 2.7. *For all G , $Z(G)$ is a subgroup of G .*

Proof. First, $e \in Z(G)$ since for all $g \in G$,

$$eg = g = ge.$$

Let $a, b \in Z(G)$. We want to show that $ab \in Z(G)$. So for any $g \in G$, we need to show that $(ab)g = g(ab)$. To prove this, take any $g \in G$. Then,

$$\begin{aligned}(ab)g &= a(bg) \quad (\text{associativity}) \\ &= a(gb) \quad (\text{since } b \in Z(G)) \\ &= (ag)b \quad (\text{associativity}) \\ &= (ga)b \quad (\text{since } a \in Z(G)) \\ &= g(ab) \quad (\text{associativity})\end{aligned}$$

So $ab \in Z(G)$.

Now, let $a \in Z(G)$ and take any $g \in G$. So $g^{-1} \in G$, and since $a \in Z(G)$,

$$ag^{-1} = g^{-1}a.$$

Taking the inverse of both sides gives

$$ga^{-1} = (ag^{-1})^{-1} = (g^{-1}a)^{-1} = a^{-1}g.$$

So for any $a \in Z(G)$ and any $g \in G$,

$$a^{-1}g = ga^{-1},$$

i.e. $a^{-1} \in Z(G)$. □

Example 2.2.9. If every proper subgroup of group G is abelian, is G abelian?

Proof. No. D_4 is not abelian. However, all proper subgroups are abelian.

$$\begin{aligned}H_1 &= \{R_0, R_{90}, R_{180}, R_{270}\} \\ H_2 &= \{R_0, R_{180}\} \\ H_3 &= \{R_0, D_1\} \\ H_4 &= \{R_0, D_2\} \\ H_5 &= \{R_0, V\} \\ H_6 &= \{R_0, H\} \\ H_7 &= \{R_0, D_{1,2}, H, V\}\end{aligned}$$

□

3 Special groups

3.1 Cyclic groups

So how do we find subgroups? Here's one way to construct subgroups:

Definition 3.1. Fix an $a \in G$. Then, $\langle a \rangle = \{a^n \mid n \in \mathbb{Z}\}$

Example 3.1.1. Consider $G = D_4$. Then, since $R_{90} \in G_4$,

$$\begin{aligned}\langle R_{90} \rangle &= \{R_{90}^{-1}, R_0, R_{90}, R_{90} \circ R_{90}, \dots\} \\ &= \{R_0, R_{90}, R_{180}, R_{270}\}.\end{aligned}$$

Example 3.1.2. If $G = \mathbb{Z}_6$ and $2 \in G$, then

$$\begin{aligned}\langle 2 \rangle &= \{2 - 2 - 2, 2 - 2, 2, 2 + 2, 2 + 2 + 2, \dots\} \\ &= \{2, 4, 0\}.\end{aligned}$$

Theorem 3.1. For any $a \in G$, $\langle a \rangle$ is a subgroup of G and it is the smallest subgroup of G that contains a .

Proof. First, $e \in \langle a \rangle$ since $e = a^0$. Now, suppose that $g_1, g_2 \in \langle a \rangle$. So $g_1 = a^{n_1}$ and $g_2 = a^{n_2}$. But then,

$$g_1 g_2 = a^{n_1} a^{n_2} = a^{n_1 + n_2} \in \langle a \rangle.$$

Finally, if $a^n \in \langle a \rangle$ then $(a^n)^{-1} = a^{-n} \in \langle a \rangle$. So $\langle a \rangle$ is a subgroup.

To prove that it is the smallest subgroup, consider a subgroup H with $a \in H$. Then, a^1, a^2, a^3 and a^0, a^{-1}, a^{-2} are also in H . So $\langle a \rangle \subseteq H$. \square

Definition 3.2. If G contains an element a such that $G = \langle a \rangle$, then we say G is cyclic and a is the generator.

Example 3.1.3. \mathbb{Z}_6 is cyclic since $\mathbb{Z}_6 = \langle 5 \rangle$.

Definition 3.3. If $a \in G$, then the order of a is the smallest positive integer such that $a^n = e$. We write $|a| = n$. If order is not finite, $|a| = \infty$.

Example 3.1.4. Consider $G = \mathbb{Z}_6$. Then,

- $|3| = 2$ since $3 + 3 = 0$.
- $|5| = 6$ since $5 + 5 + 5 + 5 + 5 + 5 = 0$.

Example 3.1.5. Consider \mathbb{Z} with addition. Then, $|1| = \infty$.

Example 3.1.6. Consider \mathbb{Z}_n with addition. Then, $|1| = n$.

Example 3.1.7. Consider $u(8) = \{1, 3, 5, 7\}$ under multiplication. Observe that

$$\begin{aligned}|1| &= 1 \\ |3| &= 2 \\ |5| &= 2 \\ |7| &= 2\end{aligned}$$

$u(8)$ is not cyclic because no element with $|a| = |u(8)| = 4$.

Theorem 3.2. Every cyclic group is abelian.

Proof. Let $g_1, g_2 \in \langle a \rangle$. So $g_1 = a^{n_1}$ and $g_2 = a^{n_2}$ for some n_1, n_2 . Then,

$$g_1 g_2 = a^{n_1} a^{n_2} = a^{n_1+n_2} = a^{n_2+n_1} = a^{n_2} a^{n_1} = g_2 g_1.$$

□

Theorem 3.3. *If G is cyclic, all subgroups are cyclic.*

Proof. Let $H \subseteq G$ be a subgroup of $G = \langle a \rangle$. If $H = \{e\}$ and if $H = G$, then H is cyclic.

So assume that $\{e\} \subset H \subset G$. If $g \in H$, then $g = a^n$ for some $n \in \mathbb{Z}$. Since $g^{-1} = a^{-n}$, we know that at least one of n or $-n$ is positive.

Let m be the smallest positive integer such that $a^m \in H$. We claim that $H = \langle a^m \rangle$. If $a^m \in H$, then $\langle a^m \rangle \subseteq H$. Take $g = a^n \in H$. Then, we can divide n by m using the division algorithm, i.e.,

$$n = mq + r,$$

with $0 \leq r < m$. If $0 < r < m$, then

$$a^n = a^{mq+r} = a^{mq} a^r.$$

Since $a^{mq} \in H$, $a^{-mq} \in H$. So

$$a^n a^{-mq} = a^{n-mq} = a^r \in H.$$

However, this contradicts our assumption that m is the smallest positive exponent in H . Therefore, $r = 0$. Hence, $n = mq$, so $g = a^n = (a^m)^q \in \langle a^m \rangle$. So H is cyclic. □

Recall that the order of $a \in G$, denoted $|a|$, is smallest positive integer n such that $a^n = e$. The order of G , denoted $|G|$, is number of elements in G .

Theorem 3.4. *Let $a \in G$.*

- *If $|a| = \infty$, then $a^i = a^j$ if and only if $i = j$.*
- *If $|a| = n$, then $a^i = a^j$ if and only if $n|(i - j)$.*
- *If $|a| = n$, Then, $\langle a \rangle = \{a^0, a^1, a^2, \dots, a^{n-1}\}$. Also, $|a| = |\langle a \rangle|$.*

Proof. (1) Because $|a| = \infty$, all elements of $\langle a \rangle$ are distinct. Indeed, if $a^i = a^j$, then $a^i a^{-j} = e$. So $a^{i-j} = e$. But $|a| = \infty$, so $a^{i-j} = e$ iff $i - j = 0$, i.e., $i = j$.

(2) Suppose that $a^i = a^j$. Without loss of generality, we can assume that $i > j$. So $a^i a^{-j} = e$. Now, we can divide $(i - j)$ by n using division algorithm, i.e.,

$$(i - j) = nq + r,$$

with $0 \leq r < n$. If $0 < r < m$, then

$$a^{i-j} = (a^n)^q a^r = a^r = e.$$

This means $a^r = e$ with $r < n$. However, this contradicts the assumption that $|a| = n$. So $r = 0$ and $n|(i - j)$. To prove the other direction, assume that $n|(i - j)$. Then, $(i - j) = nq$ and $i = nq + j$. Then,

$$a^i = a^{nq+j} = (a^n)^q a^j = a^q a^j = a^j.$$

(3) We want to show that $\langle a \rangle = \{a^0, a^1, \dots, a^{n-1}\}$. Take $a^k \in \langle a \rangle$. Then, we divide k by n using division algorithm:

$$k = nq + r,$$

with $0 \leq r < n$. So

$$a^k = a^{nq+r} = a^r.$$

So $a^k = a^r \in \{a^0, a^1, \dots, a^{n-1}\}$. □

Example 3.1.8. Consider

$$\mathbb{Z}_5 = \{0, 1, 2, 3, 4\}.$$

This is a cyclic group generated by 1. So $|1| = 5$. Note that $7 \cdot 1 = 2 = 22 \cdot 1$ and $5|(22 - 7)$.

Corollary 3.1. *For any cyclical group $G = \langle a \rangle$, if $|a| = n$, and $a^k = e$, then $n|k$.*

Proof. Apply (2) with $i = k$ and $j = 0$. □

Theorem 3.5. *If $|a| = n$, then $|a^k| = n/\gcd(n, k)$.*

Example 3.1.9. Consider

$$\mathbb{Z}_6 = \{0, 1, 2, 3, 4, 5\}.$$

Then,

$$\langle 1 \rangle = \{1, 1+1, 1+1+1, 1+1+1+1, 1+1+1+1+1, 1+1+1+1+1+1\}.$$

Then, $|1| = 6$. So

$$\langle 1 \rangle = \{1 \cdot 1, 2 \cdot 1, 3 \cdot 1, 4 \cdot 1, 5 \cdot 1, 0 \cdot 1\}.$$

So

$$|2 \cdot 1| = \frac{6}{\gcd(2, 6)} = \frac{6}{2} = 3$$

$$|3 \cdot 1| = \frac{6}{\gcd(3, 6)} = \frac{6}{3} = 2$$

$$|4 \cdot 1| = \frac{6}{\gcd(4, 6)} = \frac{6}{2} = 3$$

Corollary 3.2. *For any $k \in \mathbb{Z}$, $\mathbb{Z}_n = \langle k \rangle$ iff $\gcd(n, k) = 1$.*

Proof. Observe that

$$k = 1 + 1 + 1 + \cdots + 1 = k \cdot 1$$

with $|1| = n$. So

$$|k| = \frac{n}{\gcd(n, k)}.$$

So $\langle k \rangle = \mathbb{Z}_n$ iff $|k| = n$ iff $n = n / \gcd(n, k)$ iff $\gcd(n, k) = 1$. \square

Now, recall that $\mathbb{Q}^* = \mathbb{Q} \setminus \{0\}$, $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$, and $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$ are multiplicative groups. We are interested in finding finite multiplicative subgroups.

Theorem 3.6. *In \mathbb{Q}^* and \mathbb{R}^* , there are only two finite subgroups, which are $\{1\}$ and $\{1, -1\}$.*

Proof. Take any $H \subseteq \mathbb{Q}^*$ be a subgroup with $|H| < \infty$. Let $a \in H$. Then, $a^n = 1$ for some n . So, a satisfies

$$a^n - 1 = 0 = (a - 1)(a^{n-1} + a^{n-2} + \cdots + a + 1).$$

If $n = 1$, then $a = 1$. If $n = 2$, then $a^2 = 1$ so $a = \pm 1$.

If $n < 3$, a would have to take a root of

$$x^{n-1} + x^{n-2} + \cdots + x + 1 = 0,$$

or $(x - 1)$. But the former equation does not have real or rational roots. So $a = 1$.

Therefore, $H = \{1\}$ or $H = \{-1, 1\}$. \square

Recall the following properties of complex numbers:

- $(a + bi)(c + di) = (ac - bd) + (ad + bc)i$.
- $|a + bi| = \sqrt{a^2 + b^2}$.
- $(a + bi)^{-1} = (a - bi)/(a^2 + b^2)$.

We can represent complex numbers using polar coordinates:

$$a + bi = z = |z|(\cos \theta + i \sin \theta),$$

and we denote it by $r \operatorname{cis} \theta$. It is convenient to use polar coordinates due to the following property:

Theorem 3.7. *If $z_1 = r_1 \operatorname{cis} \theta_1$ and $z_2 = r_2 \operatorname{cis} \theta_2$. Then,*

$$\begin{aligned} z_1 z_2 &= r_1 r_2 (\operatorname{cis}(\theta_1 + \theta_2)) \\ z_1^{-1} &= r_1^{-1} \operatorname{cis}(-\theta_1) \end{aligned}$$

Definition 3.4 (Circle subgroup).

$$\mathbb{T} = \{z \in \mathbb{C}^* \mid |z| = 1\}$$

is a subgroup.

Proof.

- (identity) $1 \in \mathbb{T}$ since $|1| = 1$.
- (closure) Suppose $z_1, z_2 \in \mathbb{T}$. So $z_1 = 1 \operatorname{cis} \theta_1$ and $z_2 = 1 \operatorname{cis} \theta_2$. So $z_1 z_2 = 1 \cdot 1 \operatorname{cis}(\theta_1 + \theta_2) \in \mathbb{T}$.
- (inverse) If $z = 1 \operatorname{cis} \theta \in \mathbb{T}$, then $z^{-1} = 1 \operatorname{cis}(-\theta) \in \mathbb{T}$.

□

Definition 3.5. Fix $n \geq 1$. The complex numbers that satisfy $x^n - 1 = 0$ are called n -th root of unity.

Remark. $x^n - 1$ has n roots (up to multiplicity) in \mathbb{C} .

Example 3.1.10. Consider $n = 3$,

$$x^3 - 1 = 0.$$

Roots are $1, w, w^2, \dots$, where

$$w = \frac{-1 + \sqrt{3}i}{2}, w^2 = \frac{-1 - \sqrt{3}i}{2}.$$

Theorem 3.8. The set of n -th root of unity form a cyclic group of order n in \mathbb{C}^* . Furthermore, the n -th root of unity are

$$z = \operatorname{cis} \left(\frac{2k\pi}{n} \right),$$

for $k = 0, 1, 2, \dots, n-1$.

Definition 3.6. A generator of the n -th group of units is called a primitive n -th root.

Example 3.1.11. If $n = 8$, primitive roots are

$$w, w^3, w^5, w^7,$$

and the rest are non-primitive roots.

Example 3.1.12. Find all cyclic subgroups of \mathbb{Z}_8 .

$$\begin{aligned} \langle 0 \rangle &= \{0\} \\ \langle 1 \rangle &= \langle 3 \rangle = \langle 5 \rangle = \langle 7 \rangle = \mathbb{Z}_8 \\ \langle 4 \rangle &= \{0, 4\} \\ \langle 2 \rangle &= \langle 6 \rangle = \{0, 2, 4, 6\} \end{aligned}$$

Example 3.1.13. Find all cyclic subgroups of $u(9)$.

$$\begin{aligned}\langle 1 \rangle &= \{1\} \\ \langle 2 \rangle &= u(9) = \langle 5 \rangle \\ \langle 4 \rangle &= \{4, 7, 1\} = \langle 7 \rangle \\ \langle 8 \rangle &= \{1, 8\}.\end{aligned}$$

Example 3.1.14. Prove that the order of every element in a cyclic group G divides $|G|$.

Example 3.1.15. Suppose $|G| = p$, a prime and G cyclic. Show that every nonidentity element has order p .

3.2 Permutation groups

Definition 3.7. A permutation of a set X is a bijection:

$$\sigma : X \rightarrow X$$

Definition 3.8. A permutation group of a set X is the set of all permutations of X with binary operation composition of functions.

Example 3.2.1. Consider

$$X = \{1, 2, 3, 4, 5\}$$

Then, given

$$\begin{aligned}\sigma : X &\rightarrow X \\ 1 &\rightarrow 1 \\ 2 &\rightarrow 3 \\ 3 &\rightarrow 4 \\ 4 &\rightarrow 2 \\ 5 &\rightarrow 5\end{aligned}$$

and

$$\begin{aligned}\tau : X &\rightarrow X \\ 1 &\rightarrow 2 \\ 2 &\rightarrow 3 \\ 3 &\rightarrow 1 \\ 4 &\rightarrow 5 \\ 5 &\rightarrow 4\end{aligned},$$

we have

$$\begin{aligned}\sigma \cdot \tau : X &\rightarrow X \\ 1 &\rightarrow 3 \\ 2 &\rightarrow 4 \\ 3 &\rightarrow 1 \\ 4 &\rightarrow 5 \\ 5 &\rightarrow 2\end{aligned}$$

To avoid writing like this, we introduce a better notation:

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 3 & 4 & 2 & 5 \end{pmatrix}, \tau = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 1 & 5 & 4 \end{pmatrix}.$$

Then,

$$\sigma \cdot \tau = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 3 & 4 & 2 & 5 \end{pmatrix} \cdot \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 1 & 5 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 4 & 1 & 5 & 2 \end{pmatrix}$$

Likewise,

$$\tau \cdot \sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 1 & 5 & 3 & 4 \end{pmatrix}$$

Note that $\sigma \cdot \tau \neq \tau \cdot \sigma$. In general, a permutation group is not abelian.

Definition 3.9. Fix an integer $n \geq 1$. The symmetric group on n letters, denoted S_n , is the set of all permutations of $\{1, 2, 3, \dots, n\}$.

Theorem 3.9. S_n is a non-abelian group (if $n \geq 3$).

Proof.

- S_n has an identity

$$\begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 1 & 2 & 3 & \cdots & n \end{pmatrix}$$

- Each element has an inverse (reverse the map the permutation)
- Composition is associative

□

Remark. Note that there are $n!$ permutations.

Example 3.2.2.

$$S_3 = \left\{ \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} \right\}$$

Now, we introduce a cyclic notation. Consider

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 4 & 3 & 6 & 2 & 5 \end{pmatrix}$$

Note that 1 and 3 map to themselves whereas we have

$$2 \rightarrow 4 \rightarrow 6 \rightarrow 5.$$

So we write

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 4 & 3 & 6 & 2 & 5 \end{pmatrix} = (2465),$$

which means that

- each element is mapped to the one to right
- the last element is mapped to the front
- elements that do not appear are mapped to themselves

Example 3.2.3.

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 1 & 4 & 6 & 5 & 3 \end{pmatrix} = (12)(346)$$

Example 3.2.4.

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 2 & 1 & 3 & 5 & 6 \end{pmatrix} = (143) = (314) = (431)$$

Definition 3.10. A permutation of form (a_1, a_2, \dots, a_k) is called a k -cycle.

Theorem 3.10. If two cycles, σ and τ , are disjoint cycles, (i.e., they don't share any common values), then $\sigma \cdot \tau = \tau \cdot \sigma$.

Proof. Let $\sigma = (a_1, \dots, a_k)$ and $\tau = (b_1, \dots, b_l)$. We know that

$$\sigma \cap \tau = \emptyset.$$

Then,

- if $x \in \{1, 2, \dots, n\}$ but $x \notin \sigma \cup \tau$, then $\sigma(x) = x$ and $\tau(x) = x$ so $\sigma(\tau(x)) = x = \tau(\sigma(x))$.
- suppose $x \in \sigma$ so $x = a_i$ for some i and $x \notin \tau$. Now, $\sigma(x) = \sigma(a_i) = a_{i+1}$. Also, $\tau(x) = x$ and $\tau(a_{i+1}) = a_{i+1}$. So

$$\sigma(\tau(a_i)) = \sigma(a_i) = a_{i+1} = \tau(a_{i+1}) = \tau(\sigma(a_i))$$

□

Example 3.2.5.

$$(12)(346) = (346)(12)$$

Remark. Not every permutation can be expressed as a cycle

Theorem 3.11. *Every permutation can be expressed as a product of disjoint cycles.*

We will illustrate this with an example, rather than a proof. Consider

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 3 & 7 & 5 & 2 & 1 & 8 & 4 & 6 \end{pmatrix}$$

We start with an element that is not mapped to itself, i.e.,

$$1 \rightarrow 3 \rightarrow 5 \implies (135)$$

Now, take another element not in previous step and is not mapped to itself

$$2 \rightarrow 7 \rightarrow 4 \implies (274)$$

We can do the same thing for the rest and get

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 3 & 7 & 5 & 2 & 1 & 8 & 4 & 6 \end{pmatrix} = (135)(274)(68)$$

The advantage of doing this is that it's easy to compute the order of σ .

Theorem 3.12. *Suppose $\sigma = \sigma_1 \sigma_2 \dots \sigma_t$ is a product of t disjoint cycles. Then,*

$$|\sigma| = \text{lcm}(|\sigma_1|, |\sigma_2|, \dots, |\sigma_t|)$$

Remark. If $\sigma = (a_1 a_2 a_3 \dots a_k)$ is a k -cycle, $|\sigma| = k$.

Proof. Let $d_i = |\sigma_i|$ and $d = \text{lcm}(d_1, \dots, d_t)$. Since the cycles are disjoint,

$$\sigma^d = (\sigma_1 \dots \sigma_t)^d = \sigma_1^d \sigma_2^d \dots \sigma_t^d$$

For each i , $d = d_i m_i$ for some m_i . So

$$\begin{aligned} \sigma^d &= \sigma_1^{d_1 m_1} \sigma_2^{d_2 m_2} \dots \sigma_t^{d_t m_t} \\ &= \left(\sigma_1^{d_1}\right)^{m_1} \left(\sigma_2^{d_2}\right)^{m_2} \dots \left(\sigma_t^{d_t}\right)^{m_t} \\ &= e^{m_1} e^{m_2} \dots e^{m_t} \\ &= e \end{aligned}$$

So $|\sigma| \leq d$.

Now, let $l = |\sigma|$. So

$$e = \sigma^l = (\sigma_1 \sigma_2 \dots \sigma_t)^l = \sigma_1^l \dots \sigma_t^l$$

Since the cycles are disjoint, this implies that

$$\sigma_i^l = e$$

for each i . Since $|\sigma_i| = d_i$, we have that $d_i | l$ for all i . So l is a common multiple of d_1, d_2, \dots, d_t . So

$$\text{lcm}(d_1, \dots, d_t) \leq l.$$

Thus,

$$|\sigma| \leq d = \text{lcm}(d_1, d_2, \dots, d_t) \leq l = |\sigma|.$$

Hence, $|\sigma| = \text{lcm}(d_1, \dots, d_t)$. □

Example 3.2.6. Going back to the example, since

$$\sigma = (135)(274)(68),$$

we get

$$|\sigma| = \text{lcm}(3, 3, 2) = 6.$$

Definition 3.11. A 2-cycle is called a transposition.

Example 3.2.7. Consider the cycle (1423). We can write it as a product of transpositions:

$$(1423) = (13)(12)(14)$$

Theorem 3.13. Every permutation can be expressed as a product of transpositions.

Proof. We only need to verify this for cycles. Consider

$$(a_1 a_2 \cdots a_k) = (a_1 a_k)(a_1 a_{k-1})(a_1 a_{k-2}) \cdots (a_1 a_3)(a_1 a_2)$$

□

Example 3.2.8.

$$\begin{aligned} \sigma &= (135)(247)(68) \\ &= (15)(13)(27)(24)(68) \end{aligned}$$

Remark. Factorization in the transposition is not unique.

Example 3.2.9.

$$\begin{aligned} (123) &= (13)(12) \\ &= (13)(23)(12)(13) \\ (1235) &= (15)(13)(12) \\ &= (13)(24)(35)(14)(24) \end{aligned}$$

Observe that (123) is a product of an even number of transpositions whereas (1235) is a product of an odd number of transpositions. So we want to make this into a theorem but we need to prove a lemma first:

Lemma 3.1. If $(id) = e = \sigma_1 \sigma_2 \cdots \sigma_t$, then t is even.

Proof. Since no transposition is the identity, we must have $t > 1$. If $t = 2$, we are done. We can perform induction on t .

e have the following 4 cases for $\sigma_{t-1} \sigma_t$:

	$\sigma_{t-1}\sigma_t$	=	$\sigma'_{t-1}\sigma'_t$
1	$(ab)(ab)$		e
2	$(bc)(ab)$		$(ac)(bc)$
3	$(cd)(ab)$		$(ab)(cd)$
4	$(ac)(ab)$		$(ab)(bc)$

In case 1, since $(ab)(ba) = e$, we remove $\sigma_{t-1}\sigma_t$ from $e = \sigma_1 \cdots \sigma_{t-2}$, and by inducting $t-2$ is even, so t is even.

In cases 2, 3 and 4, we can replace $\sigma_{t-1}\sigma_t$ with $\sigma'_{t-1}\sigma'_t$. In all cases, the last occurrence of a moves left by 1.

Now, we look at $\sigma_{t-2}\sigma_{t-1}$. If in case 1, remove the pair $\sigma_{t-2}\sigma_{t-1}$ and finish by inducting. Else, use cases 2, 3 and 4 to move left one transpositions. We eventually get into case 1. If not, we end with

$$(id) = (ab')\sigma_2\sigma_3 \cdots \sigma_t,$$

but the right hand side sends a to b' , contradicting the fact that this is identity. \square

Theorem 3.14. *No permutation can be expressed as both of odd number of transpositions and even number of transpositions*

Proof. Suppose

$$\sigma = \sigma_1 \cdots \sigma_t = \tau_1 \cdots \tau_l$$

with t even and l odd. Then,

$$\begin{aligned} (id) &= \sigma(\sigma^{-1}) = (\sigma_1 \cdots \sigma_t)(\tau_1 \cdots \tau_l)^{-1} \\ &= \sigma_1 \cdots \sigma_t \tau_1^{-1} \cdots \tau_l^{-1} \\ &= \sigma_1 \cdots \sigma_t \tau_1 \cdots \tau_l \end{aligned}$$

So (id) is a product of $t+l$ transpositions. But this is odd, so a contradiction to the lemma. \square

Definition 3.12. *A permutation of $\sigma \in S_n$ is even if it can be written as an even number of transpositions and odd if it can be written as an odd number of transpositions.*

3.3 Alternating groups

Definition 3.13. *The alternating group A_n is*

$$A_n = \{\sigma \in S_n \mid \sigma \text{ is even}\}.$$

Theorem 3.15. *A_n is a group and a subgroup of S_n .*

Proof. To prove closure, let $\sigma, \tau \in A_n$. So

$$\sigma = \sigma_1 \cdots \sigma_t$$

and

$$\tau = \tau_1 \cdots \tau_l$$

with t, l even. But then

$$\sigma\tau = \sigma_1 \cdots \sigma_t \tau_1 \cdots \tau_l \in A_n$$

since $t + l$ is even. Also, $(id) \in A_n$ by the lemma above.

Finally, if $\sigma \in A_n$ and $\sigma = \sigma_1 \cdots \sigma_t$ with t even, then

$$\begin{aligned} \sigma^{-1} &= (\sigma_1 \cdots \sigma_t)^{-1} \\ &= \sigma_t^{-1} \cdots \sigma_1^{-1} \\ &= \sigma_t \cdots \sigma_1 \in A_n \end{aligned}$$

□

3.4 Group of rigid motions


Recall that D_4 is a set of all rigid motions of the square. We can now think of the rotations as permutations

$$\begin{aligned} R_0 &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \end{pmatrix}, \\ R_{90} &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 1 \end{pmatrix}, \\ R_{180} &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 1 & 2 \end{pmatrix}, \\ &\vdots \\ D &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 4 & 3 & 2 \end{pmatrix} \end{aligned}$$

In fact, we can apply this for any regular polygons:

Definition 3.14. D_n is the group of rigid motions of the regular n -gon.

Note that we are going to write vertices in clockwise fashion:



A rigid motion is determined by 2 pieces of information:

- where 1 is sent to (n choices)
- do the numbers go clockwise or counter clockwise (2 choices)

So the total number of rigid motions is $2n$.

Theorem 3.16. D_n is a group of order $2n$.

Proof. We already showed that $|D_n| = 2n$. We want to show that it's actually a group:

- Clearly, $e \in D_n$ since this is the motion where we leave unchanged.
- If $\sigma, \tau \in D_n$, they are both rigid motion, but so it $\sigma\tau \in D_n$.
- If $\sigma \in D_n$ is a rigid motion, we can always reverse the motion to back the original configuration. So $\sigma^{-1} \in D_n$.

□

Remark. D_n is a subgroup of S_n .

Example 3.4.1. $D_3 = S_3$

We saw that D_4 is not cyclic. In general, D_n is not cyclic. However, D_n can be generated by 2 elements.

Theorem 3.17. For $n \geq 3$, D_n consists of all products of elements r and s such that rotation, r , and reflection, s , satisfy


$$r^n = 1 \text{ and } s^n = 1,$$

Proof. Notice that any rigid motion is a rotation and/or a reflection.

Let $r = \frac{2\pi}{n}$. Then, there are n rotations:

$$(id), r, r^2 = 2\left(\frac{2\pi}{n}\right), r^3 = 3\left(\frac{2\pi}{n}\right), \dots, r^{n-1} = (n-1)\left(\frac{2\pi}{n}\right)$$

Likewise, there are n reflections $s_1, s_2, s_3, \dots, s_n$, where each s_i leaves i fixed. For example, s_1 in D_5 will look like this:



If n is odd, s_i only fixes i , whereas if n is even, s_i fixes two or no elements (e.g., reflection of a square along the vertical axis fix no elements).

Let $s = s_1$. I claim that every element of D_n can be written in terms of r and s . Recall that a rigid motion is determined by (1) where 1 is sent and (2) whether numbers are clockwise or counter clockwise. If 1 is sent to k clockwise, the motion is given by r^{k-1} . If 1 is sent to k in counter clockwise, the motion is given by $r^{k-1}s$. So

$$D_n = \{r^a s^b \mid 0 \leq a \leq n-1, 0 \leq b \leq 1\}.$$

Finally, consider $rsrs$. Then, $rsrs = 1$ so $r(srs) = 1$ and $r^{-1} = srs$. □

Example 3.4.2. Show D_n is not abelian for all $n \geq 3$.

Proof. Suppose that D_n is abelian. We showed that $rsrs = 1$. Since D_n is abelian, $rs^2r = 1$. But $s^2 = 1$. So $r^2 = 1$. But $n \geq 3$ and $|r| = 3 > 2$. □

Example 3.4.3. Use cycle notation to write out all elements of D_5 .

3.5 Lagrange's Theorem

Definition 3.15. Let G be a group with subgroup $H \subseteq G$. The left coset of H with representative $g \in G$ is the set

$$gH = \{gh \mid h \in H\}.$$

The right coset of H is

$$Hg = \{hg \mid h \in H\}.$$

Example 3.5.1. Consider

$$\begin{cases} G = u(8) = \{1, 3, 5, 7\} \\ H = \{1, 5\} \subseteq G \end{cases}$$

Then,

$$\begin{aligned} 1H &= \{1, 5\} \\ 3H &= \{3, 7\} \\ 5H &= \{5, 1\} \\ 7H &= \{7, 3\} \end{aligned}$$

Example 3.5.2. Consider

$$\begin{cases} G = \mathbb{Z}_8 = \{0, 1, 2, 3, \dots, 7\} \\ H = \{0, 4\} \subseteq G \end{cases}$$

Then,

$$\begin{aligned} 0 + H &= \{0, 4\} \\ 1 + H &= \{1, 5\} \\ 2 + H &= \{2, 6\} \\ &\vdots \\ 7 + H &= \{7, 3\} \end{aligned}$$

Remark. If G is abelian, then left and right cosets are same, i.e.

$$gH = \{gh \mid h \in H\} = \{gh \mid h \in H\} = Hg.$$

This is false if G is not abelian.

Example 3.5.3. Consider $G = D_4$ and $T = \{R_0, H\}$, where H is the horizontal flip. Then,

$$R_{90}T = \{R_{90} \circ R_0, R_{90} \circ H\} \neq \{R_0 \circ R_{90}, H \circ R_{90}\} = TR_{90}$$

Lemma 3.2 (Properties of cosets). *Let $H \subseteq G$ be a subgroup. Then,*

- $g \in gH$.
- $gH = H$ iff $g \in H$.
- $g_1H = g_2H$ iff $g_1 \in g_2H$.
- $g_1H = g_2H$ or $g_1H \cap g_2H = \emptyset$.
- $g_1H = g_2H$ iff $g_1^{-1}g_2 \in H$.
- $|g_1H| = |g_2H|$.
- $|gH| = |Hg|$.

Proof.

(1) Since $e \in H$, $ge = g \in gH$.

(2) (\Rightarrow) Suppose $gH = H$. Since $g \in gH$, $g \in H$ because $H = gH$. (\Leftarrow)

Now we want to show that $gH \subseteq H$ given $g \in H$. Since $g \in H$, and H is a subgroup, $gh \in H$ for all $h \in H$. So $gH \subseteq H$. Now, we take $h \in H$. Because $g \in H$, $g^{-1} \in H$, and so is $g^{-1}h$. But then

$$h = g(g^{-1}h) \in gH.$$

Thus, $H \subseteq gH$. So $H = gH$.

(3) (\Rightarrow) Suppose $g_1H = g_2H$. Since $g_1 \in g_1H$, this implies that $g_1 \in g_2H$.

(\Leftarrow) Take $t \in g_1H$ so $t = g_1h$ for some h . And we are given that $g_1 \in g_2H$, so $g_1 = g_2h'$ for some h' . Then, $t = g_h = (g_2h)h' = g_2(hh') \in g_2H$. So $g_1H \subseteq g_2H$. Now, take $t \in g_2H$ so $t = g_2h$ and we know $g_1(h')^{-1} = g_2$. So

$$\begin{aligned} t &= g_2h = (g_1(h')^{-1})h \\ &= g_1[(h')^{-1}h] \in g_1H. \end{aligned}$$

So $g_2H \subseteq g_1H$. Hence, $g_1H = g_2H$.

(4) Since g_1H and g_2H are sets, we can have (a) $g_1H \cap g_2H = \emptyset$, (b) $g_1H = g_2H$, or (c) $g_1H \neq g_2H$ and $g_1H \cap g_2H$. Suppose $x \in g_1H \cap g_2H$. So $x \in g_1H$ implies that $g_1H = xH$. Also, $x \in g_2H$ implies that $g_2H = xH$. So $g_1H = xH = g_2H$. So $g_1H = g_2H$. So (c) cannot happen.

(5) Details are same as the proof of (3)

(6) Define a map

$$f : g_1H \rightarrow g_2H$$

by $f(g_1h) = g_2h$. I claim that f is a bijection.

(one-to-one) If $f(g, h) = f(g, h')$, we have $g_2h = g_2h'$. By cancellation, $h = h'$ so $g_1h = g_1h'$.

(onto) Take $t = g_2h \in g_2H$. Then, $g_1h \in g_1H$ and $f(g_1h) = g_2h = t$. Since f is a bijection,

$$|g_1H| = |g_2H|$$

(7) Same idea but we use a map

$$f : gH \rightarrow Hg$$

by $f(gh) = hg$. □

Theorem 3.18 (Lagrange's Theorem). *If G is a finite group and $H \subseteq G$ is a subgroup, then, $|H||G|$. Also, the number of distinct cosets is $\frac{|G|}{|H|}$.*

Proof. Suppose that there are n distinct left cosets of H in G , say g_1H, g_2H, \dots, g_nH . For each $g \in G$,

$$g \in g_iH = g_iH$$

for some g_i . Thusm,

$$G = g_1H \cup g_2H \cup \dots \cup g_nH.$$

Since cosets are distinct,

$$\begin{aligned} |G| &= |g_1H| + |g_2H| + |g_3H| + \dots + |g_nH| \\ &= |H| + |H| + \dots + |H| \\ &= n|H| \end{aligned}$$

So $|H||G|$ and $\frac{n=|G|}{|H|}$ is the number of distinct cosets. □

Definition 3.16. *The index of H in G is the number of distinct left cosets and denoted $[G : H]$. So $[G : H] = \frac{|G|}{|H|}$.*

Example 3.5.4. Consider

$$\begin{cases} G = u(8) = \{1, 3, 5, 7\} \\ H = \{1, 5\} \subseteq G \end{cases}$$

Then, $[G : H] = 4/2 = 2$.

Note that Lagrange is not true if $|G| = \infty$.

Example 3.5.5. Consider $G = \mathbb{Z}$ and $H = \{2n \mid n \in \mathbb{Z}\}$, the set of even integers. Then, there are only two distinct left cosets: $0 + H = H$ and $1 + H$. So $[G : H] = 2$. However,

$$\frac{|G|}{|H|} = \frac{\infty}{\infty}.$$

Corollary 3.3. For any $g \in G$ (G finite), then $|g| \mid |G|$.

Proof. For any $g \in G$, $|g| = |\langle g \rangle|$. Since $\langle g \rangle$ is a subgroup of G , $|g| \mid |G|$. \square

Corollary 3.4. If $|G| = p$ is a prime, then G must be cyclic and is generated by any non-identity element.

Proof. Let $g \in G$ with $g \neq e$. Then, $1 < |g| \mid |G| = p$ so $|g| = p$, i.e. $\langle g \rangle = G$. \square

Roughly this says all cyclic groups of order p are the same as \mathbb{Z}_p .

Corollary 3.5. Let H and K be subgroup of G such that $K \subset H \subset G$. Then,

$$[G : K] = [G : H][H : K]$$

Proof.

$$[G : K] = \frac{|G|}{|K|} = \frac{|G|}{|H|} \times \frac{|H|}{|K|} = [G : H][H : K]$$

\square

Note that the converse of Lagrange's theorem is false, i.e. if $d \mid |G|$, then G has a subgroup of order d .

Example 3.5.6. Consider an alternating group A_4 . Then,

$$|A_4| = 4!/2 = 12$$

Note that $6 \mid 12$, but we will show that A_4 has no subgroup of order 6.

Suppose $H \subseteq A_4$ was a subgroup of order t . So $[A_4 : H] = 12/6$. For all $g \in A_4$, $gH = Hg$. So

1. if $g \in H$, then $gH = H = Hg$
2. if $g \notin H$, then $gH \neq H$.

Since $[A_4 : H] = 2$, this means $A_4 = H \cup gH$ but we also would have $Hg \neq H$ and $A_4 = H \cup Hg$. Thus,

$$H \cup gH = H \cup Hg \implies gH = Hg,$$

since those unions are disjoint. So

$$gHg^{-1} = H,$$

for all $g \in A_4$.

Note that the group A_4 has 8 three cycles:

$$(123), (132), (124), (142), (134), (143), (234), (243).$$

So H has at least one of three cycles, say $(123) \in H$. This implies that $(123)^{-1} = (132) \in H$. Then,

$$(124)(123)(124)^{-1} = (243) \in H, (243)(123)(243)^{-1} = (142) \in H,$$

But then H has at least 7 elements:

$$(id), (123), (132), (243), (243)^{-1}, (142), (142)^{-1}$$

But $|14| = 6$. So H does not exist.

4 Fermat's little theorem

4.1 Fermat's little theorem

Definition 4.1. Euler's ϕ -function $\phi : \mathbb{N} \rightarrow \mathbb{N}$ is defined as

$$\begin{aligned}\phi(1) &= 1 \\ \phi(n) &= \{m \mid 1 \leq m < n, \gcd(m, n) = 1\} = |U(n)|.\end{aligned}$$

Theorem 4.1. Let a and n be integers with $n > 1$ and $\gcd(a, n) = 1$. Then,

$$a^{\phi(n)} \equiv 1 \pmod{n}$$

Proof. Since $\gcd(a, n) = 1$, this means that $a \in U(n)$. Then, $|a| |U(n)| = \phi(n)$. So

$$\phi(n) = |a|l$$

and

$$a^{\phi(n)} = a^{|a|l} = \left(a^{|a|}\right)^l = 1$$

in $U(n)$. As a result,

$$a^{\phi(n)} \equiv 1 \pmod{n}.$$

□

Theorem 4.2. Let p be prime. Then, for all integers a ,

$$a^p \equiv a \pmod{p}$$

Proof. If $p|a$, then $a^p \equiv a \pmod{p}$. If $p \nmid a$, then $\gcd(a, p) = 1$. By Euler's Theorem,

$$a^{\phi(p)} \equiv 1 \pmod{p}.$$

But p prime means $\phi(p) = p - 1$. So

$$a^{p-1} \equiv 1 \pmod{p} \implies a^p \equiv a \pmod{p}.$$

□

Example 4.1.1. Consider $a = 32$ and $p = 7$. Then,

$$32^7 \equiv 32 \pmod{7}.$$

This is extremely useful for modular computation.

5 Isomorphisms

5.1 Isomorphisms

Informally, two sets are isomorphic if they are the same, but just have different labels.

Definition 5.1. Let G and H be groups with operations $*$ and \circ , respectively. Then, G is isomorphic to H if there is a bijection $\phi : G \rightarrow H$ that preserves the operation, i.e.,

$$\phi(a * b) = \phi(a) \circ (\phi b),$$

where $*$ is an operation in G and \circ is an operation in H . Then, we write $G \simeq H$.

Example 5.1.1. Consider

$$\begin{aligned} G = u(8) &= \{1, 3, 5, 7\} \\ H = u(12) &= \{1, 5, 7, 11\} \end{aligned}$$

Prove that $u(8) \simeq u(12)$.

Define our map $\phi : u(8) \rightarrow u(12)$ by

$$\begin{aligned} 1 &\rightarrow 1 \\ 3 &\rightarrow 5 \\ 5 &\rightarrow 7 \\ 7 &\rightarrow 11 \end{aligned}$$

This is a bijection. To check the operation is preserved, we can compare cayley tables:

	1	3	5	7		1	3	5	7
1	1	3	5	7	1	1	5	7	11
3	3	1	7	5	5	5	1	11	7
5	5	7	1	3	7	7	11	1	5
7	7	5	3	1	11	11	7	5	1

5.2 Cyclic groups

Theorem 5.1. (A) Every infinite cyclic group is isomorphic to \mathbb{Z} . (B) Every finite cyclic group G with $|G| = n$ is isomorphic to \mathbb{Z}_n .

Proof. (A) Let $G = \langle a \rangle = \{a^i \mid i \in \mathbb{Z}\}$. Define a map $\phi : \mathbb{Z} \rightarrow G$ by $\phi(i) = a^i$. Then this is a bijection:

- (onto) For any $g \in G$, $g = a^i$ for some $i \in \mathbb{Z}$. Then, $\phi(i) = a^i = g$.
- (one-to-one) Suppose that $\phi(k_1) = a^{k_1} = a^{k_2} = \phi(k_2)$. Since G is infinite, $k_1 = k_2$. So ϕ is one-to-one.

Let $i, j \in \mathbb{Z}$. Then,

$$\phi(i+j) = a^{i+j} = a^i a^j = \phi(i)\phi(j).$$

(B) We are given that $G = \{a^0, a^1, \dots, a^{n-1}\} = \langle a \rangle$. Define

$$\phi : \mathbb{Z}_n \rightarrow G$$

by $\phi(i) = a^i$. This is clearly a bijection. Given $i, j \in \mathbb{Z}_n$, suppose $i+j = k \in \mathbb{Z}_n$. Then,

$$\phi(i+j) = \phi(k) = a^k = a^{i+j} = a^i a^j = \phi(i)\phi(j)$$

□

Theorem 5.2 (Properties of isomorphisms). *If $\phi : G \rightarrow H$ is an isomorphism, then*

1. $\phi(e_G) = e_H$.
2. $\phi(g)^{-1} = \phi(g^{-1})$
3. $|G| = |H|$
4. If G is abelian, so is H
5. If G is cyclic, then so is H
6. If G has a subgroup of order m , then so does H
7. For all $g \in G$, $|g| = |\phi(g)|$

Proof. (1) We know $e_G e_G = e_G$. So $\phi(e_G) = \phi(e_G e_G) = \phi(e_G)\phi(e_G)$. Then,

$$e_H \phi(e_G) = \phi(e_G)\phi(e_G).$$

By the cancellation property, $e_H = \phi(e_G)$.

(2) Observe that

$$e_H = \phi(e_G) = \phi(gg^{-1}) = \phi(g)\phi(g^{-1}).$$

Since inverses are unique, $\phi(g)^{-1} = \phi(g^{-1})$.

(3) Since ϕ is a bijection, $|G| = |H|$.

(4) Let $h_1, h_2 \in H$. Then,

$$\begin{aligned} h_1 h_2 &= \phi(g_1)\phi(g_2) \\ &= \phi(g_1 g_2) \\ &= \phi(g_2 g_1) \\ &= \phi(g_2)\phi(g_1) = h_2 h_1 \end{aligned}$$

(5) Same type of proof

(6) Homework

(7) Let $a = |g|$, i.e. $g^a = e_G$. Then,

$$e_H = \phi(e_G) = \phi(g^a) = \phi(g) \cdots \phi(g) = \phi(g)^a$$

So $b = |\phi(g)| \leq a$.

Now, let $b = |\phi(g)|$. So

$$\phi(g)^b = \phi(g \cdots g) = \phi(g^b) = e_H = \phi(e_G).$$

Since ϕ is one-to-one and $g^b = e_G$, so $a \leq b$. Therefore, $a \leq b \leq a$. So $a = b$. \square

Example 5.2.1. D_4 and \mathbb{Z}_8 are not isomorphic because \mathbb{Z}_4 is cyclic whereas D_4 is not.

Example 5.2.2. $u(8)$ and \mathbb{Z}_4 are not isomorphic because $|1| = 4$ in \mathbb{Z}_4 but every element has order to in $u(8)$.

Example 5.2.3. Let G be any finite group of order p (prime). Then, $G \simeq \mathbb{Z}_p$ because G is cyclic with order p .

Fundamental problem of finite group theory. Classify all finite groups up to isomorphism (memorize this table):

n	all groups of order n up to isomorphism
1	$\{e\}$
2	\mathbb{Z}_2
3	\mathbb{Z}_3
4	$\mathbb{Z}_4, u(8) = \mathbb{Z}_2 \times \mathbb{Z}_2$
5	\mathbb{Z}_5
6	\mathbb{Z}_6, S_3
7	\mathbb{Z}_7

5.3 Cayley's Theorem

Theorem 5.3. Every group is isomorphism to a group of permutations.

Example 5.3.1. Consider

$$U(8) = \{1, 3, 5, 7\}.$$

For each $g \in U(8)$, we acn define a bijection

$$\lambda_g : U(8) \rightarrow U(8)$$

by letting $\lambda_g(x) = gx$. So λ_3 is defined by $\lambda_3(x) = 3x$:

$$1 \rightarrow 3 \cdot 1 = 3$$

$$3 \rightarrow 3 \cdot 3 = 1$$

$$5 \rightarrow 3 \cdot 5 = 7$$

$$7 \rightarrow 3 \cdot 7 = 5$$

So we can write λ_3 as a permutation:

$$\begin{pmatrix} 1 & 3 & 5 & 7 \\ 3 & 1 & 7 & 5 \end{pmatrix}$$

Likewise,

$$\lambda_1 = \begin{pmatrix} 1 & 3 & 5 & 7 \\ 1 & 3 & 5 & 7 \end{pmatrix}, \lambda_5 = \begin{pmatrix} 1 & 3 & 5 & 7 \\ 5 & 7 & 1 & 3 \end{pmatrix}, \lambda_7 = \begin{pmatrix} 1 & 3 & 5 & 7 \\ 7 & 5 & 3 & 1 \end{pmatrix}$$

Example 5.3.2. Let $\bar{U}(8) = \{\lambda_1, \lambda_3, \lambda_5, \lambda_7\}$ as a set of bijections from $U(8)$ to itself. Then, $U(8) \simeq \bar{U}(8)$ as groups.

Proof. For each $g \in G$, define $\lambda_g : G \rightarrow G$ given by $\lambda_g(x) = gx$. □

Theorem 5.4. Each $\lambda_g : G \rightarrow G$ is a bijection, i.e., a permutation of elements of G .

Proof. (surjective) Let $g \in G$. Since $g^{-1} \in G$, so is $g^{-1}G$. Then,

$$\lambda_g(g^{-1}G) = g(g^{-1}G) = (gg^{-1})G = G.$$

(injective) Suppose $\lambda_g(x) = gx = gy = \lambda_g(y)$. But, by cancellation, $gx = gy$ implies $x = y$. So λ_g is injective.

Therefore,

$$H = \{\lambda_g \mid g \in G\} = \bar{G}.$$

□

Theorem 5.5. H is a subgroup of all the permutations of the elements of G under composition.

Proof. (closed) Take $\lambda_g, \lambda_h \in H$. Then,

$$\begin{aligned} (\lambda_g \circ \lambda_h)(x) &= \lambda_g(\lambda_h(x)) \\ &= \lambda_g(hx) \\ &= ghx \end{aligned}$$

But $g, h \in G$ so $gh \in G$ and $\lambda_{gh} \in H$ and $\lambda_{gh}(x) = ghx$. So

$$(\lambda_g \lambda_h)(x) = ghx = \lambda_{gh}(x).$$

(identity) Since $e \in G$, $\lambda_e \in H$. For all $x \in G$, $\lambda_e(x) = ex = x$. So λ_e is the identity function.

(inverse) Consider $\lambda_g \in H$. Since $g \in G$, $g^{-1} \in G$, and so $\lambda_{g^{-1}} \in H$. Then, for all $x \in G$,

$$(\lambda_g \circ \lambda_{g^{-1}})(x) = g(g^{-1}(x)) = x = \lambda_e(x)$$

□

Theorem 5.6. $G \simeq \bar{G} = H$

Proof. Define $\phi : G \rightarrow H$ by $g \rightarrow \lambda_g$. We check that this is an isomorphism.

(surjective) If $\lambda_g \in H$, then $g \in G$, and $\phi(g) = \lambda_g$.

(injective) Suppose $\phi(g) = \lambda_g = \lambda_h = \phi(h)$. Since $e \in G$, we have

$$\lambda_g(e) = ge = he = \lambda_h(e).$$

So $g = h$ (ϕ preserves operation).

So

$$\phi(g) \circ \phi(h) = (\lambda_g \circ \lambda_h) = \lambda_{gh} = \phi(gh)$$

So $H \simeq G$, as desired. \square

5.4 Direct Products

Let $(G, *)$ and (H, \circ) be two groups.

Definition 5.2 (External direct product). $G \times H = \{(g, h) \mid g \in G \text{ and } h \in H\}$ is a group under the operation

$$(g_1, h_1)(g_2, h_2) = (g_1 * g_2, h_1 \circ h_2)$$

Example 5.4.1. Consider $\mathbb{Z}_2 = \{0, 1\}$. Then,

$$\mathbb{Z}_2 \times \mathbb{Z}_2 = \{(0, 0), (0, 1), (1, 0), (1, 1)\}$$

Note that $|\mathbb{Z}_2 \times \mathbb{Z}_2| = 4$ but $\mathbb{Z}_2 \times \mathbb{Z}_2 \neq \mathbb{Z}_4$ since $\mathbb{Z}_2 \times \mathbb{Z}_2$ has no element of order 4.

Example 5.4.2. $\mathbb{Z}_2 \times \mathbb{Z}_3 = \{(0, 0), (0, 1), (0, 2), (1, 0), (1, 1), (1, 2)\}$

Theorem 5.7. If $(g, h) \in G \times H$ and $|g| = r$ and $|h| = s$, then $|(g, h)| = \text{lcm}(r, s)$.

Example 5.4.3. In \mathbb{Z}_3 , $|1| = 3$ and in \mathbb{Z}_5 , $|1| = 5$. So in $\mathbb{Z}_3 \times \mathbb{Z}_5$,

$$|(1, 1)| = 15 = \text{lcm}(3, 5)$$

So $\mathbb{Z}_3 \times \mathbb{Z}_5 \simeq \mathbb{Z}_{15}$

Theorem 5.8. $\mathbb{Z}_n \times \mathbb{Z}_m \simeq \mathbb{Z}_{nm}$ if and only if $\text{gcd}(m, n) = 1$.

Proof. (\Rightarrow) Suppose $\mathbb{Z}_n \times \mathbb{Z}_m \simeq \mathbb{Z}_{nm}$, but $\text{gcd}(m, n) = d > 1$. So

$$\frac{mn}{d} = m \left(\frac{n}{d} \right) = \left(\frac{m}{d} \right) n < mn.$$

But then, for all $(a, b) \in \mathbb{Z}_n \times \mathbb{Z}_m$,

$$\underbrace{(a, b) + \cdots (a, b)}_{\frac{mn}{d}} = \left(an \left(\frac{m}{d} \right), bm \left(\frac{n}{d} \right) \right) = (0, 0)$$

So every element in $\mathbb{Z}_n \times \mathbb{Z}_m$ has order less than mn/d but \mathbb{Z}_{nm} has at least 1 element of order mn , which allows a contradiction to arise.

(\Leftarrow) $|1| = n$ in \mathbb{Z}_n and $|1| = m$ in \mathbb{Z}_m . Since $\gcd(m, n) = 1$ and $\text{lcm}(m, n) = mn$,

$$|(1, 1)| = \text{lcm}(m, n) = mn,$$

i.e., $\langle (1, 1) \rangle = \mathbb{Z}_{nm}$ since $|\mathbb{Z}_n \times \mathbb{Z}_m| = mn$. \square

Remark. $\gcd(a, b) \cdot \text{lcm}(a, b) = ab$.

Given a group G , can we find subgroups H and K such that $G = H \times K$?

Definition 5.3. Let G be a group with two subgroups H and K . Then, G is an internal direct product of H and K if

- $G = HK = \{hk \mid h \in H, k \in K\}$
- $H \cap K = \{e\}$
- $hk = kh$ for all $k \in K$ and $h \in H$.

Example 5.4.4. Let $G = U(8) = \{1, 3, 5, 7\}$, $H = \{1, 3\}$ and $K = \{1, 5\}$. Then,

- $HK = \{1 \cdot 1, 1 \cdot 5, 3 \cdot 1, 3 \cdot 5\} = \{1, 5, 3, 7\}$.
- $H \cap K = \{e\} = \{1\}$
- Since $U(8)$ is abelian, $hk = kh$ for all $k \in K$ and $h \in H$.

So $U(8)$ is the internal direct product of H and K .

Theorem 5.9. If G is the internal direct product of H and K , then $G = H \times K$.

Example 5.4.5. $U(8) = H \times K$

Proof. Observe that every $g \in G$ can be written as $g = hk$ for some $h \in H, k \in K$. So we define $\phi : G \rightarrow H \times K$, by $\phi(g) = (h, k)$. First, we need to show that ϕ is well defined. Suppose $g = h_1k_1 = h_2k_2$. Then, $h_2^{-1}h_1 = k_2k_1^{-1}$. Since $h_2^{-1}h_1 \in H$ and $k_2k_1^{-1} \in K$,

$$h_2^{-1}h_1 = k_2k_1^{-1} \in H \cap K = \{e\}.$$

So $h_1 = h_2$ and $k_1 = k_2$ so $g = hk$ is the unique way of factoring g .

We claim that ϕ is one-to-one and onto (check this fact) so we just need to show that the operation is preserved.

$$\begin{aligned} \phi(g_1g_2) &= \phi(h_1k_1h_2k_2) \\ &= \phi(h_1h_2k_1k_2) \\ &= (h_1h_2, k_1k_2) \\ &= (h_1, k_1)(h_2, k_2) \\ &= \phi(g_1)\phi(g_2) \end{aligned}$$

\square

Example 5.4.6. Let $T = \{2^n 3^m \mid n, m \in \mathbb{Z}\} \subseteq \mathbb{Q}$. Prove that $T \simeq \mathbb{Z} \times \mathbb{Z}$.

Proof. So we want to show that T is an internal direct product of $T_1 = \{2^n \mid n \in \mathbb{Z}\}$ and $T_2 = \{3^m \mid m \in \mathbb{Z}\}$.

- Note that $T_1 \cap T_2 = \{1\} = \{2^0\} = \{3^0\}$
- Also, note that $t_1 t_2 = t_2 t_1$ for all $t_1 \in T_1$ and $t_2 \in T_2$, since T is abelian.
- Let $t \in T$. So $t = 2^n 3^m$. But $2^n \in T_1$ and $3^m \in T_2$, so $T = T_1 T_2$.

So T is an internal direct product of T_1 and T_2 . Then, we apply the previous theorem to conclude that $T \simeq T_1 \times T_2$. \square

Remark. For finite groups, we can prove cancellation:

$$G \times H = G \times K \implies H = K$$

If G is not finite, this is false. Let $G = \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z} \times \cdots$. Let $H = \mathbb{Z}$ and $K = \mathbb{Z} \times \mathbb{Z}$. Then, $H \times G = K \times G$ but $H \neq K$.

6 Factor groups

6.1 Factor groups

Recall that if $H \subseteq G$ is a subgroup, then we have left and right cosets: gH and Hg . Hereafter, we write $\frac{G}{H}$ to denote the set of all distinct left cosets.

Example 6.1.1. Consider $G = \mathbb{Z}_8$ and $H = \{0, 4\}$. Then,

$$\frac{G}{H} = \{0 + H, 1 + H, 2 + H, 3 + H\}$$

So $\frac{G}{H}$ is a set. Does $\frac{G}{H}$ have extra structure? In particular, is $\frac{G}{H}$ a group?

Definition 6.1. $\frac{G}{H}$ is a factor group (or quotient group) if $\frac{G}{H}$ is a group.

So how should an operation on $\frac{G}{H}$ be defined? Our first guess is to use

$$(aH)(bH) = (abH),$$

We have defined our operation in terms of coset representative but the problem is that there are many a' such that $aH = a'H$.

Example 6.1.2. Consider the following in $\frac{\mathbb{Z}_8}{H}$:

$$\begin{aligned} (1 + H) &= (5 + H) \\ (2 + H) &= (6 + H) \\ (1 + H) + (2 + H) &= (3 + H) \\ (5 + H) + (6 + H) &= (11 + H) = (3 + H) \end{aligned}$$

Example 6.1.3. Consider

$$\begin{aligned} S_3 &= \{(1), (12), (13), (23), (132), (123)\} \\ N &= \{(1), (12)\} \end{aligned}$$

Then,

$$\frac{S_3}{N} = \{(1)N, (123)N, (23)N\}$$

In this case, $(123)N$ and $(13)N$ are same cosets but $(123)N(23)N$ and $(13)N(23)N$ are different:

$$\begin{aligned} (123)N(23)N &= (123)(23)N = (12)N \\ (13)N(23)N &= (13)(23) = (132)N \end{aligned}$$

So when is $\frac{G}{H}$ a group? It depends on H . H needs an extra property!

Definition 6.2. A subgroup $N \subseteq G$ is a normal subgroup if $gN = Ng$ for all $g \in G$.

Theorem 6.1. If G is abelian, every subgroup is normal.

Proof. For any $g \in G$ and subgroup $N \subseteq G$,

$$gN = \{gn \mid n \in N\}$$

Since G is abelian, $gn = ng$ for all $n \in N$. Thus,

$$gN = Ng,$$

and N is normal. □

Example 6.1.4. Consider

$$N = \{(1), (12)\} \subseteq S_3.$$

Then, N is not normal since

$$(123)N \neq N(123).$$

Theorem 6.2 (Normal subgroup test). *Let $N \subseteq G$ be a subgroup. Then, following are equivalent:*

1. N is normal in G
2. $gNg^{-1} \subseteq N$ for all $g \in G$
3. $gNg^{-1} = N$ for all $g \in G$

Proof. (1 \implies 2) Suppose N is normal, i.e., $gN = Ng$ for all $g \in G$. So for any $g \in G$, there exists n and $n' \in N$ such that $gn = n'g$. So, for any $g \in G$ and $n \in N$,

$$gn g^{-1} n'$$

for some $n' \in N$. But this means

$$gNg^{-1} = \{gn g^{-1} \mid n \in N\} \subseteq N$$

(2 \implies 3) It suffices to show that $N \subseteq gNg^{-1}$ for $g \in G$. Let $n \in N$ and let $g \in G$. By (2),

$$(g^{-1})N(g^{-1})^{-1} \subseteq N.$$

Thus,

$$g^{-1}n(g^{-1})^{-1} \in N$$

but then

$$n = g(g^{-1}n(g^{-1})^{-1})g^{-1} \in gNg^{-1}$$

(3 \implies 1) We are given $gNg^{-1} = N$ for all $g \in G$. So

$$gn g^{-1} = n'$$

for some $n \in N$ and $n' \in N$. Then,

$$gn = n'g.$$

So for any $g \in G, n \in N$, $gn \in gN$ is also in Ng since $ng = n'g$. So $gN \subseteq Ng$. By a similar argument $Ng \subseteq gN$. □

Remark. $gN = Ng$ does not imply $gn = ng$ for all $g \in G$ and $n \in N$. It means that there exists n and n' such that $gn = n'g$.

Theorem 6.3. *Let G be a group with a normal subgroup N . Then,*

$$G/N = \{\text{all distinct left cosets}\}$$

is a group where the operation is

$$(aN)(bN) = (ab)N$$

Definition 6.3. G/N is called the factor group or quotient group.

Proof. (Step 1) First, we show that the operation is closed and well defined. The operation is closed since $abN \in G/N$. To check for well definedness, we need to show that if $aN = a'N$ and $bN = b'N$, then

$$abN = a'b'N.$$

Let $t \in abN$, so $t = abn$. Since $bn \in bN = b'N$, there is n_2 such that

$$bn = b'n_2.$$

Also, since N is normal,

$$b'N = Nb'.$$

So $b'n_2 = n_3b'$ for some $n_3 \in N$. Then,

$$t = abn = a(b'n_2) = a(n_3b') = (an_3)b'.$$

So $an_3 = aN = a'N$. So $an_3 = a'n_4$ for some $n_4 \in N$. Then,

$$t = (an_3)b' = (a'n_4)b' = a'(n_4b').$$

Since $n_4b' \in Nb' = b'N$, we have

$$n_4b' = b'n_5$$

for $n_5 \in N$ since N is normal. So

$$t = a'(n_4b') = a'b'n_5 \in a'b'N.$$

Thus, $abN \subseteq a'b'N$. A similar argument can be made for the other direction.

(Step 2 - identity) eN is the identity.

(Step 3 - inverse) If $gN \in G/N$, $g^{-1}N \in G/N$ and

$$(gN)(g^{-1}N) = (gg^{-1})N = eN.$$

(Step 4 - associativity) Since operations in G is associative, this operation is associative. \square

Example 6.1.5. Consier the following normal set:

$$D = \{R_0, R_{180}\} \subseteq D_4.$$

Then,

$$D_4/N = \{R_0N, R_{90}N, HN, D_1N\},$$

where

$$R_0N = \{R_0, R_{180}\}$$

$$R_{90}N = \{R_{90}, R_{270}\}$$

$$HN = \{H, V\}$$

$$D_1N = \{D_1, D_2\}$$

Then, we can make the following operation table:

	R_0N	$R_{90}N$	HN	D_1N
R_0N	R_0N	$R_{90}N$	HN	D_1N
$R_{90}N$	$R_{90}N$	R_0N	D_1N	HN
HN	HN	D_1N	R_0N	$R_{90}N$
D_1N	D_1N	HN	$R_{90}N$	R_0N

Note that properties of G/N are related to property of G and N .

Theorem 6.4. *Suppose $N \subseteq G$ is normal. Then, the quotient G/N is abelian if and only if $ghg^{-1}h^{-1} \in N$ for all $g, h \in G$.*

Proof. G/N is abelian if and only if

- $(gN)(hN) = (hN)(gN)$
- $ghN = (hg)N$ for all $h, g \in N$
- $(gh)(hg)^{-1} \in N$
- $(gh)(g^{-1}h^{-1}) \in N$

□

Theorem 6.5 (Cayley's theorem for finite abelian groups). *Let G be a finite abelian group. If p is a prime such that $p \mid |G|$, then G has an element of order p (a partial converse of Lagrange's theorem)*

Proof. If $n = 2 \simeq \mathbb{Z}_2$. Now, $2 \mid |G| = 2$ is the only prime that divides 2 and \mathbb{Z}_2 has an element of order 2.

Now, I claim that if $\frac{G}{N}$ has an element of order m and $|G| < \infty$, then G has an elemnt of order m . First, suppose that gN has order m . So

$$(gN)^m = g^mN = eN.$$

If $d = |g|$, then $(gN)^d = g^dN = eN$. This means that

$$m \mid d \iff d = mk.$$

But then, $|g| = mk$ implies $|g^k| = m$.

Suppose $n = |G| > 2$. Let $e \neq x \in G$ and let $|x| = qm \geq 2$ where q is a prime. If $q = p$, then $|x^m| = p$, and we are done. If $q \neq p$, then $\langle x^m \rangle$ is a cyclic group of order q . This is normal in G since G is abelian, so

$$\bar{G} = \frac{G}{\langle x^m \rangle}$$

is a group with $|\bar{G}| = |G|/q = n/q < n$. Since $p \nmid |G| = n$ and $p \neq q$ and $p \nmid n/q$, so $p \nmid |\bar{G}|$. By definition, \bar{G} has an element of order p . By the claim, we then have G has an element of order p . \square

Example 6.1.6. Observe that

$$|U(23)| = 22.$$

Since $2|22$ and $11|22$, this group has elements of order 2 and 11.

Example 6.1.7. Since $|U(43)| = 42 = 2 \times 3 \times 7$, $U(43)$ has an element of order 2, 3, 7.

6.2 Simple groups

Definition 6.4. A group G is a simple group if it does not have any non-trivial normal subgroups.

Theorem 6.6. If p is prime, \mathbb{Z}_p is simple.

Proof. If p is prime, the only subgroups of $\mathbb{Z}\mathbb{Z}_p$ are the trivial subgroups. So it can't have nontrivial normal subgroups. \square

Theorem 6.7. For all $n \geq 3$, A_n is simple.

Proof. We provide an outline of the proof instead:

1. For all $n \geq 3$, A_n is generated by 3 cycles.
2. If N is a normal subgroup in A_n with $n \geq 3$ and if N contains a 3 cycle, then $N = A_n$.
3. If $n \geq 5$, any normal subgroup $N \leq A_n$ contains a 3 cycle.

Then, steps 2 and 3 imply the theorem. \square

Example 6.2.1. If $n = 3$, A_3 is also simple since $|A_3| = \frac{3!}{2} = 3$, so $A_3 \simeq \mathbb{Z}_3$.

Example 6.2.2. If $n = 4$, then A_n has a normal subgroup:

$$N = \{(1), (12)(34), (13)(24), (14)(23)\} \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$$

N does not have a 3-cycle.

Remark. While A_n is simple if $n \geq 5$, S_n is never simple except if $n = 1$ or $n = 2$.

Theorem 6.8. *For all $n \geq 3$, A_3 is normal in S_n .*

Proof. We want to show that

$$\sigma A \sigma^{-1} \subseteq A_n$$

for all $\sigma \in S_n$.

If $\sigma = \sigma_1 \sigma_2 \cdots \sigma_{2l}$ is an even permutation, then so is σ^{-1} . Let $\tau = \tau_1 \cdots \tau_{2m} \in A_n$. Then, $\sigma \tau \sigma^{-1}$ consists of even number of permutations. So

$$\sigma \tau \sigma^{-1} \in A_n.$$

If σ is odd, then $\sigma \tau \sigma^{-1}$ still consists of even number of permutations. So

$$\sigma \tau \sigma^{-1} \in A_n.$$

□

Corollary 6.1. $\frac{S_n}{A_n} \simeq \mathbb{Z}_2$

Proof. Note that

$$\left| \frac{S_n}{A_n} \right| = \frac{|S_n|}{|A_n|} = \frac{n!}{\left(\frac{n!}{2}\right)} = 2$$

Since there is only one group with order 2, $\frac{S_n}{A_n} \simeq \mathbb{Z}_2$. □

Example 6.2.3. If G/N is abelian and N abelian, is G abelian? No, $S_3/A_3 \simeq \mathbb{Z}_2$ is abelian and $A_3 \simeq \mathbb{Z}_3$ is abelian but S_3 is not.

7 Homomorphism

7.1 Homomorphism

Definition 7.1. Let $(G, *)$ and (H, \circ) be two groups. A homomorphism is a function, $\phi : G \rightarrow H$ such that

$$\phi(a * b) = \phi(a) \circ \phi(b),$$

for all $a, b \in G$.

Note that we drop the bijection condition of an isomorphism.

Example 7.1.1. Define a map $\phi : \mathbb{Z} \rightarrow \mathbb{Z}_n$ by

$$\phi(a) = a \mod n.$$

This is an isomorphism since

$$\begin{aligned}\phi(a + b) &= (a + b) \mod n \\ &= (a \mod n) + (b \mod n) \\ &= \phi(a) + \phi(b)\end{aligned}$$

Example 7.1.2. Consider $\phi : \mathbb{R}^* \rightarrow \mathbb{R}^*$ defined by

$$\phi(x) = |x|.$$

This is an isomorphism because

$$\phi(ab) = |ab| = |a||b| = \phi(a)\phi(b)$$

Example 7.1.3. Consider $\phi : \text{GL}_2(\mathbb{R}) \rightarrow \mathbb{R}^*$ defined by

$$\phi(A) = \det(A)$$

This is an isomorphism because

$$\begin{aligned}\phi(AB) &= \det(AB) \\ &= \det(A)\det(B) \\ &= \phi(A)\phi(B)\end{aligned}$$

Theorem 7.1. Let $\phi : G \rightarrow H$ be any group homomorphism. Then,

1. $\phi(e_G) = e_H$
2. $\phi(a)^n = \phi(a^n)$ for all $n \in \mathbb{Z}$
3. $\phi(a)^{-1} = \phi(a^{-1})$

4. If $N \subseteq G$ is a subgroup, then

$$\phi(N) = \{\phi(n) \mid n \in N\}$$

is a subgroup of H

5. If K is any group of H , then

$$\phi^{-1}(K) = \{g \in G \mid \phi(g) \in K\}$$

is a subgroup of G . If K is normal, so is $\phi^{-1}(K)$.

Proof. (1) Note that $e_G = e_G e_G$. So

$$\phi(e_G) = \phi(e_G e_G) = \phi(e_G) \phi(e_G)$$

We multiply both sides by $\phi(e_G)^{-1}$, we have

$$e_H = \phi(e_G)^{-1} \phi(e_G) = \phi(e_G)^{-1} \phi(e_G) \phi(e_G) = \phi(e_G)$$

(2) If $n = 0$, this is just 1. So suppose $n \geq 1$:

$$\phi(a)^n = \underbrace{\phi(a) \cdots \phi(a)}_n = \phi(a^n)$$

by homomorphism property. If $n \leq -1$, then $-n \geq 1$ so

$$\phi(a)^{-n} = \phi(a^{-n})$$

On the other hand,

$$\phi(a)^n (\phi(a)^n)^{-1} = \phi(a)^n \phi(a)^{-n} = e_H$$

Also, since $a^n a^{-n} = e_G$, we have

$$\phi(e_G) = e_H = \phi(a^n) \phi(a^{-n}).$$

So

$$\phi(a^n) \phi(a^{-n}) = \phi(a)^n \phi(a)^{-n}$$

Finally, we can cancel $\phi(a^{-n})$ with $\phi(a)^{-n}$ to obtain

$$\phi(a^n) = \phi(a)^n.$$

(3) This is property 2 with $n = -1$.

(4) Apply the subgroup test to

$$\phi(N) = \{\phi(n) \mid n \in N\}$$

- (identity) Since $e_G \in N$, and $\phi(e_G) = e_H$, so $e_H \in \phi(N)$.

- (closure) Let $a, b \in \phi(N)$. Then,

$$\begin{aligned} a &= \phi(n_1), \\ b &= \phi(n_2), \end{aligned}$$

for some $n_1, n_2 \in N$. Then,

$$ab = \phi(n_1)\phi(n_2) = \phi(n_1n_2).$$

But $n_1n_2 \in N$. Since $n_1n_2 \in N$, so $ab \in \phi(N)$.

- (inverse) Let $a \in \phi(N)$, so $a = \phi(n_1)$ for some $n_1 \in N$. Then,

$$a^{-1} = \phi(n_1)^{-1} = \phi(n_1^{-1}).$$

But $n_1^{-1} \in N$ since $n_1 \in N$. So $a^{-1} \in \phi(N)$.

- (5) We use a subgroup test to show that

$$\phi^{-1}(K) = \{g \in G \mid \phi(a) \in K\}$$

is a subgroup.

- (identity) Since $e_H \in K$ and $\phi(e_G) = e_H$, $e_G \in \phi^{-1}(K)$.
- (closure) take $a, b \in \phi^{-1}(K)$. So $\phi(a)$ and $\phi(b) \in K$. So $\phi(a)\phi(b) \in K$ since K is closed. But since $\phi(a)\phi(b) = \phi(ab)$, we have $ab \in \phi^{-1}(K)$.
- (inverse) Take $a \in \phi^{-1}(K)$. So $\phi(a) \in K$ and $\phi(a)^{-1} = \phi(a^{-1}) \in K$. So $a^{-1} \in \phi^{-1}(K)$. Now, assume K is normal. We want to show that

$$g\phi^{-1}(K)g^{-1} \subseteq \phi^{-1}(K).$$

Take $t \in g\phi^{-1}(K)g^{-1}$ so $t = gag^{-1}$ with $\phi(a) \in K$. Then,

$$\begin{aligned} \phi(t) &= \phi(gag^{-1}) \\ &= \phi(g)\phi(a)\phi(g^{-1}) \\ &= \phi(g)\phi(a)\phi(g)^{-1} \in \phi(g)K\phi(g)^{-1} \subseteq K \end{aligned}$$

Since K is normal, so it $\phi(t) \in K$ so $t \in \phi^{-1}(K)$.

□

7.2 Kernels

Definition 7.2. If $\phi : G \rightarrow H$ is a group homomorphism, then the kernel of ϕ is

$$\text{Ker}(\phi) = \{g \in G \mid \phi(g) = e_H\}$$

Example 7.2.1. Consider

$$\phi : \mathbb{Z} \rightarrow \mathbb{Z}_n$$

given by $\phi(a) = a \pmod n$. Then,

$$\text{Ker}(\phi) = \{9, \pm n, \pm 2n, \dots\} = n\mathbb{Z}.$$

Theorem 7.2. Let $\phi : G \rightarrow H$ be any group homomorphism. Then, $\text{Ker}(\phi)$ is a normal subgroup of G .

Proof. $\{e_H\}$ is a normal subgroup in H . So $\phi^{-1}(\{e_H\})$ is normal subgroup in G . But

$$\begin{aligned} \phi^{-1}(\{e_H\}) &= \{g \in G \mid \phi(g) \in \{e_H\}\} \iff \phi(g) = e_H \\ &= \text{Ker}\phi \end{aligned}$$

□

Remark. Given $\phi : G \rightarrow H$ a group homomorphism, $\frac{G}{\text{Ker}(\phi)}$ is a group also.

Theorem 7.3 (First isomorphism theorem). Let $\phi : G \rightarrow H$ be any group homomorphism with $k = \text{Ker}(\phi)$ and $L = \text{Im}(\phi)$. Then, $G/K = \text{Im}(\phi)$

Example 7.2.2. $f : \mathbb{Z} \rightarrow \mathbb{Z}_4$ is a group homomorphism when $f(a) = a \pmod 4$. Then, $\text{Im}f = \mathbb{Z}_4$ because f is onto. Furthermore,

$$\text{Ker}f = \{0, \pm 4, \pm 8, \dots\} = \langle 4 \rangle.$$

By the first isomorphism theorem, $\mathbb{Z}/\langle 4 \rangle = \mathbb{Z}_4$.

Lemma 7.1. Let $\phi : G \rightarrow H$ be a group homomorphism with $K = \text{Ker}(\phi)$. Then, $\phi(a) = \phi(b)$ iff $aK = bK$.

Proof. (\Rightarrow) Suppose $\phi(a) = \phi(b)$. Then, $\phi(a)\phi(b)^{-1} = e_H$. So

$$e_H = \phi(a)\phi(b)^{-1} = \phi(a)\phi(b^{-1}) = \phi(ab^{-1}).$$

So $ab^{-1} \in K$ but this means $aK = bK$.

(\Leftarrow) If $aK = bK$, then $ab^{-1} \in K$. So

$$\phi(ab^{-1}) = \phi(a)\phi(b^{-1}) = \phi(a)\phi(b) = e_H.$$

But then $\phi(a) = \phi(b)$. □

Definition 7.3. Let G be a group with normal subgroup N . Then, the map $\pi : G \rightarrow G/N$ defined by $\pi(g) = gN$ is called the natural homomorphism.

Proof. We claim that $\varphi : G/K \rightarrow \text{Im}\phi$ given by $\varphi(gK) = \phi(g)$ is an isomorphism.

First, we want to show that φ is well-defined. Suppose $aK = bK$ with respect to $\varphi(aK) = \varphi(bK)$. Now, $\varphi(aK) = \phi(a)$ and $\varphi(bK) = \phi(b)$, and by the Lemma, $\phi(a) = \phi(b)$. So $\phi(aK) = \phi(bK)$.

Now, we want to show that φ is a group homomorphism. Let $aK, bK \in G/K$. Then,

$$\begin{aligned}\varphi(aKbK) &= \varphi(abK) \\ &= \phi(ab) \\ &= \phi(a)\phi(b) \\ &= \varphi(aK)\varphi(bK)\end{aligned}$$

Now, we want to show that φ is surjective. Let $t \in \text{Im}\phi$. So there is a $g \in G$ such that $\phi(g) = t$. But then $gK \in G/K$, and $\varphi(gK) = \phi(g) = t$.

Finally, we want to show that φ is injective. Let $aK, bK \in G/K$. Suppose

$$\varphi(aK) = \phi(a) = \pi(b) = \varphi(bK).$$

Since $\phi(a) = \phi(b)$, by the Lemma, $aK = bK$ so φ is injective.

So by these facts, $G/K \simeq \text{Im}\phi$. □

Example 7.2.3. For any $n \geq 1$, $\mathbb{Z}/\langle n \rangle \simeq \mathbb{Z}$.

Proof. We have a group homomorphism $f : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ given by $f(a) = a \pmod n$. Then,

$$\begin{aligned}\text{Im}f &= \mathbb{Z}_n \\ \text{Ker}f &= \{a_1 \pm n, \dots\} = \langle m \rangle\end{aligned}$$

So by the first isomorphism theorem, $\mathbb{Z}/\langle a \rangle = \mathbb{Z}_n$. □

Example 7.2.4. Let S_n be the symmetric group with $n \geq 3$. Define $f : S_n \rightarrow \mathbb{Z}_2$ by

$$f(\sigma) = \begin{cases} 0 & \text{if } \sigma \text{ is even} \\ 1 & \text{if } \sigma \text{ is odd} \end{cases}$$

This is a homomorphism.

Proof. Let $\sigma_1, \sigma_2 \in S_n$. Then,

1. σ_1, σ_2 both even:

$$f(\sigma_1\sigma_2) = 0 = 0 + 0 = f(\sigma_1) + f(\sigma_2)$$

2. σ_1 even, σ_2 odd:

$$f(\sigma_1\sigma_2) = 1 = 0 + 1 = f(\sigma_1) + f(\sigma_2)$$

3. σ_1 odd, σ_2 even: same as above

4. σ_1, σ_2 both odd:

$$f(\sigma_1\sigma_2) = 0 = 1 + 1 = f(\sigma_1) + f(\sigma_2)$$

□

Note that $\text{Ker}(f) = A_n$ and $\text{Im}(\phi) = \mathbb{Z}_2$. So $S_n/A_n \simeq \mathbb{Z}_2$.

Corollary 7.1. *For $n \geq 3$, A_n is normal in S_n .*

Proof. Since A_n is the kernel of some homomorphism, and because all kernels are normal, A_n is normal. \square

Example 7.2.5. Suppose $\phi : \mathbb{Z}_{30} \rightarrow \mathbb{Z}_{30}$ is a group homomorphism and $\text{Ker}(\phi) = \{0, 10, 20\}$. If $\phi(23) = 9$, find all elements of \mathbb{Z}_{30} that get mapped to 9, i.e. find all b such that $\phi(b) = 9$.

We want all b such that

$$\begin{aligned}\phi(23) = 9 &\iff 23 + K = b + K \\ &\iff (23 - b) \in K = \{0, 10, 20\} \\ &\iff b = 3, 13, 23\end{aligned}$$

Theorem 7.4. *Let $\phi : G \rightarrow H$ be a group homomorphism. Then, ϕ is one-to-one if and only if $\text{Ker}(\phi) = \{e_G\}$. (Kernel measures injectivity)*

Proof. (\Rightarrow) Assume ϕ is one to one with respect to $\text{Ker}(\phi) = \{e_G\}$. Take $g \in \text{Ker}(\phi)$. So

$$\phi(g) = e_H = \phi(e_G).$$

Since ϕ is one-to-one, $g = e_G$.

(\Leftarrow) Assume $\text{Ker}(\phi) = \{e_G\}$. Suppose $\phi(a) = \phi(b) \in H$. Hence,

$$\begin{aligned}e_H &= \phi(a)\phi(b)^{-1} \\ &= \phi(a)\phi(b^{-1}) \\ &= \phi(ab^{-1})\end{aligned}$$

So $ab^{-1} \in \text{Ker}(\phi)$. So $ab^{-1} = e_G \implies a = b$. \square

Theorem 7.5. *If $\phi : G \rightarrow H$ is injective, H has a subgroup isomorphic to G , namely $\text{Im}(\phi)$.*

Proof. ϕ injective implies that $\text{Ker}(\phi) = \{e_G\}$. By the first isomorphism theorem, we find that $G/\{e_G\} \simeq \text{Im}(\phi)$. But $G \simeq G/\{e_G\}$. So $G \simeq \text{Im}(\phi)$ \square

Theorem 7.6. *Suppose $\phi : G \rightarrow H$ is an onto homomorphism with $\text{Ker}(\phi) = K$. If $|G|, |H| < \infty$, then,*

$$|G| = |K||H|$$

Proof. Since ϕ is onto, by the first isomorphism theorem, we have $G/K = H$. So

$$|H| = |G/K| = |G|/|K| \implies |G| = |H||K|,$$

by the Lagrange's theorem. \square

Definition 7.4. *The trivial homomorphism $\phi : G \rightarrow H$ is the homomorphism $\phi(g) = e_H$ for all $g \in G$.*

Here are some useful facts:

1. $G/\{e_G\} \simeq G$.
2. $G/G \simeq \{e_G\}$.

Example 7.2.6. Suppose $\phi : \mathbb{Z}_p \rightarrow H$ with p a prime such that if ϕ is not the trivial homomorphism, then ϕ is one-to-one.

For this example, it is sufficient to show that $\text{Ker}(\phi) = \{0\}$. Note that \mathbb{Z}_p only has trivial subgroups. Since the Kernel of ϕ is a subgroup of \mathbb{Z}_p , we know that the kernel of ϕ is either equal to $\{0\}$ or \mathbb{Z}_p . If $\text{Ker}(\phi) = \mathbb{Z}_p$, then ϕ is the trivial homomorphism so it must be that $\text{Ker}(\phi) = \{0\}$.

Theorem 7.7 (Second isomorphism theorem). *Let H be a subgroup of G (not necessarily normal) and N a normal subgroup of G . Then,*

1. $HN = \{hn | h \in H, n \in N\}$ is a subgroup of G
2. N is normal in HN
3. $H \cap N$ is normal in N
4. $HN/N \simeq H/(H \cap N)$

Proof. Here's an idea of the proof:

1. Prove that HN is subgroup of G
2. Prove that N is normal in HN and $H \cap N$ is normal in H
3. Define a map $\phi : H \rightarrow HN/N$ by $\phi(h) = hN$
 - (a) Show ϕ is a homomorphism
 - (b) Show ϕ is onto
4. So 3 implies that $H/\text{Ker}(\phi) \simeq HN/N$ by the first isomorphism theorem
5. Show $\text{Ker}(\phi) = H \cap N$.

□

Theorem 7.8 (Correspondance theorem). *Let G be a group with a normal subgroup N .*

- *If K is a subgroup such that $N \subseteq K \subseteq G$, then K/N is a subgroup of G/N .*
- *If L is a subgroup of G/N , then there is a subgroup $N \subseteq K \subseteq G$ such that $L = K/N$.*

Example 7.2.7. Consider $G = \mathbb{Z}_8$ and $N = \{0, 4\}$.

- Subgroups of $N \subseteq K \subseteq G$:
 1. $N \subseteq N = \{0, 4\} \subseteq G$
 2. $N \subseteq \{0, 2, 4, 6\} \subseteq G$
 3. $N \subseteq G \subseteq G$
- Subgroups of $G/N \simeq \mathbb{Z}_8/N \simeq \mathbb{Z}_4$
 1. $N/N = \{0 + N\}$
 2. $\{0, 2, 4, 6\}/N = \{0 + N, 2 + N\}$
 3. $G/N = \{0 + N, 1 + N, 2 + N, 3 + N\}$

Theorem 7.9 (Third isomorphism theorem). *Let G be a group with H and N normal subgroups such that $N \subseteq H \subseteq G$. Then,*

$$G/H \simeq (G/N)/(H/N)$$

Proof. Here's a sketch of the proof:

1. Define a group homomorphism $\phi : G/N \rightarrow G/H$ by $\phi(gN) = gH$ (need to check that this is well-defined and a homomorphism)
2. Show that ϕ is onto
3. Show $\text{Ker}(\phi) = H/N \subseteq G/N$.

Combining (1), (2), (3), and the first isomorphism gives

$$(G/N)/(H/N) \simeq G/H,$$

as required. □

8 Rings

8.1 Introduction

Informally, a ring is an additive group with more structure.

Definition 8.1. A ring R is a set with two binary operations (usually addition and multiplication) such that for all $a, b \in R$,

1. $a + b = b + a$
2. $(a + b) + c = a + (b + c)$
3. There is an additive identity $0 \in R$ such that $a + 0 = 0 + a = a$
4. There is an additive inverse $(-a)$, i.e., for all $a \in R$, we have

$$a + (-a) = (-a) + a = 0$$

5. $(ab)c = a(bc)$
6. $a(b + c) = ab + ca$ and $(a + b)c = ac + bc$

Note that the first four conditions imply that a ring is an abelian group under addition

Definition 8.2. A ring R has an identity if there is $1 \in R$ such that $a1 = 1a = a$ for all a .

Definition 8.3. A ring R is commutative if $ab = ba$ for all $a, b \in R$.

Definition 8.4. A ring R is an (integral) domain if R has identity, is commutative, and $ab = 0$ implies $a = 0$ or $b = 0$


Definition 8.5. A ring R is a division ring if R has an identity and if every element $a \in R$ has a multiplicative inverse, i.e., there exists $a^{-1} \in R$ such that $aa^{-1} = 1$.

Definition 8.6. A field is a division ring that is commutative.

Remark. If $a \in R$ has a multiplicative inverse, we say a is a unit.

Example 8.1.1. Consider \mathbb{Z} , a set of all integers.

- binary operations are regular $+$ and \times .
- has an identity
- is commutative
- domain
- not a field since if $a \neq 1, -1, 0$, then a is not a unit



Example 8.1.2. $\mathbb{Q}, \mathbb{R}, \mathbb{C}$ are all rings. In fact, they are fields.

Example 8.1.3. \mathbb{Z}_n is a ring where operations are addition and multiplication modulo n .

- binary operations are addition and multiplication modulo n
- commutative with identity
- not a domain because $2 \times 3 = 0 \pmod{6}$ but $2 \neq 0$ and $3 \neq 0$. However, \mathbb{Z}_2 is a domain (in fact a field).

Example 8.1.4. $E = \{2n | n \in \mathbb{Z}\}$ is a ring that is commutative without identity.

Example 8.1.5. $M_2(\mathbb{R})$, a set of all two by two real matrices, is a ring that is not commutative but has an identity (the identity matrix).

Theorem 8.1 (Properties of a ring). *Let $a, b \in R$, then*

1. $a \cdot 0 = 0 \cdot a = 0$
2. $a(-b) = (-a)b = -(ab)$
3. $(-a)(-b) = ab$

If $1 \in R$, then

4. $(-1)a = -a$
5. $(-1)(-1) = 1$
6. *identity is unique*

Proof. (uniqueness of identity) Suppose 1 and $1'$ are identities of R . Then,

$$1 = 11' = 1'$$

So $1 = 1'$. □

Theorem 8.2 (uniqueness of inverses). *If $a \in R$ and a is a unit, then a^{-1} is unique.*

8.2 Subring

Definition 8.7. *A subset S of a ring R is a subring if S is also a ring under the operation of R .*

Theorem 8.3. *Let $S \subseteq R$ be a subset of a ring R . Then, S is a subring if*

1. $S \neq \emptyset$
2. $rs \in S$ for all $r, s \in S$ (closed under multiplication)
3. $r - s \in S$ for all $r, s \in S$ (closed under subtraction)

Example 8.2.1. $\{0\} \subseteq R$ and $R \subseteq R$ are the trivial subrings.

Example 8.2.2. $E = \{2n | n \in \mathbb{Z}\}$ is a subring of \mathbb{Z} . Note that \mathbb{Z} has an identity but E does not.