

MATH 2C03 - Differential Equations

Sang Woo Park

January 9, 2017

Course Outline

- Website: <https://ms.mcmaster.ca/lovric/2C3.html>
- Textbook: *Elementary Differential Equations with Boundary-Value Problems*
- Course pack must be bought.
- Assignments are online assignments.

Contents

1	Introduction	2
1.1	Differential equations	2
1.2	Ordinary differential equations	3
2	First-order ODEs	5
2.1	First-order ODEs	5

1 Introduction

1.1 Differential equations

Definition 1.1. *Differential equation is an equation where the unknown object is a function or a set of functions, and which involve their derivatives.*

Definition 1.2. An ordinary differential equation (ODE) is a differential equation involving one variable: $f'(x)$

Example 1.1.1 (ODE). The following differential equation represents exponential growth:

$$P'(t) = kP(t), k > 0$$

Definition 1.3. If a function, $f(x, y, z, \dots)$, involves more than one variable, we must use partial derivatives, e.g. $\partial f / \partial x, \partial^2 f / \partial x \partial y, \dots$. This is called a partial differential equation (PDE).

Example 1.1.2 (PDE). Let $c(x, t)$ represent the concentration at a location that is x units away from the source at time t . Then, we have

$$c_t(x, t) = Ac_{xx}(x, t)$$

Remark. Note that example 1.1.1 can be rearranged as follows:

$$\frac{P'(t)}{P(t)} = k$$

$P'(t)/P(t)$ represents the *relative rate of change*.

More often, k is not a constant. For example, we can incorporate seasonal variation to the exponential growth model.

Example 1.1.3.

$$\frac{P'(t)}{P(t)} = k \sin(at)$$

Definition 1.4. If a model does not involve any chance effects, it's a deterministic model. Otherwise, it's a stochastic model.

Example 1.1.4 (Deterministic model). Example 1.1.1 is a deterministic model.

Example 1.1.5 (Stochastic model). Going back to example 1.1.1, we may define k as follows:

$$k = \begin{cases} 0.6 & 35\% \text{chance} \\ 0.5 & 65\% \text{chance} \end{cases}$$

This is a stochastic model.

1.2 Ordinary differential equations

All ordinary differential equations (ODEs) contain the following:

- An independent variable
- Unknown function
- Derivative of the function

In other words, all ordinary equations have the following form:

$$F(x, y, y', y'', \dots, y^{(n)}) = 0.$$

Example 1.2.1. If $F(s, t) = s^2 + 2t - 4$, then

$$F(x, y) = x^2 + 2y - 4.$$

Note that this function is defined *implicitly*.

Example 1.2.2. If $F(s, t, u) = e^{su} - t^2$, then

$$F(x, y, y') = e^{xy'} = y^2.$$

This is a first order ODE.

Example 1.2.3. If $F(s, t, u, v, w) = w^2 - s + 4u$, then

$$F(x, y, y', y'', y''') = (y''')^2 - xy + 4y' = 0$$

This is a third order ODE.

Definition 1.5 (Order of an ODE). *Order of an ODE is determined by the highest non-zero derivative. If the highest non-zero derivative is the n -th derivative, the ODE also has an order of n .*

Remark. We categorize differential equations because each category requires a different methods to find its solution.

Example 1.2.4. Consider the following differential equation:

$$y'' - 3x^2y' + xy - 7e^x = 0$$

Since the highest non-zero derivative is the second derivative, this is a second order ODE. Note that this equation has the following form:

$$F(x, y, y', y'') = 0$$

We can rearrange this equation by solving for the highest derivative:

$$y'' = 3x^2y' - xy + 7e^x$$

We can also decide to put all y terms on LHS:

$$y'' - 3x^2y' + xy = 7e^x$$

Here, $7e^x$ is referred to as a *homogeneous term* since it does not contain y .

Definition 1.6 (Homogeneous ODE). *If the homogeneous term of an ODE is equal to 0, the ODE is called homogeneous.*

Definition 1.7 (Linear ODE). *Let $y^{(n)}(x)$ be the n -th derivative of $y(x)$. If an ODE can be written in the following form,*

$$a_n(x)y^{(n)}(x) + a_{n-1}(x)y^{(n-1)}(x) + \cdots + a_1(x)y'(x) + a_0(x)y(x) = p(x),$$

it is a linear.

Example 1.2.5. Consider the following equations:

1. $y''' - 3x^2y'' + 4y' = 76 \sin x$
2. $y'' - 3x^2\sqrt{y} = 76 \sin x$
3. $4y'' - 3y' + 4y = 6$
4. $y^{(4)} - 3y''y' - 4x^3y = 0$

Only equation 1 and 3 are linear.

Equation 1

$$\begin{cases} a_3(x) = 1 \\ a_2(x) = -3x^2 \\ a_1(x) = 4 \\ a_0(x) = 0 \end{cases}$$

Equation 2

\sqrt{y} is not linear.

Equation 43

$$\begin{cases} a_3(x) = 4 \\ a_2(x) = -3 \\ a_0(x) = 4 \\ p(x) = 6 \end{cases}$$

Equation 4

$y''y'$ is not linear.

2 First-order ODEs

2.1 First-order ODEs

Commonly, first order ordinary equations are written as

$$y' = G(x, y)$$

Example 2.1.1. $y' = 3x^3y^4 + 7 \ln x$.

Definition 2.1 (Pure time ODE). $y' = f(x)$

Definition 2.2 (Autonomous ODE). $y' = f(y)$

Example 2.1.2 (Pure time ODE). Consider the following ODE:

$$y' = 3x^2 - e^x$$

Since this is a pure time ODE, we can solve this ODE by integrating both sides with respect to the time variable x .

$$\begin{aligned} y &= \int (3x^2 - e^x) dx \\ &= x^3 - e^x + C \end{aligned}$$

Note that this solution gives a *family* of curves (we get a different curve for each value of C). Note that changing c shifts the curve vertically. To identify one solution from the family (i.e. to find the exact value of C), we need to know *initial condition*. If $y(0) = 4$, we have $C = 5$.

Definition 2.3 (General solution). *Integrating the ordinary differential equation gives us a solution with integration constant. This solution is called a general solution.*

Definition 2.4 (Particular solution). *Once we are given the initial condition, we can determine the exact value of the integration constant. Solution obtained by solving the Initial Value Problem is called a particular solution.*

Example 2.1.3. Consider the following ODE:

$$y'' = 4x^2,$$

where $y(0) = 3$ and $y'(0) = -7$. Since both values are given at $x = 0$, they are called *initial conditions*. We can also be asked to find the particular solution where $y(0) = 2$ and $y(4) = -3$. Since they occur at different x values, they are called *boundary conditions*.

When we're given the initial conditions, we can find the particular solution by integrating the ODE once:

$$y' = \frac{4}{3}x^3 + C.$$

However, when we're given the boundary conditions, we must integrate the ODE twice:

$$y = \frac{1}{3}x^4 + Cx + D$$

Definition 2.5 (General IVP). *Consider the following ODE:*

$$F(x, y, y', y'', \dots, y^{(n)}) = 0$$

IVP requires us to find the particular solution of the ODE given the initial conditions: $y(x_0) = y_0, y(x_1) = y_1, \dots, y^{(n-1)}(x_0) = y_{n-1}$. Note that the solution of an IVP, BVP is a function, and it can be algebraic, geometric, numeric, and/or qualitative.

Example 2.1.4. Show that

$$y = 1 - x + 4x \ln x$$

is a solution of the following IVP:

$$\begin{aligned} x^2 y'' - xy' + y &= 1 \\ y(1) &= 0 \\ y'(1) &= 3 \end{aligned}$$

Proof. First, we start by differentiating y :

$$\begin{aligned} y' &= -1 + 4 \ln x + 4x \frac{1}{x} = 4 \ln x + 3 \\ y'' &= \frac{4}{x} \end{aligned}$$

Then, by substitution, we have

$$\begin{aligned} &x^2 y'' - xy' + y \\ &= x^2 \left(\frac{4}{x} \right) - x(4 \ln x + 3) + 4 \ln x + 3 \\ &= 1 \end{aligned}$$

It is also easy to verify that $y(1) = 0$ and $y'(1) = 3$. □