
Contents

6	Many-Body Perturbation Theory	3
6.1	Rayleigh-Schrödinger (RS) Perturbation Theory	3
*6.2	Diagrammatic Representation of RS Perturbation Theory	3
6.2.1	Diagrammatic Perturbation Theory for Two States	3
	Exercise 6.1	3
6.2.2	Diagrammatic Perturbation Theory for N States	4
	Exercise 6.2	4
6.2.3	Summation of Diagrams	6
6.3	Orbital Perturbation Theory: One-Particle Perturbations	6

CHAPTER 6

Many-Body Perturbation Theory

6.1 Rayleigh-Schrödinger (RS) Perturbation Theory

*6.2 Diagrammatic Representation of RS Perturbation Theory

6.2.1 Diagrammatic Perturbation Theory for Two States

Exercise 6.1

Write down and evaluate all fifth-order diagrams that have the property that an imaginary horizontal line crosses only one hole and one particle line. Show that the sum of such diagrams is

$$\frac{V_{12}V_{21}(V_{22} - V_{11})^3}{(E_1^{(0)} - E_2^{(0)})^4}$$

Hint: There are eight such diagrams, and they can be generated by adding three dots to the second-order diagram in all positive ways.

Solution 6.1

The final results are listed below firstly.




Figure 6.1: All fifth-order diagrams, which have the property that an imaginary horizontal line crosses only one hole and one particle line, and their mathematical expressions.

Note that the diagrams, which have the property that an imaginary horizontal line crosses only one hole

and one particle line, have no pair of hole/particle lines whose overlap is nonempty. For any pair of hole/particle lines which connect the dots m_1 and n_1 , m_2 and n_2 , where $n_1 > m_1$ and $n_2 > m_2$, their overlap will be

- $[\max\{m_1, m_2\}, \min\{n_1, n_2\}]$, if $\min\{n_1, n_2\} > \max\{m_1, m_2\}$;
- empty, otherwise.

For example, if there is a pair of hole lines, one connecting the dot 1 and 3, and the other connecting 2 and 5, their overlap will be $[2, 3]$. Take an another example, if there is a pair of particle lines, one connecting the dot 5 and 4, and the other connecting 2 and 1, their overlap will be empty.

Thus, there are four cases.

- Four hole lines and one particle line. There is only one method, hole lines are $(1, 2)$, $(2, 3)$, $(3, 4)$, $(4, 5)$ and the only particle line is $(5, 1)$, as the first subdiagram in Fig 6.1.
- Three hole lines and two particle lines. There are three methods as follows.
 - Hole lines are $(1, 2)$, $(2, 3)$, and $(3, 5)$ while particle lines are $(5, 4)$, $(4, 1)$.
 - Hole lines are $(1, 2)$, $(2, 4)$, and $(4, 5)$ while particle lines are $(5, 3)$, $(3, 1)$.
 - Hole lines are $(1, 3)$, $(3, 4)$, and $(4, 5)$ while particle lines are $(5, 2)$, $(2, 1)$.

They correspond to the second, third, fourth subdiagram in Fig 6.1.

- Two hole lines and three particle lines. There are three methods as follows.
 - Hole lines are $(1, 3)$, $(3, 5)$ while particle lines are $(5, 4)$, $(4, 2)$, and $(2, 1)$.
 - Hole lines are $(1, 4)$, $(4, 5)$ while particle lines are $(5, 3)$, $(3, 2)$, and $(2, 1)$.
 - Hole lines are $(1, 2)$, $(2, 5)$ while particle lines are $(5, 4)$, $(4, 3)$, and $(3, 1)$.

They correspond to the fifth, sixth, seventh subdiagram in Fig 6.1.

- One hole line and four particle lines. There is only one method, the only hole line is $(1, 5)$ and particle lines are $(5, 4)$, $(4, 3)$, $(3, 2)$, and $(2, 1)$, as the eighth subdiagram in Fig 6.1.

Thus, the sum of such diagrams is


$$-\frac{V_{12}V_{21}(V_{11}^3 - 3V_{11}^2V_{22} + 3V_{11}V_{22}^2 - V_{22}^3)}{(E_1^{(0)} - E_2^{(0)})^4} = -\frac{V_{12}V_{21}(V_{11} - V_{22})^3}{(E_1^{(0)} - E_2^{(0)})^4} = \frac{V_{12}V_{21}(V_{22} - V_{11})^3}{(E_1^{(0)} - E_2^{(0)})^4}.$$

In fact, as the textbook says, these eight diagrams can be generated by adding three dots to the second-order diagram in all positive ways. In fact, any pair of hole/particle lines in them has also empty overlap. I think the calculation of the overlap is much direct than inspecting the property of lines.

6.2.2 Diagrammatic Perturbation Theory for N States

Exercise 6.2

Use diagrammatic techniques to obtain the fourth-order perturbation energy of a particular state (say, i) of an N -state system. That is, evaluate the diagrams



where the indices m, n, k, \dots exclude i . Using the approach of Section 6.1, obtain an algebraic expression for the fourth-order energy and compare it to the diagrammatic result.

Solution 6.2




Figure 6.2: All fourth-order diagrams and their mathematical expressions.

htrETWREARHSTJ

$$\begin{aligned}
E_i^{(4)} &= \langle i | \mathcal{V} | \Psi_i^{(3)} \rangle = \sum'_n \langle i | \mathcal{V} | n \rangle \langle n | \Psi_i^{(3)} \rangle \\
&= \sum'_n \frac{V_{in}}{E_i^{(0)} - E_n^{(0)}} \left[\langle n | \mathcal{V} | \Psi_i^{(2)} \rangle - E_i^{(1)} \langle n | \Psi_i^{(2)} \rangle - E_i^{(2)} \langle n | \Psi_i^{(1)} \rangle \right] \\
&= \sum'_n \frac{V_{in} \langle n | \mathcal{V} | \Psi_i^{(2)} \rangle}{E_i^{(0)} - E_n^{(0)}} - E_i^{(1)} \sum'_n \frac{V_{in} \langle n | \Psi_i^{(2)} \rangle}{E_i^{(0)} - E_n^{(0)}} - E_i^{(2)} \sum'_n \frac{V_{in} \langle n | \mathcal{V} | \Psi_i^{(1)} \rangle}{E_i^{(0)} - E_n^{(0)}} \\
&= \sum'_{mn} \frac{V_{in} \langle n | \mathcal{V} | m \rangle \langle m | \Psi_i^{(2)} \rangle}{E_i^{(0)} - E_n^{(0)}} - E_i^{(1)} \sum'_n \frac{V_{in}}{E_i^{(0)} - E_n^{(0)}} \langle n | \Psi_i^{(2)} \rangle - E_i^{(2)} \sum'_{mn} \frac{V_{in} \langle n | \mathcal{V} | m \rangle \langle m | \Psi_i^{(1)} \rangle}{E_i^{(0)} - E_n^{(0)}} \\
&= \sum'_{mn} \frac{V_{in} V_{nm}}{E_i^{(0)} - E_n^{(0)}} \langle m | \Psi_i^{(2)} \rangle - \sum'_n \frac{V_{ii} V_{in}}{E_i^{(0)} - E_n^{(0)}} \langle n | \Psi_i^{(2)} \rangle - \left(\sum'_k \frac{V_{ik} V_{ki}}{E_i^{(0)} - E_k^{(0)}} \right) \sum'_{mn} \frac{V_{in} V_{nm} \langle m | \Psi_i^{(1)} \rangle}{E_i^{(0)} - E_n^{(0)}} \\
&= \sum'_{mn} \frac{V_{in} V_{nm}}{E_i^{(0)} - E_n^{(0)}} \frac{\langle m | \mathcal{V} | \Psi_i^{(1)} \rangle - E_i^{(1)} \langle m | \Psi_i^{(1)} \rangle}{E_i^{(0)} - E_m^{(0)}} - \sum'_n \frac{V_{ii} V_{in}}{E_i^{(0)} - E_n^{(0)}} \frac{\langle n | \mathcal{V} | \Psi_i^{(1)} \rangle - E_i^{(1)} \langle n | \Psi_i^{(1)} \rangle}{E_i^{(0)} - E_n^{(0)}} \\
&\quad - \left(\sum'_k \frac{V_{ik} V_{ki}}{E_i^{(0)} - E_k^{(0)}} \right) \sum'_{mn} \frac{V_{in} V_{nm}}{E_i^{(0)} - E_n^{(0)}} \langle m | \Psi_i^{(1)} \rangle \\
&= \sum'_{mn} \frac{V_{in} V_{nm}}{(E_i^{(0)} - E_n^{(0)})(E_i^{(0)} - E_m^{(0)})} \left[\sum'_k \langle m | \mathcal{V} | k \rangle \langle k | \Psi_i^{(1)} \rangle - E_i^{(1)} \langle m | \Psi_i^{(1)} \rangle \right] \\
&\quad - \sum'_n \frac{V_{ii} V_{in}}{(E_i^{(0)} - E_n^{(0)})^2} \left[\sum'_m \langle n | \mathcal{V} | m \rangle \langle m | \Psi_i^{(1)} \rangle - E_i^{(1)} \langle n | \Psi_i^{(1)} \rangle \right] - \sum'_{mnk} \frac{V_{ik} V_{ki} V_{in} V_{nm} \langle m | \Psi_i^{(1)} \rangle}{(E_i^{(0)} - E_n^{(0)})(E_i^{(0)} - E_k^{(0)})}
\end{aligned}$$

$$\begin{aligned}
&= \sum'_{mnk} \frac{V_{in} V_{nm} V_{mk}}{(E_i^{(0)} - E_n^{(0)})(E_i^{(0)} - E_m^{(0)})} \langle k | \Psi_i^{(1)} \rangle - \sum'_{mn} \frac{V_{ii} V_{in} V_{nm}}{(E_i^{(0)} - E_n^{(0)})(E_i^{(0)} - E_m^{(0)})} \langle m | \Psi_i^{(1)} \rangle \\
&\quad - \sum'_{mn} \frac{V_{ii} V_{in} V_{nm}}{(E_i^{(0)} - E_n^{(0)})^2} \langle m | \Psi_i^{(1)} \rangle + \sum'_n \frac{V_{ii}^2 V_{in}}{(E_i^{(0)} - E_n^{(0)})^2} \langle n | \Psi_i^{(1)} \rangle - \sum'_{mnk} \frac{V_{ik} V_{ki} V_{in} V_{nm}}{(E_i^{(0)} - E_n^{(0)})(E_i^{(0)} - E_k^{(0)})} \langle m | \Psi_i^{(1)} \rangle
\end{aligned}$$

6.2.3 Summation of Diagrams

6.3 Orbital Perturbation Theory: One-Particle Perturbations