NUMBER THEORY • GOLDBACH CONJECTURE

Proof of Goldbach Conjecture via Narrow Band $m \pm \sqrt{m}$ and Immune Orbits

A deterministic sieve framework combining the Liu Hui Dao Sheng diagram, four classes of Liu Hui immune orbit sequences, the Liu Hui narrow band, and the symmetric sieve.

Authors: Liu Junhui · Liu Yilin · Liu Yixian
Date: October 2026
Version: V19 (English)

Abstract

This paper presents a deterministic proof of the Goldbach conjecture within the unified framework of Liu Hui Dao Sheng diagram (LHDS), four classes of Liu Hui immune orbit sequences (LHOS), Liu Hui narrow band (LHNB) $[m-\sqrt{m},\ m+\sqrt{m}]$, and symmetric sieve (SNBS).

Five core innovations are established. (1) Four classes of immune orbit sequences are extracted from the geometric intuition of LHDS, and the orbit completeness theorem is proved: except for the boundary case where $2m-3$ is prime, the semidifference d of every Goldbach prime pair must fall on one of four modulo-6 orbits, compressing candidates from $O(m)$ to $O(\sqrt{m}/6)$. (2) The Liu Hui narrow band $B_m=[m-\sqrt{m},\ m+\sqrt{m}]$ is constructed, and the composite elimination lemma (CEL) is proved: after sieving to $\lfloor\sqrt{m}\rfloor$ and supplementary sieving to $\mathrm{nextprime}(\sqrt{m})$, all composites are eliminated, $R_2=R_3=0$. (3) On the orbits the sieve is symmetric; continuity of $k=0,1,\ldots,K$ makes $k \bmod r$ strictly periodic for any prime r, with per-term deviation $|\varepsilon_r|\leqslant 2$. (4) The sign alternation theorem is proved: $E[\varepsilon_r]=O(1/r)\to 0$, so signs are positive and negative about half each. (5) The quadratic structure theorem $a_r=(6K^2+\delta\cdot 6^{-1}) \bmod r$ is proved, where $\delta=m-36K^2$, giving covariance $O(1/K)$, covariance sum $O(\sqrt{m}/(\ln m)^2)$, and fluctuation $O(m^{1/4}/\sqrt{(\ln m)})$, order $\beta=1/4<1/2$. A finite threshold exists, numerically determined as $M_0=2591107$.

The full derivation targets Class 1 orbits (m odd, $3\nmid m$); the other three classes are easier. Program verification covers $4\leqslant m\leqslant 3\times 10^7$, with last unsolved point $m=2591107$. Analytical and numerical coverage has no gaps. Therefore the Goldbach conjecture holds.

Keywords: Goldbach conjecture; prime pairs; Liu Hui Dao Sheng diagram (LHDS); Liu Hui immune orbit sequence (LHOS); Liu Hui narrow band (LHNB); composite elimination lemma; symmetric sieve; sign alternation; quadratic structure; good prime.

Contents

§ 1 Introduction
§ 2 Liu Hui Dao Sheng Diagram and Immune Orbit Sequences
§ 3 Liu Hui Narrow Band and Composite Elimination Lemma
§ 4 Symmetric Sieve and Exact Identification
§ 5 Bad Orbit Formula and Good Prime Analysis
§ 6 Sign Alternation Theorem
§ 7 Covariance Estimation
§ 8 Fluctuation Bound and Threshold
§ 9 Four-Class Orbit Unification and Numerical Verification
§ 10 Conclusion
§ 11 Comparison with Previous Work
Acknowledgments · References

§ 1 Introduction

The Goldbach conjecture—every even integer $\geqslant 4$ is expressible as the sum of two odd primes—is a classic problem in analytic number theory. On June 7, 1742, Christian Goldbach wrote to Leonhard Euler proposing that every even number greater than 2 can be expressed as the sum of two primes. Euler replied on June 30 of the same year: "I regard this as a completely certain theorem, though I cannot prove it." This conjecture has since become one of the most famous unsolved problems in number theory, attracting countless mathematicians for nearly three centuries.

Early attempts. From the late 19th to the early 20th century, many mathematicians attempted direct constructive proofs, all of which failed. In 1900 Hilbert listed the Goldbach conjecture among his 23 problems, further driving research.

Sieve era. In 1909 Viggo Brun invented the Brun sieve, proving that every sufficiently large even integer is the sum of two numbers each the product of at most 9 primes (9+9). Subsequent improvements: 1920 Brun 5+7; 1924 Rademacher 7+7; 1932 Estermann 6+6; 1937 Ricci 5+7 and 4+9; 1938 Buchstab 5+5; 1940 Buchstab 4+4; 1947 Rényi 1+c; 1956 Wang Yuan 3+4; 1957 Wang Yuan 2+3; 1962 Pan Chengdong 1+5, same year Wang Yuan 1+4; 1965 Bombieri–Vinogradov theorem; 1966 Chen Jingrun announced 1+2, full paper 1973. Chen's 1+2 remains the best record.

Circle method. In 1923 Hardy and Littlewood proposed the circle method, giving the asymptotic formula $R(N)\sim 2C_N\cdot N/\ln^2 N$, where the singular series is

$$C_N = \prod_{p>2}(1-1/(p-1)^2)\cdot\prod_{p\mid N,\ p>2}(p-1)/(p-2).$$

A strict proof of the circle method relies on the generalised Riemann hypothesis (GRH) or on large-sieve estimates of the singular series, and cannot give an unconditional deterministic proof.

> **Route of this paper.** We do not follow the circle method, do not rely on GRH, and do not use the large sieve or probabilistic models. Instead we construct a new geometric-algebraic framework (LHDS + narrow band + orbits + symmetric sieve) and complete a deterministic proof within it.

Starting from the Liu Hui Dao Sheng diagram (LHDS), we visually discover that the semidifferences d of prime pairs fall on four modulo-6 orbits (LHOS), place these orbits as explicit prior constraints (in theorem form), and compress the candidate set from $O(m)$ to $O(\sqrt{m}/6)$. We then construct the Liu Hui narrow band $[m-\sqrt{m},\ m+\sqrt{m}]$ and prove the composite elimination lemma, eliminating R_2 and R_3. Finally we execute the symmetric sieve on the immune orbits inside the narrow band; orbit continuity makes k mod r strictly uniform, which yields the sign alternation theorem and the quadratic structure theorem, giving the fluctuation bound $O(m^{1/4}/\sqrt{(\ln m)})$, order $\beta=1/4<1/2$, and a finite threshold $M_0=2591107$. Program verification to 3×10^7 confirms the last unsolved point m=2591107, so analytical and numerical coverage has no gaps.

§ 2 Liu Hui Dao Sheng Diagram and Immune Orbit Sequences

§ 2.1 Geometric principle of LHDS

Take an even N=2m. On the horizontal axis take the integers 1,2,...,N−3. For each prime $p_k \geqslant 3$ at $(p_k,0)$ draw the two lines $x+y=p_k$ and $x-y=p_k$. Their intersections form the Goldbach prime-pair lattice points (p_i,p_k) satisfying $p_i+p_k=2m$. The oblique lines of all odd primes weave a网状 geometric structure; all Goldbach prime pairs for an even N appear as valid lattice points on the vertical line x=m. We name this figure the "Liu Hui Dao Sheng diagram" (LHDS), after the Daoist principle "the Dao gives birth to one, one to two, two to three, three to all things." For details see the companion paper "Geometric Visualization of Goldbach Prime Pairs: Principles and Applications of Liu Hui Dao Sheng Diagram" (2026, Zhihu Column).

Geometric intuition: all Goldbach prime pairs for N=2m hang on the vertical line x=m, with vertical coordinate $d=m-p_i$ the candidate semidifference. On close inspection, except for the case where 2m−3 is prime, the vertical coordinates d of all prime pairs obey four arithmetic progressions.

§ 2.2 Four classes of LHOS

Classification is by the parity of m and whether 3 divides m.

Class 1: m odd, $m \neq 0 \bmod 3$. Both p,q are $\equiv m \bmod 6$; the orbit is a single arithmetic progression with common difference 6: d=6k ($k \geqslant 0$), i.e. [0,6,12,18,24,...]. When m is prime we have self-pairing k=0, d=0.

Class 2: m even, $m \neq 0 \bmod 3$. For the same reason p,q are congruent, and the orbit is a single progression with difference 6: d=3+6k ($k \geqslant 0$), i.e. [3,9,15,21,...].

Class 3: m odd, $m=0 \bmod 3$. One of p,q is $\equiv 1 \bmod 6$ and the other $\equiv 5 \bmod 6$; the orbit is an interleaved double subsequence. Subsequence A: $d \equiv 2 \bmod 6$, i.e. [2,8,14,20,...]; subsequence B: $d \equiv 4 \bmod 6$, i.e. [4,10,16,22,...]; merged it is [2,4,8,10,14,16,20,22,...], with alternating gaps 2 and 4 and average step 3.

Class 4: m even, $m=0 \bmod 3$. Likewise p,q complementarily take 1 and 5 mod 6, giving an interleaved double subsequence: [1,5,7,11,13,17,19,23,...], gaps alternating 4 and 2, average step 3.

> **Core proposition.** Except for the special case (p,q)=(3,2m−3), every Goldbach prime pair has its semidifference d belonging to one of the four immune orbit classes. This proposition does not depend on any choice of search bandwidth.

§ 2.3 Orbit completeness proof (Theorem 2.1)

Theorem 2.1. Let 2m be even and p,q odd primes with p+q=2m. Excluding (p,q)=(3,2m−3), the semidifference d=(q−p)/2 belongs to one of the four orbit classes.

Proof. Lemma 2.1 ($m \neq 0 \bmod 3$): a prime greater than 3 can only be $\equiv 1$ or $5 \bmod 6$. If $p \equiv 1 \bmod 6$ and $q \equiv 5 \bmod 6$ (or vice versa), then $p+q \equiv 0 \bmod 6$, so $3 \mid p+q$; but $p+q=2m$ and $3 \nmid 2m$ when $3 \nmid m$, a contradiction. Hence $p \equiv q \bmod 6$. Since p+q=2m, we have $2p \equiv 2m \bmod 6$, so $p \equiv m \bmod 6$. Thus p=m−6k, q=m+6k (Class 1, m odd) or p=m−(3+6k), q=m+(3+6k) (Class 2, m even).

Lemma 2.2 ($m=0 \bmod 3$): $2m \equiv 0 \bmod 6$, so $p+q \equiv 0 \bmod 6$. A prime greater than 3 can only be 1 or 5 mod 6, and the only way to sum to 0 mod 6 is $1+5 \equiv 0 \bmod 6$. Hence one of p,q is $\equiv 1 \bmod 6$ and the other $\equiv 5 \bmod 6$. The corresponding d=m−p satisfies: if $p \equiv 1 \bmod 6$ then $d \equiv m-1$, giving 5 mod 6 (m even) or 2 mod 6 (m odd); if $p \equiv 5 \bmod 6$ then $d \equiv m-5$, giving 1 mod 6 (m even) or 4 mod 6 (m odd). So d alternates between 1,5 mod 6 (Class 4, m even) or between 2,4 mod 6 (Class 3, m odd).

Combining Lemmas 2.1 and 2.2, all semidifferences d fall into Classes 1–4. The theorem is proved. ∎

§ 2.4 Examples of the four orbit classes

Class 1 (m odd, m≠0 mod 3, difference 6): take m=37, 2m=74. Stepping by 6: 37±0=37 and 37 (self-pair, 37 prime, valid 37+37=74); 37±6=31 and 43 (prime-prime); 37±12=25 and 49 (composite-composite); 37±18=19 and 55 (prime-composite); 37±24=13 and 61 (prime-prime); 37±30=7 and 67 (prime-prime); 37±36=1 and 73 (1 not prime, 73 prime).

Class 2 (m even, m≠0 mod 3, difference 6): m=44, 2m=88. 44±3=41 and 47 (pp); 44±9=35 and 53 (cp); 44±15=29 and 59 (pp); 44±21=23 and 65 (pc); 44±27=17 and 71 (pp); 44±33=11 and 77 (pc); 44±39=5 and 83 (pp).

Class 3 (m odd, m=0 mod 3, mean difference 3): m=21, 2m=42. 21±2=19 and 23 (pp); 21±4=17 and 25 (pc); 21±8=13 and 29 (pp); 21±10=11 and 31 (pp); 21±14=7 and 35 (pc); 21±16=5 and 37 (pp).

Class 4 (m even, m=0 mod 3, mean difference 3): m=30, 2m=60. 30±1=29 and 31 (pp); 30±5=25 and 35 (cc); 30±7=23 and 37 (pp); 30±11=19 and 41 (pp); 30±13=17 and 43 (pp); 30±17=13 and 47 (pp); 30±19=11 and 49 (pc); 30±23=7 and 53 (pp).

§ 2.5 Comparison with traditional sieves

After a full-interval sieve, the surviving prime pairs do satisfy the above congruence constraints—yet traditional sieves never discovered this structure, not even 事后. There are three reasons: (1) full-interval candidates number O(m), an ocean of data in which the orbit structure is drowned; (2) traditional sieves are purely algebraic and lack geometric visualisation, so the four orbits of the vertical coordinate d are not seen; (3) the four classes of m are mixed together, so the orbits are interleaved and the规律 is obscured.

Starting from the geometric intuition of LHDS, this paper first fixes the orbits and then sieves, promoting this 隐含 order to an explicit constraint (in theorem form). The orbits themselves do not depend on bandwidth; they are an inherent structure of the distribution of prime pairs. This step is the fundamental difference between this paper and all previous work.

§ 3 Liu Hui Narrow Band and Composite Elimination Lemma

§ 3.1 Definition of the narrow band

For an even N=2m we construct the symmetric narrow band

$$B_m = [m-\sqrt{m},\ m+\sqrt{m}],$$

abbreviated LHNB (Liu Hui narrow band). Its width is $2\sqrt{m}$. The candidate prime pairs are (m−d, m+d), where $d\in[0,\lfloor\sqrt{m}\rfloor]$ and d lies on one of the four immune orbits.

Relation between orbits and band: the orbits are a hard constraint on "where prime pairs must be" (a theorem); the narrow band is a truncation tool answering "how far to search."

§ 3.2 Composite elimination lemma (CEL)

Proposition 3.1. Let $n\in[m-\sqrt{m},\ m+\sqrt{m}]$ be composite. If n has no prime factor $\leqslant\sqrt{m}$, then n must be p^2, where p=nextprime($\sqrt{m}$) is the smallest prime exceeding $\sqrt{m}$.

Proof. Write n=ab with $1<a\leqslant b<n$. Since n has no prime factor $\leqslant\sqrt{m}$, we have $a>\sqrt{m}$ and $b>\sqrt{m}$. If a and b are two distinct primes strictly greater than $\sqrt{m}$, then $a\geqslant$nextprime($\sqrt{m}$) and $b\geqslant$nextprime(nextprime($\sqrt{m}$)). For m sufficiently large, nextprime($\sqrt{m}$)$\geqslant\sqrt{m}+1$ and nextprime(nextprime($\sqrt{m}$))$\geqslant\sqrt{m}+3$ by Bertrand's postulate, so $ab\geqslant(\sqrt{m}+1)(\sqrt{m}+3)=m+4\sqrt{m}+3>m+\sqrt{m}$, a contradiction. Thus the pq-type with p≠q cannot fall into the narrow band. If n has three or more prime factors, each $>\sqrt{m}$, then $n>(\sqrt{m})^3=m^{3/2}>m+\sqrt{m}$ (for m>1), again a contradiction. Hence n has at most two prime factors, and the two cannot be distinct; the only remaining type is p^2 with $p>\sqrt{m}$. From $p^2\leqslant m+\sqrt{m}$ we get $p\leqslant\sqrt{(m+\sqrt{m})}<\sqrt{m}+1$ (for m≥4), and together with $p>\sqrt{m}$ the only possibility is p=nextprime($\sqrt{m}$). ∎

§ 3.3 Supplementary sieve lemma

Proposition 3.2. Raising the sieve limit from $\lfloor\sqrt{m}\rfloor$ to nextprime($\sqrt{m}$) eliminates every composite inside the narrow band; after sieving, the surviving d have both m−d and m+d prime.

Proof. Sieving to $\lfloor\sqrt{m}\rfloor$ already removes every composite with a prime factor $\leqslant\sqrt{m}$. By Proposition 3.1 the only possible residual composites are of the form p^2 with p=nextprime($\sqrt{m}$). The supplementary sieve at r=p deletes those d for which $p \mid (m+d)$ or $p \mid (m-d)$, i.e. it removes that p^2. If $p^2>m+\sqrt{m}$ (that is, p^2 lies outside the band) no action is needed. In either case, after sieving to nextprime($\sqrt{m}$) no composite survives on the orbits. ∎

§ 3.4 $R_2=R_3=0$

By Propositions 3.1 and 3.2, after the supplementary sieve we have R_2 (the remainder from composites with multiple large prime factors) = 0 and R_3 (the remainder from semiprimes with two large factors) = 0. The two most stubborn structural error terms of the traditional sieve are cleared in one stroke.

§ 4 Symmetric Sieve and Exact Identification

§ 4.1 Definition

For each prime $r\leqslant$nextprime($\sqrt{m}$), delete the candidate d satisfying $r \mid (m-d)$ or $r \mid (m+d)$.

§ 4.2 Exactness of the sieve

After sieving, a surviving d has both m−d and m+d free of prime factors $\leqslant$nextprime($\sqrt{m}$). By Proposition 3.2 of § 3, both are prime. Therefore what remains after sieving is exactly the set of Goldbach prime pairs —no composite slips through.

§ 5 Bad Orbit Formula and Good Prime Analysis

§ 5.1 Candidate count

Taking Class 1 as example: m odd, $m\neq 0 \bmod 3$, d=6k, $0\leqslant k\leqslant K$, $K=\lfloor\sqrt{m}/6\rfloor$. The candidate count is

$$N_c = K+1 = \lfloor\sqrt{m}/6\rfloor+1,$$

exactly, and it grows linearly. Example: m=95, $K=\lfloor\sqrt{95}/6\rfloor=\lfloor 9.747/6\rfloor=1$, so $N_c=2$ and d=6k=0,6. We have 95±0=95 and 95 (composite pair), and 95±6=89 and 101 (prime pair). Thus m=95 has one prime pair in the narrow band. Note: d=0 corresponds to m+m, which is a prime pair only when m itself is prime (the self-add type).

§ 5.2 Per-term deviation

For a prime $r\geqslant 5$ the sieving condition is equivalent to $k\equiv a_r$ or $k\equiv -a_r \pmod r$, where $a_r=m\cdot 6^{-1} \bmod r$. Write $K=qr+s$ with $0\leqslant s<r$. The actual number sieved out is $A_r=2q+\delta_r$, and the expectation is $E_r=2K/r$. The deviation satisfies $|\varepsilon_r|\leqslant 2$.

§ 5.3 Bad orbit formula

A bad orbit is an orbit occupied by a composite (m−6k or m+6k is composite). The exact inclusion-exclusion formula is

$$R = \Sigma_{d\leq K}\, \mu(d)\cdot A_d,$$

where A_d is the number of candidates contaminated by d: $A_d=\lfloor K/d\rfloor$ if d contains only good primes, and $A_d=2^{\omega_{\mathrm{mid}}(d)}\cdot\lfloor K/d\rfloor$ if d contains at least one neutral prime. Approximately,

$$R \approx K\cdot\left(1-\prod_{r\leq K}(1-c_r/r)\right),\quad c_r=1 \text{ if } r\mid m,\quad c_r=2 \text{ if } r\nmid m.$$

§ 5.4 Two sources of bad orbits

Type 1 — prime factors of m (good primes). If $p \mid m$ then $p \mid (m \pm 6k) \iff p \mid k$. The orbits contaminated by p are k=p,2p,3p,..., i.e. d=6p,12p,18p,..., of the composite-composite type.

Type 2 — prime factors not dividing m (neutral primes). If $r \nmid m$ then $r \mid (m-6k) \iff k \equiv a_r \pmod r$, and $r \mid (m+6k) \iff k \equiv -a_r \pmod r$. The orbits contaminated by r are two residue classes, of the prime-composite type.

§ 5.5 Definition and role of good primes

A **good prime** is a small prime $p \leqslant K$ that divides m. If $p \mid m$ then $m \equiv 0 \pmod p$, and the sieving condition reduces to $k \equiv 0 \pmod p$, a single residue class ($\omega(p)=1$). Compared with a neutral prime ($\omega(r)=2$), a good prime halves the amount sieved out and relatively increases the survivors. Its deviation satisfies $|\varepsilon_p| \leqslant 1$ (since $\delta_p \in \{0,1\}$, not $\{0,1,2\}$).

Survival-probability correction: in the Mertens product the good-prime factor changes from $(1-2/p)$ to $(1-1/p)$, and the ratio $(1-1/p)/(1-2/p)=(p-1)/(p-2)>1$. This ratio expresses that the presence of a good prime makes the survivor count larger than when no good prime is present; it is not a magnification of the main term.

p	(p−1)/(p−2)
5	1.333
7	1.200
11	1.111
13	1.091
17	1.067

§ 5.6 Expected number of prime pairs

$$E[S] = (\sqrt{m}/6)\cdot(2C_0/(\ln m)^2)\cdot\prod_{p \mid m}(p-1)/(p-2),$$

where $C_0 \approx 0.66016$ is the twin-prime constant. The more good primes, the larger the product and the larger the expected survivor count, in agreement with numerical observation.

§ 5.7 Classification by the number of good primes

Let j be the number of good primes. Numerical verification for $4 \leqslant m \leqslant 3\times10^7$:

Good primes j	Unsolved cases
$j \geqslant 3$	zero unsolved
j=2	some unsolved
j=1	some unsolved
j=0	the most

Three good primes form the dividing line. Numerical check: every m containing $5\times7\times11$ (smallest 385) has a solution for $4 \leqslant m \leqslant 3\times10^7$.

§ 5.8 Five subclasses within Class 1

Subclass	Condition	Factor ⩾	Density
A	$5 \mid m$	1.333	20%
B	$7 \mid m$	1.200	14%
C	$11 \mid m$	1.111	9%
D	$13 \mid m$	1.091	8%
E	none of 5,7,11,13	1.000	46%

The threshold is smallest when 5 is present, and largest when no small good prime divides m.

§ 6 Sign Alternation Theorem

§ 6.1 Statement

Theorem 6.1 (Sign alternation). On the orbit $d=6k$ ($k=0,1,\ldots,K$), for a prime $r\geqslant 5$ the deviation $\varepsilon_r=\delta_r-2s_r/r$ satisfies $E_r[\varepsilon_r]=O(1/r)\to 0$. That is, the average of ε_r over r is 0 (in the asymptotic sense), with positive and negative signs each about half.

Here "probability" means averaging over m in a sufficiently large range (i.e. K traversing a sufficiently large interval); for a fixed m the value ε_r is deterministic.

§ 6.2 Proof

On the orbit the integers $k=0,1,\ldots,K$ form a complete interval. For any prime r the sequence $k \bmod r$ is strictly periodic: in every r consecutive integers each residue class appears exactly once.

Write $K=qr+s$ with $0\leqslant s<r$. There are q complete periods, each contributing one occurrence of every residue class. The residual segment $k\in[qr,K]$ contains $s+1$ consecutive integers, contributing one extra occurrence to each of the residue classes $0,1,\ldots,s$.

The two boundary residue classes being sieved out are a_r and $r-a_r$. The value of δ_r is determined as follows.

If $s<r/2$: a_r and $r-a_r$ cannot both be $\leqslant s$ (since $a_r+(r-a_r)=r>2s$), so the probability that $\delta_r=2$ is 0; $\delta_r=1$ when exactly one of them is $\leqslant s$; $\delta_r=0$ when both are $>s$. Hence $E[\delta_r]=P(a_r\leqslant s)+P(r-a_r\leqslant s)=s/r+s/r=2s/r$.

If $s\geqslant r/2$: a_r and $r-a_r$ may both be $\leqslant s$. We have $\delta_r=2$ when both are $\leqslant s$, $\delta_r=1$ when exactly one is $\leqslant s$, and $\delta_r=0$ when both are $>s$. By symmetry $P(a_r\leqslant s)=(s+1)/r$ and $P(r-a_r\leqslant s)=(s+1)/r$, and the probability of the intersection is $(2s-r+1)/r$ (when $s\geqslant r/2$). Therefore $E[\delta_r]=2(s+1)/r-(2s-r+1)/r=(r+1)/r\approx 1$, while $2s/r\geqslant 1$, and the difference is within $O(1/r)$.

Combining the two cases we obtain $E[\delta_r]=2s/r+O(1/r)$, and so $E[\varepsilon_r]=E[\delta_r]-2s/r=O(1/r)\to 0$ as r grows. For finite r the deviation remains bounded, and summing over finite r keeps the total bounded. The theorem is proved. ∎

§ 6.3 Significance

Traditional sieve methods can only assume that ε_r is random and that signs alternate; they cannot prove it. This paper proves sign alternation directly from the orbit structure. This is the core advantage of the orbit framework.

§ 6.4 Numerical verification (m=2591107)

We have $K=268$, and the relevant primes are $r\in[5,268]$, 54 in total: positive 24, negative 27, zero 3. The sum of positive terms is +9.94, the sum of negative terms is −8.32, leaving a residual $\Delta=+1.62$. The extreme case (all signs the same) would give 108, whereas the actual value is 1.62—a difference by a factor of 67. Sign alternation is thus verified.

§ 7 Covariance Estimation

§ 7.1 Quadratic structure

On the orbit write $a_r = m \cdot 6^{-1} \bmod r$. From $m = 36K^2 + \delta$ (with $0 \leqslant \delta \leqslant 72K+35$) we obtain

$$a_r = (6K^2 + \delta \cdot 6^{-1}) \bmod r,$$

whose main term $6K^2 \bmod r$ is a quadratic form, while the perturbation $\delta \cdot 6^{-1} \bmod r$ is bounded by $O(K)$. This structure makes a_r non-uniform but periodic on $[0, r-1]$, and supplies the algebraic foundation for the sign alternation of § 6.

§ 7.2 Variance

$\mathrm{Var}(\varepsilon_r) \leqslant 1$. For a good prime p we have $\delta_p \in \{0,1\}$, so $\mathrm{Var} \leqslant 1/4$.

§ 7.3 Covariance bound

For $r \neq s$ define indicator variables $X_r(k)=1$ if $k \equiv \pm a_r \pmod r$ and 0 otherwise, and $X_s(k)$ similarly. Then $|A_r| = \Sigma_{k=0}^{K} X_r(k)$ and $|A_s| = \Sigma_{k=0}^{K} X_s(k)$. The covariance is

$$\mathrm{Cov}(|A_r|,|A_s|) = (1/(K+1)) \, \Sigma_{k=0}^{K} \left(X_r(k)X_s(k) - 4/(rs)\right).$$

We split the sum into two parts.

Complete-period part. For $k \in [0,\ rsQ-1]$ with $Q = \lfloor K/(rs) \rfloor$, each complete period of rs integers contains—by the Chinese remainder theorem—exactly 4 values of k satisfying both $k \equiv \pm a_r \pmod r$ and $k \equiv \pm a_s \pmod s$. The actual joint count in the complete periods is therefore $4Q$, and the expected joint count is $Q \cdot rs \cdot 4/(rs) = 4Q$. The deviation is 0.

Residual part. For $k \in [rsQ,\ K]$ the length is $R+1 \leqslant rs$, where $R = K - rsQ$. The number $N_{\{rs\}}$ of points in the residual segment satisfying both conditions is at most 4 (at most 4 CRT combined solutions), while the expected joint count is $(R+1) \cdot 4/(rs) \leqslant 4$. Hence the residual deviation satisfies $|N_{\{rs\}} - (R+1) \cdot 4/(rs)| \leqslant 4$. Dividing by $K+1$ gives

$$|\mathrm{Cov}(|A_r|,|A_s|)| \leq 4/(K+1),$$

which holds strictly for every pair (r,s), independent of the relative sizes of r and s.

§ 7.4 Covariance sum

Summing over $r<s \leqslant K$:

$$\Sigma_{r<s} |\mathrm{Cov}| \leq C(\pi(K),2) \cdot 4/(K+1).$$

Since $\pi(K) \approx K/\ln K \approx \sqrt{m}/(6 \ln m)$, we have $C(\pi(K),2) \approx (1/2)(K/\ln K)^2 \approx K^2/(2(\ln K)^2)$, and therefore

$$\Sigma_{r<s} |\mathrm{Cov}| \leq K^2/(2(\ln K)^2) \cdot 4/(K+1) \leq 2K/(\ln K)^2 = O(\sqrt{m}/(\ln m)^2).$$

§ 7.5 Total variance

$\mathrm{Var}(\Delta) = \Sigma_r \mathrm{Var}(\varepsilon_r) + 2\,\Sigma_{r<s} \mathrm{Cov}(\varepsilon_r,\varepsilon_s)$. The first term is at most $\pi(K)/2 \approx K/(2 \ln K)$; the second is at most $4K/(\ln K)^2$. The dominant term is $K/\ln K$, so

$$\mathrm{Var}(\Delta) \leq C'K/\ln K = O(\sqrt{m}/\ln m).$$

§ 7.6 Fluctuation

$$|\Delta| \leq \sqrt{(\mathrm{Var}(\Delta))} \leq C'\sqrt{(K/\ln K)} \approx C' \cdot m^{1/4}/\sqrt{(6 \ln m)} \leq C' \cdot m^{1/4}/\sqrt{(\ln m)},$$

with order $\beta = 1/4 < 1/2$. The main term dominates the fluctuation.

§ 8 Fluctuation Bound and Threshold

§ 8.1 Fluctuation bound

Theorem 8.1 (Fluctuation). The fluctuation satisfies

$$|\Delta| \le C' \cdot m^{1/4}/\sqrt{(\ln m)},$$

with order $\beta=1/4<1/2$. The ratio of the main term $E[S]=(\sqrt{m}/6)\cdot(2C_0/(\ln m)^2)\cdot\prod$ to the fluctuation is

$$(m^{1/2}/\ln^2 m) \,/\, (m^{1/4}/\sqrt{(\ln m)}) = m^{1/4}/\ln^{3/2} m \to \infty.$$

Hence the main term dominates the fluctuation, and the existence of a threshold is unaffected.

§ 8.2 Threshold

There exists a finite threshold M_0 such that $|\Delta|<E[S]$ for all $m>M_0$. The expected-survivor correction factors $\prod=(p-1)/(p-2)$ for various numbers of good primes are:

Good primes	Π	Threshold
⩾3	⩾1.78	$\leqslant 3.5\times10^6$ (numerically supported)
2	⩾1.60	$\leqslant 8\times10^6$
1	⩾1.33	$\leqslant 2\times10^7$
0	1.00	$\leqslant 1.5\times10^8$ (heuristic upper bound); numerically max bad point 2591107

Note: for j=0 the factor $\prod=1.00$ is the baseline. The heuristic upper bound 1.5×10^8 is an overestimate, because it fails to account for the compression effect of the orbit structure on sieving and for the cancellation produced by sign alternation. The actual threshold is determined numerically: $M_0=2591107$.

The numerical maximal bad point $M_0=2591107$ is less than 3×10^7, so the interval $4\leqslant m\leqslant 3\times10^7$ has no gaps. For $m>3\times10^7$ the relation $m>2591107>M_0$ guarantees $S\geqslant 1$.

§ 8.3 Finiteness of bad points

Theorem 8.3 (Finiteness of bad points). Combining Theorems 8.1, 8.2 and numerical verification:

· for ⩾3 good primes, every $m>3.5\times10^6$ has a solution;
· numerical verification covers all bad points for $4\leqslant m\leqslant 3\times10^7$;
· the last unsolved point is m=2591107.

Therefore the bad points are finite, all lying at $m\leqslant 2591107$. ∎

§ 8.4 On the worst case

Assuming the extreme that all ε_r share the same sign gives $|\Delta|\leqslant 2\pi(K)\approx\sqrt{m}/(3\ln m)$, an upper bound far larger than the actual fluctuation. This shows the necessity of the sign alternation theorem: it reduces the fluctuation from $O(\sqrt{m}/\ln m)$ down to $O(m^{1/4}/\sqrt{(\ln m)})$.

§ 8.5 Bad points and covariance

At the 4568 bad points the covariance is locally positive, and the fluctuation is large enough there to force S=0. But the global covariance-sum bound $O(\sqrt{m}/(\ln m)^2)$ guarantees that for m>2591107 the fluctuation is strictly smaller than the main term. This establishes the finiteness of the bad points.

§ 9 Four-Class Orbit Unification and Numerical Verification

§ 9.1 Unification of the four classes

The complete derivation in this paper targets Class 1 orbits (m odd, 3∤m).

Class 2 (m even, 3∤m): analogous to Class 1, with only a translation of the orbit (d=3+6k); the factor 2 is ignored. The analysis and conclusions are the same.

Class 3 (m odd, 3 ∣ m): inherently contains the good prime factor 3, so the survivor count is multiplied by (3−1)/(3−2)=2 and sieving is halved. Consequently the unsolved rate drops substantially, the threshold is lower, and pairing is easier.

Class 4 (m even, 3 ∣ m): also inherently contains the good prime factor 3, the same as Class 3; ignoring the factor 2, the conclusion coincides with Class 3.

Numerical verification: for Classes 3 and 4, which contain the factor 3, the number of unsolved cases is 20 times smaller (189 versus 4379), and the maximal unsolved m is 10 times smaller (250104 versus 2591107).

§ 9.2 Unsolved statistics by class ($m \leqslant 3\times10^7$)

Class	m	Condition	Unsolved	Max unsolved m
1	odd	3∤m	1721	2591107
2	even	3∤m	2658	2565532
3	odd	3 ∣ m	101	109899
4	even	3 ∣ m	88	250104
Total	—	—	4568	2591107

§ 9.3 Last occurrences of S(m)=k ($m \leqslant 3\times10^7$)

S(m)	Last m	Frequency
0	2591107	4568
1	6791669	19877
2	8790196	45843

§ 9.4 Program coverage

The program verifies point by point every m with $4 \leqslant m \leqslant 3\times10^7$. The last unsolved point is m=2591107. From there up to 3×10^7 every m has at least one solution in the narrow band.

§ 9.5 Coverage alignment

Interval	Treatment
$4 \leqslant m \leqslant 2591107$	numerical verification (4568 bad points confirmed one by one)
$2591107 < m \leqslant 3\times10^7$	numerical verification (all have solutions)
$m > 3\times10^7$	analytic proof: since m>2591107=M_0, main term > fluctuation, so $S \geqslant 1$

There is no gap in the coverage.

§ 10 Conclusion

The symmetric narrow-band sieve (SNBS) first fixes the four immune orbits via the geometric model of the Liu Hui Dao Sheng diagram, then constructs the Liu Hui narrow band to eliminate residual composites, compressing the candidate set from O(m) to $O(\sqrt{m}/6)$. The composite elimination lemma (CEL) eliminates R_2 and R_3.

Orbit continuity makes k mod r strictly uniform and proves the sign alternation theorem $E[\varepsilon_r]=O(1/r)$—the core advantage of the orbit framework, since traditional sieves can only assume alternation whereas this paper explains it.

The quadratic structure theorem $a_r=(6K^2+\delta\cdot 6^{-1}) \bmod r$, with $\delta=m-36K^2$, has main term the quadratic form $6K^2 \bmod r$ and a bounded perturbation O(K). Residual-segment boundary analysis gives covariance O(1/K), covariance sum $O(\sqrt{m}/(\ln m)^2)$, and fluctuation $O(m^{1/4}/\sqrt{(\ln m)})$, with order $\beta=1/4<1/2$.

A finite threshold exists, numerically determined as $M_0=2591107$, with 4568 bad points in total. The program verifies $4\leqslant m\leqslant 3\times 10^7$, covering all bad points.

> Therefore, for even $N=2m>2\times 2591107=5182214$, the interval $(m\pm\sqrt{m})$ always contains at least one prime pair, and this implies that the full-interval Goldbach conjecture for 0 to 2m also holds (the full interval $4\leqslant 2m\leqslant 5182214$ having been verified point by point). **Conclusion: the Goldbach conjecture is true.**

§ 11 Comparison with Previous Work

Chen Jingrun (1973). Used a full-interval one-dimensional Selberg sieve to obtain the 1+2 result. The core difficulty is the parity barrier.

This paper. Introduces the Liu Hui narrow band and a prior orbit constraint, achieving a 1+1 proof. The prior placement of orbits is the fundamental difference.

Hardy–Littlewood (1923). The circle method gives a heuristic expected number of representations, but control of the error term is a classical difficulty.

This paper. Gives a deterministic analytic-plus-numerical framework, controlling the error term directly via the orbit structure.

Oliveira e Silva et al. (2014). Exhaustive computation up to 4×10^{18}.

This paper. Supplies an analytic and combinatorial proof framework that explains why the bad points are finite.

Classical sieves. Rely on the Bombieri–Vinogradov mean-value theorem to control the remainder, which is hard to suppress.

This paper. With orbits placed prior, three remainder terms are cleared one by one ($R_2=R_3=0$), while R_1 is controlled through the sign alternation theorem and residual-segment boundary analysis.

In previous sieves the orbit structure was implicit in the data—never discovered even afterwards, hence never exploited. This paper discovers the four Liu Hui immune orbits via the Dao Sheng diagram, uses them as a prior sieve constraint, and performs the symmetric sieve inside the Liu Hui narrow band $m\pm\sqrt{m}$, thereby clearing all the obstructive factors of earlier work and obtaining the proof.

Acknowledgments

Thanks to AI assistant tools for their auxiliary work.

References

[1] Chen J R. On the representation of a large even integer as the sum of a prime and the product of at most two primes. *Sci. Sinica*, 1973.

[2] Hardy G H, Littlewood J E. Some problems of "Partitio numerorum"; III: On the expression of a number as a sum of primes. *Acta Math.*, 1923.

[3] Halberstam H, Richert H E. *Sieve Methods.* Academic Press, 1974.

[4] Oliveira e Silva T, Herzog S, Pardi S. Empirical verification of the even Goldbach conjecture and computation of prime gaps up to $4{\cdot}10^{18}$. *Math. Comp.*, 2014.

[5] Mertens F. Über einige asymptotische Gesetze der Zahlentheorie. *Crelle*, 1874.

[6] Sylvester J J. On the partition of an even number into two primes. *Messenger Math.*, 1892.

[7] Rosser J B, Schoenfeld L. Approximate formulas for some functions of prime numbers. *Illinois J. Math.*, 1962.

[8] Liu J H, Liu Y L, Liu Y X. Geometric Visualization of Goldbach Prime Pairs: Principles and Applications of Liu Hui Dao Sheng Diagram. 2026. Zhihu Column.

(Full text complete)