

# Quantile Regression in the Presence of Monotone Missingness with Sensitivity Analysis

**SUMMARY:** In this paper, we consider the problem of longitudinal quantile regression using pattern mixture models when there is monotone missingness. Our proposed pattern mixture models specify the marginal quantile regression parameters, as well as the conditional distribution, thus allowing sensitivity analysis, a key component in inference for incomplete data. For computation of the likelihood of the parametric model, we proposed a novel way to simplify the integral. Both frequentist and Bayesian methods are illustrated. The model is applied on a clinical trial study for weight management.

**KEY WORDS:** Monotone missingness; Nonignorable missingness; Quantile regression; Sensitivity analysis.

## 1. Introduction

Quantile regression is used to study the relationship between a response and covariates when one (or several) quantiles are of interest as opposed to mean regression. The dependence between upper or lower quantiles of the response variable and the covariates often vary differentially relative to that of the mean. How quantiles depend on covariates is of interest in econometrics, educational studies, biomedical studies, and environment studies (Yu and Moyeed, 2001; Buchinsky, 1994, 1998; He et al., 1998; Koenker and Machado, 1999; Wei et al., 2006; Yu et al., 2003). A comprehensive review of applications of quantile regression was presented in Koenker (2005).

Quantile regression is more robust to outliers than mean regression and provides information about how covariates affect quantiles, which offers a more complete description of the conditional distribution of the response. Different effects of covariates can be assumed for different quantiles.

The traditional frequentist approach was proposed by Koenker and Bassett (1978) for a single quantile with estimators derived by minimizing a loss function. The popularity of this approach is due to its computational efficiency, well-developed asymptotic properties, and straightforward extensions to simultaneous quantile regression and random effect models. However, asymptotic inference may not be accurate for small sample sizes and the approach does not naturally extend to missing data.

Bayesian approaches offer exact inference in small samples. Motivated by the loss (check) function, Yu and Moyeed (2001) proposed an asymmetric Laplace distribution for the error term, such that maximizing the posterior distribution is equivalent to minimizing the check function. Also semiparametric methods have been proposed for median regression. Walker and Mallick (1999) used a diffuse finite Pólya Tree prior for the error term. Kottas and Gelfand (2001) modeled the error by two families of median zero distribution using a mixture

Dirichlet process priors, which is very useful for unimodal error distributions. Hanson and Johnson (2002) adopted mixture of Pólya Tree prior in median regression, which is more robust in terms of multimodality and skewness. Other recent approaches include quantile pyramid priors, mixture of Dirichlet process priors of multivariate normal distributions and infinite mixture of Gaussian densities which place quantile constraints on the residuals (Hjort and Petrone, 2007; Hjort and Walker, 2009; Kottas and Krnjajić, 2009; Reich et al., 2010).

The above methods focus on complete data. There are a few articles about quantile regression with missingness. Wei et al. (2012) proposed a multiple imputation method for quantile regression model when there are some covariates missing at random (MAR). They impute the missing covariates by specifying its conditional density given observed covariates and outcomes, which comes from the estimated conditional quantile regression and specification of conditional density of missing covariates given observed ones. However, they put more focus on the missing covariates rather than missing outcomes. Bottai and Zhen (2013) illustrated an imputation method using estimated conditional quantiles of missing outcomes given observed data. Their approach does not make distributional assumptions. They assumed the missing data mechanism (MDM) is ignorable. However, because their imputation method is not derived from a joint distribution, the joint distribution with such conditionals may not exist. In addition, their approach does not allow for MNAR.

Yuan and Yin (2010) introduced a fully parametric Bayesian quantile regression approach for longitudinal data with nonignorable missing data. They used shared latent subject-specific random effects to explain the within-subject correlation and to associate the response process with missing data process, and applied multivariate normal priors on the random terms to match the traditional quantile regression check function with penalties. However, the quantile regression coefficients are conditional on the random effects, which is not of interest if we are interested in interpreting regression coefficients unconditional on random

effects. In addition, they are conditional on random effects, which tie together the responses and missingness process, so they have slightly different interpretation than regular random effects in longitudinal methods. Moreover, due to their full parametric specification for the full data, their model does not allow for sensitivity analysis, which is a key component in inference for incomplete data (NAS 2010).

Pattern mixture models were originally proposed to model missing data in Rubin (1977). Later mixture models were extended to handle MNAR in longitudinal data. For discrete dropout times, Little (1993, 1994) proposed a general method by introducing a finite mixture of multivariate distribution for longitudinal data. When there are many possible dropout time, Roy (2003) proposed to group them by latent classes.

Roy and Daniels (2008) extended Roy (2003) to generalized linear models and proposed a pattern mixture model for data with nonignorable dropout, borrowing ideas from Heagerty (1999). But their approach only estimates the marginal covariate effects on the mean. We will use related ideas for quantile regression models which allows non-ignorable missingness and sensitivity analysis.

The structure of this article is as follows. First, we introduce a quantile regression method to address monotone nonignorable missingness in section 2, including sensitivity analysis and computational details. We use simulation studies to evaluate the performance of the model in section 3. We apply our approach to data from a recent clinical trial in section 4. Finally, discussion and conclusions are given in section 5.

## 2. Model

In this section, we first introduce some notations, then describe our proposed quantile regression model in section 2.1. We provide details on MAR and MNAR and computation in sections 2.2 and 2.3.

Under monotone dropout, without loss of generality, denote  $S_i \in \{1, 2, \dots, J\}$  to be the

number of observed  $Y'_{ij}$ s for subject  $i$ , and  $\mathbf{Y}_i = (Y_{i1}, Y_{i2}, \dots, Y_{iJ})^T$  to be the full data response vector for subject  $i$ , where  $J$  is the maximum follow up time. We assume  $Y_{i1}$  is always observed. We are interested in the  $\tau$ -th marginal quantile regression coefficients  $\boldsymbol{\gamma}_j = (\gamma_{j1}, \gamma_{j2}, \dots, \gamma_{jp})^T$ ,

$$\Pr(Y_{ij} \leq \mathbf{x}_i^T \boldsymbol{\gamma}_j) = \tau, \text{ for } j = 1, \dots, J,$$

where  $\mathbf{x}_i$  is a  $p \times 1$  vector of covariates for subject  $i$ .

Let

$$p_k(Y) = p(Y|S = k), \quad p_{\geq k}(Y) = p(Y|S \geq k)$$

be the densities of response  $\mathbf{Y}$  given follow-up time  $S = k$  and  $S \geq k$ . And  $\Pr_k$  be the corresponding probability given  $S = k$ .

## 2.1 Mixture Model Specification

We adopt a pattern mixture model to jointly model the response and missingness (Little, 1994; Daniels and Hogan, 2008). Mixture models factor the joint distribution of response and missingness as

$$p(\mathbf{y}, \mathbf{S}, |\mathbf{x}, \boldsymbol{\omega}) = p(\mathbf{y}|\mathbf{S}, \mathbf{x}, \boldsymbol{\omega})p(\mathbf{S}|\mathbf{x}, \boldsymbol{\omega}).$$

Thus the full-data response follows the distribution is given by

$$p(\mathbf{y}|\mathbf{x}, \boldsymbol{\omega}) = \sum_{\mathbf{S} \in \mathcal{S}} p(\mathbf{y}|\mathbf{S}, \mathbf{x}, \boldsymbol{\theta})p(\mathbf{S}|\mathbf{x}, \boldsymbol{\phi}),$$

where  $\mathcal{S}$  is the sample space for dropout time  $S$  and the parameter vector  $\boldsymbol{\omega}$  is partitioned as  $(\boldsymbol{\theta}, \boldsymbol{\phi})$ .

Furthermore, the conditional distribution of response within patterns can be decomposed as

$$p(\mathbf{y}_{obs}, \mathbf{y}_{mis}|\mathbf{S}, \boldsymbol{\theta}) = p(\mathbf{y}_{mis}|\mathbf{y}_{obs}, \mathbf{S}, \boldsymbol{\theta}_E)p(\mathbf{y}_{obs}|\mathbf{S}, \boldsymbol{\theta}_{y,O}), \quad (1)$$

where  $\boldsymbol{\theta}_E$  indexes the parameters in the extrapolation distribution, the first term on the

right hand side and  $\boldsymbol{\theta}_{y,O}$  indexes parameters in the distribution of observed responses, the second term on the right hand side.

We assume models within pattern to be multivariate normal distributions and specify a sequential model parametrization. Let the subscript  $i$  stand for subject  $i$ . First we specify the marginal quantile regression models as:

$$\Pr(Y_{ij} \leq \mathbf{x}_{ij}^T \boldsymbol{\gamma}_j) = \tau,$$

where  $\boldsymbol{\gamma}_j$  is the  $\tau^{th}$  quantile regression coefficients of interest for component  $j$ .

Then we specify the conditional distributions as:

$$\begin{aligned} p_k(y_{i1}) &= N(\Delta_{i1} + \mathbf{x}_{i1}^T \boldsymbol{\beta}_1^{(k)}, \sigma_1^{(k)}), k = 1, \dots, J, \\ p_k(y_{ij} | \mathbf{y}_{ij-}) &= \begin{cases} N(\Delta_{ij} + \mathbf{x}_{ij}^T \mathbf{h}_j^{(k)} + \mathbf{y}_{ij-}^T \boldsymbol{\beta}_{y,j-1}^{(k)}, \sigma_j^{(k)}), & k < j; \\ N(\Delta_{ij} + \mathbf{y}_{ij-}^T \boldsymbol{\beta}_{y,j-1}^{(\geq j)}, \sigma_j^{(\geq j)}), & k \geq j; \end{cases}, \text{ for } 2 \leq j \leq J, \quad (2) \\ S_{ij} = k | \mathbf{x}_{ij} &\sim \text{Multinomial}(1, \boldsymbol{\phi}), \end{aligned}$$

where  $\mathbf{y}_{ij-} = (y_{i1}, \dots, y_{i(j-1)})^T$  is the response history for subject  $i$  up to time point  $(j-1)$ ;  $\boldsymbol{\phi} = (\phi_1, \dots, \phi_J)$  is the multinomial probability vector for the number of observed responses;  $\mathbf{h}_j^{(k)} = (h_{j1}^{(k)}, \dots, h_{jp}^{(k)})$  are sensitivity parameters which we will discuss in section 2.2.  $\mathbf{x}_{ij}$  is a  $p \times 1$  covariate vector;  $\boldsymbol{\beta}_{y,j-1}^{(k)} = (\beta_{y1,j-1}^{(k)}, \dots, \beta_{y_{j-1},j-1}^{(k)})^T$  are autoregressive coefficients and  $\sigma_j^{(k)}$  is the conditional standard deviation of response component  $j$ . We specify the model as in (2) to have multivariate normal distribution within patterns such that MAR exists (Wang and Daniels, 2011). More details are presented in section 2.2.

In (2),  $\Delta_{ij}$  are functions of  $\tau, \mathbf{x}_{ij}, \boldsymbol{\beta}, \mathbf{h}, \boldsymbol{\sigma}, \boldsymbol{\gamma}_j, \boldsymbol{\phi}$  and are determined by the marginal quantile regressions,

$$\tau = \Pr(Y_{ij} \leq \mathbf{x}_{ij}^T \boldsymbol{\gamma}_j) = \sum_{k=1}^J \phi_k \Pr_k(Y_{ij} \leq \mathbf{x}_{ij}^T \boldsymbol{\gamma}_j) \text{ for } j = 1, \quad (3)$$

and

$$\begin{aligned}
\tau &= \Pr(Y_{ij} \leq \mathbf{x}_{ij}^T \boldsymbol{\gamma}_j) = \sum_{k=1}^J \phi_k \Pr_k(Y_{ij} \leq \mathbf{x}_{ij}^T \boldsymbol{\gamma}_j) \\
&= \sum_{k=1}^J \phi_k \int \cdots \int \Pr_k(Y_{ij} \leq \mathbf{x}_{ij}^T \boldsymbol{\gamma}_j | \mathbf{y}_{ij-}) \Pr_k(y_{i(j-1)} | \mathbf{y}_{i(j-1)-}) \\
&\quad \cdots \Pr_k(y_{i2} | y_{i1}) \Pr_k(y_{i1}) dy_{i(j-1)} \cdots dy_{i1}. \text{ for } j = 2, \dots, J.
\end{aligned} \tag{4}$$

Computational details will be given in section 2.3.

The idea is to model the marginal quantile regressions directly, then to embed them in the likelihood through restrictions in the mixture model. The mixture model in (2) allows the marginal quantile regression coefficients to differ by quantiles. Otherwise, the quantile lines would be parallel to each other. Moreover, the mixture model also allows sensitivity analysis.

For identifiability of the observed data distribution, we apply the following restrictions,

$$\sum_{k=1}^J \beta_{l1}^{(k)} = 0, l = 1, \dots, p,$$

## 2.2 Missing Data Mechanism and Sensitivity Analysis

In general, mixture models are not identified due to insufficient information provided by observed data. Specific forms of missingness are needed to induce constraints to identify the distributions for incomplete patterns, in particular, the extrapolation distribution in (1). In this section, we explore ways to embed the missingness mechanism and sensitivity parameters in mixture models for our setting.

In the mixture model in (2), MAR holds (Molenberghs et al., 1998; Wang and Daniels, 2011) if and only if, for each  $j \geq 2$  and  $k < j$ :

$$\Pr_k(y_j | y_1, \dots, y_{j-1}) = \Pr_{\geq j}(y_j | y_1, \dots, y_{j-1}).$$

When  $2 \leq j \leq J$  and  $k < j$ ,  $Y_j$  is not observed, thus  $\mathbf{h}_j^{(k)}$  and  $\sigma_j^{(k)}, \boldsymbol{\beta}_{y,j-1}^{(k)} = (\beta_{y_1,j}^{(k)}, \dots, \beta_{y_{j-1},j-1}^{(k)})^T$

can not be identified from the observed data. Denote

$$\log \sigma_j^{(k)} = \log \sigma_j^{(\geq j)} + \delta_j^{(k)},$$

$$\beta_{y,j-1}^{(k)} = \beta_{y,j-1}^{(\geq j)} + \eta_{j-1}^{(k)},$$

where  $\eta_{j-1}^{(k)} = (\eta_{y_1,j-1}^{(k)}, \dots, \eta_{y_{j-1},j-1}^{(k)})$  for  $k < j$ . Then  $\xi_s = (\mathbf{h}_j^{(k)}, \eta_{j-1}^{(k)}, \delta_j^{(k)})$  is a set of sensitivity parameters (Daniels and Hogan, 2008), where  $k < j, 2 \leq j \leq J$ .

When  $\xi_s = \xi_{s0} = \mathbf{0}$ , MAR holds. If  $\xi_s$  is fixed at  $\xi_s \neq \xi_{s0}$ , the missingness mechanism is MNAR. We can vary  $\xi_s$  around  $\mathbf{0}$  to examine the impact of different MNAR mechanisms.

For fully Bayesian inference, we can put priors on  $(\xi_s, \xi_m)$  as :

$$p(\xi_s, \xi_m) = p(\xi_s)p(\xi_m),$$

where  $\xi_m = (\gamma_j, \beta_{y,j-1}^{(\geq j)}, \alpha_j^{(\geq j)}, \phi)$ , the identified parameters in the data distribution. If we assume MAR with no uncertainty, the prior of  $\xi_s$  is  $p(\xi_s = \mathbf{0}) \equiv 1$ . Sensitivity analysis can be executed by putting point mass priors on  $\xi_s$  to examine the effect of priors on the posterior inference about quantile regression coefficients  $\gamma_j^\tau$ . For example, if MAR is assumed with uncertainty, priors can be assigned as  $E(\xi_s) = \xi_{s0} = \mathbf{0}$  with  $\text{Var}(\xi_s) \neq \mathbf{0}$ . If we assume MNAR with no uncertainty, we can put priors satisfying  $E(\xi_s) = \Delta_\xi$ , where  $\Delta_\xi \neq \mathbf{0}$  and  $\text{Var}(\xi_s) = \mathbf{0}$ . If MNAR is assumed with uncertainty, then priors could be  $E(\xi_s) = \Delta_\xi$ , where  $\Delta_\xi \neq \mathbf{0}$  and  $\text{Var}(\xi_s) \neq \mathbf{0}$ .

In general, each pattern  $S = k$  has its own set of sensitivity parameters  $\xi_s^{(k)}$ . However, to keep the number of sensitivity parameters at a manageable level (Daniels and Hogan, 2008) and without loss of generality, we assume  $\xi_s$  does not depend on pattern.

### 2.3 Computation

In section 2.3.1, we provide details on calculating  $\Delta_{ij}$  in (2) for  $j = 1, \dots, J$ . Then we show how to obtain maximum likelihood estimates in section 2.3.2. Finally, we present a Monte Carlo Markov Chain (MCMC) sampling algorithm for Bayesian inference in section 2.3.4.



2.3.1 *Calculation of  $\Delta$*  . From equation (3) and (4),  $\Delta_{ij}$  depends on subject-specific covariates  $\mathbf{x}_{ij}$ , thus  $\Delta_{ij}$  needs to be calculated for each subject. We now illustrate how to calculate  $\Delta_{ij}$  given all the other parameters  $\boldsymbol{\xi} = (\boldsymbol{\xi}_m, \xi_s)$ .

- $\Delta_{i1}$  : Expand equation (3):

$$\tau = \sum_{k=1}^J \phi_k \Phi \left( \frac{\mathbf{x}_{i1}^T \boldsymbol{\gamma}_1 - \Delta_{i1} - \mathbf{x}_{i1}^T \boldsymbol{\beta}_1^{(k)}}{\sigma_1^{(k)}} \right),$$

where  $\Phi$  is the standard normal CDF. Because the above equation is continuous and monotone in  $\Delta_{i1}$ , it can be solved by a standard numerical root-finding method (e.g. bisection method) with minimal difficulty.

- $\Delta_{ij}, 2 \leq j \leq J$  :

First we introduce a lemma:

LEMMA 2.1: *An integral of a normal CDF with mean  $b$  and standard deviation  $a$  over another normal distribution with mean  $\mu$  and standard deviation  $\sigma$  can be simplified to a closed form in terms of normal CDF:*

$$\int \Phi \left( \frac{x - b}{a} \right) d\Phi(x; \mu, \sigma) = \begin{cases} 1 - \Phi \left( \frac{b - \mu}{\sigma} / \sqrt{\frac{a^2}{\sigma^2} + 1} \right) & a > 0, \\ \Phi \left( \frac{b - \mu}{\sigma} / \sqrt{\frac{a^2}{\sigma^2} + 1} \right) & a < 0, \end{cases}$$

where  $\Phi(x; \mu, \sigma)$  stands for a CDF of normal distribution with mean  $\mu$  and standard deviation  $\sigma$ .

Given the result in Lemma 2.1, to solve equation (4), we propose a recursive approach. For

the first multiple integral in equation (4), apply lemma 2.1 once to obtain:

$$\begin{aligned}
\Pr_1(Y_{ij} \leq \mathbf{x}_{ij}^T \boldsymbol{\gamma}_j) &= \int \dots \int \Pr(Y_{ij} \leq \mathbf{x}_{ij}^T \boldsymbol{\gamma}_j | S = 1, \mathbf{x}_{ij}, \mathbf{Y}_{ij-}) \\
&\quad dF(Y_{i(j-1)} | S = 1, \mathbf{x}_{ij}, \mathbf{Y}_{i(j-1)-}) \dots dF(Y_{i1} | S = 1, \mathbf{x}_{ij}), \\
&= \int \dots \int \Phi \left( \frac{\mathbf{x}_{ij}^T \boldsymbol{\gamma}_j - \mu_{j|1, \dots, j-1}(\mathbf{Y}_{ij-})}{\sigma_{j|1, \dots, j-1}} \right) \\
&\quad dF(Y_{i(j-1)} | S = 1, \mathbf{x}_{ij}, \mathbf{Y}_{i(j-1)-}) \dots dF(Y_{i1} | S = 1, \mathbf{x}_{ij}), \\
&= \int \dots \int \Phi \left( \frac{Y_{i(j-2)} - b^*}{a^*} \right) dF(Y_{i(j-2)} | S = 1, \mathbf{x}_{ij}, \mathbf{Y}_{i(j-2)-}) \\
&\quad \dots dF(Y_{i1} | S = 1, \mathbf{x}_{ij}).
\end{aligned}$$

Then, by recursively applying lemma 2.1 ( $j - 1$ ) times, each multiple integral in equation (4) can be simplified to single normal CDF. Thus we can easily solve for  $\Delta_{ij}$  using standard numerical root-finding method as for  $j = 1$ .

**2.3.2 Maximum Likelihood Estimation.** The observed data likelihood for an individual  $i$  with follow-up time  $S_i = k$  is

$$\begin{aligned}
L_i(\boldsymbol{\xi} | \mathbf{y}_i, S_i = k) &= \phi_k p_k(y_k | y_1, \dots, y_{k-1}) p_k(y_{k-1} | y_1, \dots, y_{k-2}) \dots p_k(y_1), \\
&= \phi_k p_{\geq k}(y_k | y_1, \dots, y_{k-1}) p_{\geq k-1}(y_{k-1} | y_1, \dots, y_{k-2}) \dots p_k(y_1),
\end{aligned} \tag{5}$$

where  $\mathbf{y}_i = (y_1, \dots, y_k)$ .

We use derivative-free optimization algorithms by quadratic approximation to compute the maximum likelihood estimates (Bates et al., 2012). Denote  $J(\boldsymbol{\xi}) = -\log L = -\log \sum_{i=1}^n L_i$ . Then maximizing the likelihood is equivalent to minimizing the target function  $J(\boldsymbol{\xi})$ . Under an MAR assumption, we fix  $\boldsymbol{\xi}_s = \mathbf{0}$ , while under MNAR assumption,  $\boldsymbol{\xi}_s$  can be chosen as desired.

During each step of the algorithm,  $\Delta_{ij}$  has to be calculated for each subject and at each time, as well as partial derivatives for each parameter.

As an example of the speed of the algorithm, for 100 bivariate outcomes and 5 covariates, it

takes about 1.9 seconds to get convergence using R version 2.15.3 (2013-03-01) (R Core Team, 2013) and platform: x86\_64-apple-darwin9.8.0/x86\_64 (64-bit). Main parts of the algorithm are coded in Fortran such as calculation of numerical derivatives and log-likelihood to quicken computations. We have incorporated those functions implementing the algorithm into the new R (R Core Team, 2013) package “qrmissing”.

We use the bootstrap (Efron, 1979; Efron and Tibshirani, 1993; Davison and Hinkley, 1997) to construct confidence interval and make inferences. We resample subjects and use bootstrap percentile intervals to form confidence intervals.

**2.3.3 Goodness of Fit Check.** A simple goodness-of-fit check can be done by examining normal QQ plots of the fitted residuals from the model. The visual test can help to diagnose if the parametric assumptions are suitable for model.

After obtaining the MLE, we use the approach described in section 2.3.1 to get the fitted  $\Delta_{ij}$  for each subject. Then the fitted residuals can be obtained by plugging in the fitted estimates and  $\hat{\Delta}_{ij}$  to obtain,

$$\hat{\epsilon}_{ij} = \begin{cases} (y_{ij} - \hat{\Delta}_{ij} - \mathbf{x}_{ij}^T \hat{\boldsymbol{\beta}}_1^{(k)}) / \hat{\sigma}_1^{(k)}, & j = 1 \\ (y_{ij} - \hat{\Delta}_{ij} - \mathbf{y}_{ij-1}^T \hat{\boldsymbol{\beta}}_{\mathbf{y},j-1}^{(\geq j)}) / \hat{\sigma}_j^{(\geq j)}, & j > 1 \end{cases}.$$

**2.3.4 Bayesian Framework.** For Bayesian inference, we specify priors on the parameters  $\boldsymbol{\xi}$  and use a block Gibbs sampling method to draw samples from the posterior distribution. Denote all the parameters to sample as :

$$\begin{aligned} \boldsymbol{\xi}_m &= \left\{ \boldsymbol{\gamma}_j, \boldsymbol{\beta}_{\mathbf{y},j-1}^{(\geq j)}, \boldsymbol{\alpha}_j^{(\geq j)} \right\} \text{ for } j = 1, \dots, J, \\ \boldsymbol{\xi}_s &= \left\{ \mathbf{h}_j^{(k)}, \boldsymbol{\eta}_{j-1}^{(k)}, \delta_j^{(k)} \right\} \text{ for } k = 1, \dots, j-1; 2 \leq j \leq J. \end{aligned}$$

Comma separated parameters are marked to sample as a block. Updates of  $\boldsymbol{\xi}_m$  require a Metropolis-Hasting algorithm, while  $\boldsymbol{\xi}_s$  samples are drawn directly from priors as desired for missingness mechanism assumptions.

As mentioned in section 2.2, MAR or MNAR assumptions are implemented via specific priors. For example, if MAR is assumed with no uncertainty, then  $\xi_s = \mathbf{0}$  with probability

1. Details for updating parameters are:

- $\gamma_1$ : Use Metropolis-Hasting algorithm.

(1) Draw  $(\gamma_1^c)$  candidates from candidate distribution;

(2) Based on the new candidate parameter  $\xi^c$ , calculate candidate  $\Delta_{ij}^c$  for each subject  $i$  as we described in section 2.3.1.

(3) Plug in  $\Delta_{ij}^c$  in likelihood (5) to get candidate likelihood;

(4) Compute Metropolis-Hasting ratio, and accept the candidate value or keep the previous value.

- For the rest of the identifiable parameters, algorithms for updating the samples are all similar to  $\gamma_j$ .
- For sensitivity parameters, because we do not get any information from the data, we sample them from priors, which are specified based on assumptions about the missingness.

### 3. Simulation Study

In this section, we compare the performance of our proposed model in section 2.1 with the *rq* function (noted as RQ) in *quantreg* R package (Koenker, 2012) and Bottai's algorithm (Bottai and Zhen, 2013) (noted as BZ). The *rq* function minimizes the loss (check) function  $\sum_{i=1}^n \rho_\tau(y_i - \mathbf{x}_i^T \boldsymbol{\beta})$  in terms of  $\boldsymbol{\beta}$ , where the loss function  $\rho_\tau(u) = u(\tau - I(u < 0))$  and does not make any distributional assumptions. Bottai and Zhen (2013) impute missing outcomes using the estimated conditional quantiles of missing outcomes given observed data. Their approach does not make distributional assumptions similar to *rq* and assumes ignorable missing data.

We considered three scenarios corresponding to both MAR and MNAR assumptions for

a bivariate response. In the first scenario,  $Y_2$  were missing at random and we used the MAR assumption in our algorithm. In the next two scenarios,  $Y_2$  were missing not at random. However, in the second scenario, we misspecified the MDM for our algorithm and still assumed MAR, while in the third scenario, we used the correct MNAR MDM. For each scenario, we considered three error distributions: normal, student t distribution with 3 degrees of freedom and Laplace distribution. For each error model, we simulated 100 data sets. For each set there are 200 bivariate observations  $\mathbf{Y}_i = (Y_{i1}, Y_{i2})$  for  $i = 1, \dots, 200$ .  $Y_{i1}$  were always observed, while some of  $Y_{i2}$  were missing. A single covariate  $x$  was sampled from  $\text{Uniform}(0,2)$ . The three models for the full data response  $\mathbf{Y}_i$  were:

$$Y_{i1}|R = 1 \sim 2 + x_i + \epsilon_{i1},$$

$$Y_{i1}|R = 0 \sim -2 - x_i + \epsilon_{i1},$$

$$Y_{i2}|R = 1, Y_{i1} \sim 1 - x_i - 1/2Y_{i1} + \epsilon_{i2},$$

where  $\epsilon_{i1}, \epsilon_{i2} \stackrel{\text{i.i.d}}{\sim} N(0, 1)$ ,  $t_3$  or  $\text{LP}(\text{rate} = 1)$  distribution within each scenario.

For all cases,  $\Pr(R = 1) = 0.5$ . When  $R = 0$ ,  $Y_{i2}$  is not observed, so  $p(Y_{i2}|R = 0, Y_{i1})$  is not identifiable from observed data.

In the first scenario,  $Y_2$  is missing at random, thus  $p(Y_{i2}|R = 0, Y_{i1}) = p(Y_{i2}|R = 1, Y_{i1})$ . In the last two scenarios,  $Y_2$  are missing not at random. We assume  $Y_{i2}|R = 0, Y_{i1} \sim 3 - x_i - 1/2Y_{i1} + \epsilon_{i2}$ . Therefore, there is a shift of 2 in the intercept between  $p(Y_2|R = 1, Y_1)$  and  $p(Y_2|R = 0, Y_1)$ .

Under an MAR assumption, the sensitivity parameter  $\boldsymbol{\xi}_s$  is fixed at  $\mathbf{0}$  as discussed in section 2.2. For  $rq$  function from *quantreg* R package, because only  $Y_{i2}|R = 1$  is observed, the quantile regression for  $Y_{i2}$  can only be fit from the information of  $Y_{i2}|R = 1$  vs  $x$ .

In scenario 2 under MNAR, we mis-specified the MDM using the wrong sensitivity parameter  $\boldsymbol{\xi}_s$  at  $\mathbf{0}$ . In scenario 3, we assumed there was an intercept shift between distribution of  $Y_{i2}|Y_{i1}, R = 1$  and  $Y_{i2}|Y_{i1}, R = 0$ , thus fixed  $\boldsymbol{\xi}_s$  at its true value.

For each dataset, we fit quantile regression for quantiles  $\tau = 0.1, 0.3, 0.5, 0.7, 0.9$ . Parameter estimates were evaluated by mean squared error (MSE),

$$\text{MSE}(\gamma_{ij}) = \frac{1}{100} \sum_{k=1}^{100} \left( \hat{\gamma}_{ij}^{(k)} - \gamma_{ij} \right)^2, i = 0, 1$$

where  $\gamma_j$  is the true value for quantile regression coefficient,  $\hat{\gamma}_j^{(k)}$  is the maximum likelihood estimates in  $k$ -th simulated dataset ( $(\gamma_{01}, \gamma_{11})$  for  $Y_{i1}$ ,  $(\gamma_{02}, \gamma_{12})$  for  $Y_{i2}$ ).

Monte Carlo standard error (MCSE) is used to evaluate the significance of difference between methods. It is calculated by

$$\text{MCSE} = \hat{\text{sd}}(\text{Bias}^2)/\sqrt{N},$$

where  $\hat{\text{sd}}$  is the sample standard deviation and  $\text{Bias} = \hat{\gamma}_{ij} - \gamma_{ij}$  and  $N$  is the number of simulations.

Table 1, 2 and 3 present the MSE for coefficients estimates of quantile 0.1, 0.3, 0.5, 0.7, 0.9 under each scenario. Simulation results show estimates from our algorithm and Bottai's approach are closer to the true value for all quantiles from 0.1 to 0.9. As expected, under normal errors, the proposed methods dominates both *rq* and BZ in most cases for MAR, incorrect MAR, and MNAR.

For the heavier tail distributions,  $t_3$  and Laplace distribution, our approach shows better performance in middle quantiles and lose to *rq* for extreme quantiles for observed data  $Y_1$ . Nevertheless, our algorithm provides larger gains over *rq* function for each marginal quantile for the second component  $Y_2$ , which are missing for some observations. No matter what missing data mechanism (MAR or MNAR), what assumption we use in our approach (misspecification or correct specified), our method shows advantages over *rq* function, especially for  $Y_2$ , because *quantreg* does not consider the missingness mechanism. The difference in MSE becomes larger for the upper quantiles because  $Y_2|R = 0$  tends to be larger than  $Y_2|R = 1$ ; therefore, the *rq* method using only the

observed  $Y_2$  yields larger bias for upper quantiles. Bottai’s approach, however, shows great advantage over *rq* function for missing data because its imputing method for missing responses. It also has smaller MSE than ours on extreme quantiles regression when distribution has heavy tail. However, our approach has advantages on middle quantiles (30% - 70%) for marginal inference on missing responses regardless using mis-specification or correct sensitivity parameters. And we can see more gains over *BZ* in the quantile regression slope estimates for  $Y_2$ .

To assess the goodness of fit, we examined the QQ plot of fitted residuals in model (2) to check the normality assumption on the error term for a random sample of the simulated datasets. When our error assumption is correct (normal), the QQ plot reflects the fitted residuals follow exact a normal distribution. However, when we misspecified the error distribution, the proposed diagnostic method did clearly suggest heavier tail error than normal, and this also demonstrates why our approach has some disadvantages for regression on extreme quantiles when errors are not normal.

[Table 1 about here.]

[Table 2 about here.]

[Table 3 about here.]

#### 4. Real Data Analysis

We apply our quantile regression approach to data from TOURS, a weight management clinical trial (Perri et al., 2008). This trial was designed to test whether a lifestyle modification program could effectively help people to manage their weights in the long term. After finishing the six-month weight loss program, participants were randomly assigned to three treatments groups: face-to-face counseling, telephone counseling and control group. Their weights were

recorded at baseline ( $Y_0$ ), 6 months ( $Y_1$ ) and 18 months ( $Y_2$ ). Here, we are interested in how the distribution of weights at six months and eighteen months change with covariates. The regressors of interest include AGE, RACE (black and white) and weight at baseline ( $Y_0$ ). Weights at the six months ( $Y_1$ ) were always observed and 13 out of 224 observations (6%) were missing at 18 months ( $Y_2$ ). The “Age” covariate was scaled to 0 to 5 with every increment representing 5 years.

We fitted regression models for bivariate responses  $\mathbf{Y}_i = (Y_{i1}, Y_{i2})$  for quantiles (10%, 30%, 50%, 70%, 90%). We ran 1000 bootstrap samples to obtain 95% confidence intervals.

Estimates under MAR and MNAR are presented in Table 4. For weights of participants at six months, weights of whites are generally 4kg lower than those of blacks for all quantiles, and the coefficients of race are negative and significant. Meanwhile, weights of participants are not affected by age since the coefficients are not significant. Difference in quantiles are reflected by the intercept. Coefficients of baseline weight show a strong relationship with weights after 6 months.

[Table 4 about here.]

For weights at 18 months after baseline, we have similar results. Weights after 18 months still have a strong relationship with baseline weights. However, whites do not weigh significantly less than blacks at 18 months unlike at 6 months.

We also did a sensitivity analysis based on an assumption of missing not at random. Based on previous studies of pattern of weight regain after lifestyle treatment (Wadden et al., 2001; Perri et al., 2008), we assume that

$$E(Y_2 - Y_1 | R = 0) = 3.6\text{kg},$$

which corresponds to 0.3kg regain per month after finishing the initial 6-month program. We incorporate the sensitivity parameters in the distribution of  $Y_2 | Y_1, R = 0$  via the following



restriction:

$$\Delta_{i2} + \mathbf{x}_{i2}^T \mathbf{h}_2^{(1)} + E(y_{i1}|R=0)(\beta_{y,1}^{(1)} + \eta_1^{(1)} - 1) = 3.6\text{kg}.$$

Table 4 presents the estimates and bootstrap percentile confidence intervals under the above MNAR mechanism. There are not large differences for estimates for  $Y_2$  under MNAR vs MAR. This is partly due to the low proportion of missing data in this study.

We also checked the goodness of fit via QQ plots on the fitted residuals as described in section 2.3.3 for each quantile regression fit. Plots are presented in Appendix 5. The QQ plots showed minimal evidence against the assumption that the residuals were normally distributed; thus we were confident with the conclusion of our quantile regression models.

## 5. Discussion

In this paper, we have developed a marginal quantile regression model for data with monotone missingness. We use a pattern mixture model to jointly model the full data response and missingness. Here we estimate marginal quantile regression coefficients instead of coefficients conditional on random effects as in Yuan and Yin (2010). In addition, our approach allows non-parallel quantile lines over different quantiles via the mixture distribution and allows for sensitivity analysis which is essential for the analysis of missing data (NAS 2010).

Our method allows the missingness to be non-ignorable. We illustrated how to put informative priors for Bayesian inference and how to find sensitivity parameters to allow different missing data mechanisms in general. The recursive integration algorithm simplifies computation and can be easily implemented even in high dimensions. Simulation studies demonstrates that our approach has smaller MSE than the traditional frequentist method  $rq$  function for most cases, especially for inferences of partial missing responses. And it has advantages over Bottai’s approach for middle quantiles regression inference even when the distribution is mis-specified. However, our approach also shows little bias for extreme

quantiles comparing to Bottai and Zhen (2013) if error is mis-specified. We also illustrated sensitivity analysis and how to allow non-ignorable missingness assumptions.

Our model assumes a multivariate normal distribution for each component in the pattern mixture model, which might be too restrictive in some settings. Simulation results showed that the mis-specification of the error term did have a small impact on the extreme quantile regression inferences. It is possible to replace that with a semi-parametric model, for example, a Dirichlet process mixture or Pólya trees. However, computational algorithms would need to be developed. Meanwhile, even though we use a multivariate normal distributions within patterns, which can easily departures from MAR via differences in means and (co)-variances, we still need strong assumptions for sequential multivariate normal distribution within each pattern; otherwise MAR constraints do not exist (Wang and Daniels, 2011).

#### ACKNOWLEDGMENTS

This study is supported by NIH grant CA85295.

#### REFERENCES

- Bates, D., Mullen, K. M., Nash, J. C., and Varadhan, R. (2012). *minqa: Derivative-free optimization algorithms by quadratic approximation*. R package version 1.2.1.
- Bottai, M. and Zhen, H. (2013). Multiple imputation based on conditional quantile estimation. *Epidemiology, Biostatistics and Public Health* **10**,.
- Buchinsky, M. (1994). Changes in the US Wage Structure 1963-1987: Application of Quantile Regression. *Econometrica* **62**, pp. 405–458.
- Buchinsky, M. (1998). Recent advances in quantile regression models: A practical guideline for empirical research. *The Journal of Human Resources* **33**, pp. 88–126.
- Daniels, M. J. and Hogan, J. W. (2008). *Missing data in longitudinal studies*, volume 109 of

- Monographs on Statistics and Applied Probability*. Chapman & Hall/CRC, Boca Raton, FL. Strategies for Bayesian modeling and sensitivity analysis.
- Davison, A. C. and Hinkley, D. V. (1997). *Bootstrap methods and their application*, volume 1 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge. With 1 IBM-PC floppy disk (3.5 inch; HD).
- Efron, B. (1979). Bootstrap methods: another look at the jackknife. *Ann. Statist.* **7**, 1–26.
- Efron, B. and Tibshirani, R. J. (1993). *An introduction to the bootstrap*, volume 57 of *Monographs on Statistics and Applied Probability*. Chapman and Hall, New York.
- Hanson, T. and Johnson, W. O. (2002). Modeling regression error with a mixture of Polya trees. *J. Amer. Statist. Assoc.* **97**, 1020–1033.
- He, X., Ng, P., and Portnoy, S. (1998). Bivariate quantile smoothing splines. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **60**, 537–550.
- Heagerty, P. J. (1999). Marginally specified logistic-normal models for longitudinal binary data. *Biometrics* **55**, pp. 688–698.
- Hjort, N. L. and Petrone, S. (2007). Nonparametric quantile inference using Dirichlet processes. In *Advances in statistical modeling and inference*, volume 3 of *Ser. Biostat.*, pages 463–492. World Sci. Publ., Hackensack, NJ.
- Hjort, N. L. and Walker, S. G. (2009). Quantile pyramids for Bayesian nonparametrics. *Ann. Statist.* **37**, 105–131.
- Koenker, R. (2005). *Quantile regression*, volume 38 of *Econometric Society Monographs*. Cambridge University Press, Cambridge.
- Koenker, R. (2012). *quantreg: Quantile Regression*. R package version 4.91.
- Koenker, R. and Bassett, Gilbert, J. (1978). Regression quantiles. *Econometrica* **46**, pp. 33–50.
- Koenker, R. and Machado, J. A. F. (1999). Goodness of fit and related inference processes


- for quantile regression. *J. Amer. Statist. Assoc.* **94**, 1296–1310.
- Kottas, A. and Gelfand, A. E. (2001). Bayesian semiparametric median regression modeling. *Journal of the American Statistical Association* **96**, pp. 1458–1468.
- Kottas, A. and Krnjajić, M. (2009). Bayesian semiparametric modelling in quantile regression. *Scand. J. Stat.* **36**, 297–319.
- Little, R. J. A. (1993). Pattern-mixture models for multivariate incomplete data. *Journal of the American Statistical Association* **88**, 125–134.
- Little, R. J. A. (1994). A class of pattern-mixture models for normal incomplete data. *Biometrika* **81**, 471–483.
- Molenberghs, G., Michiels, B., Kenward, M. G., and Diggle, P. J. (1998). Monotone missing data and pattern-mixture models. *Statist. Neerlandica* **52**, 153–161.
- Perri, M. G., Limacher, M. C., Durning, P. E., Janicke, D. M., Lutes, L. D., Bobroff, L. B., Dale, M. S., Daniels, M. J., Radcliff, T. A., and Martin, A. D. (2008). Extended-care programs for weight management in rural communities: the treatment of obesity in underserved rural settings (tours) randomized trial. *Archives of internal medicine* **168**, 2347.
- R Core Team (2013). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0.
- Reich, B. J., Bondell, H. D., and Wang, H. J. (2010). Flexible Bayesian quantile regression for independent and clustered data. *Biostatistics* **11**, 337–352.
- Roy, J. (2003). Modeling longitudinal data with nonignorable dropouts using a latent dropout class model. *Biometrics* **59**, 829–836.
- Roy, J. and Daniels, M. J. (2008). A general class of pattern mixture models for nonignorable dropout with many possible dropout times. *Biometrics* **64**, 538–545, 668.
- Rubin, D. B. (1977). Formalizing subjective notions about the effect of nonrespondents in

- sample surveys. *J. Amer. Statist. Assoc.* **72**, 538–543.
- Wadden, T. A., Berkowitz, R. I., Sarwer, D. B., Prus-Wisniewski, R., and Steinberg, C. (2001). Benefits of lifestyle modification in the pharmacologic treatment of obesity: a randomized trial. *Archives of Internal Medicine* **161**, 218–227.
- Walker, S. and Mallick, B. K. (1999). A Bayesian semiparametric accelerated failure time model. *Biometrics* **55**, 477–483.
- Wang, C. and Daniels, M. J. (2011). A note on MAR, identifying restrictions, model comparison, and sensitivity analysis in pattern mixture models with and without covariates for incomplete data. *Biometrics* **67**, 810–818.
- Wei, Y., Ma, Y., and Carroll, R. J. (2012). Multiple imputation in quantile regression. *Biometrika* **99**, 423–438.
- Wei, Y., Pere, A., Koenker, R., and He, X. (2006). Quantile regression methods for reference growth charts. *Stat. Med.* **25**, 1369–1382.
- Yu, K., Lu, Z., and Stander, J. (2003). Quantile regression: applications and current research areas. *The Statistician* **52**, 331–350.
- Yu, K. and Moyeed, R. A. (2001). Bayesian quantile regression. *Statist. Probab. Lett.* **54**, 437–447.
- Yuan, Y. and Yin, G. (2010). Bayesian quantile regression for longitudinal studies with nonignorable missing data. *Biometrics* **66**, 105–114.

## APPENDIX

### *Goodness of Fit Check for Tours Data*

- Tours MAR:



● Tours MNAR:




Table 1

Scenario 1:  $MSE(MCSE)$  for coefficients estimates of quantiles 0.1, 0.3, 0.5, 0.7, 0.9 under MAR assumptions.  $(\gamma_{01}, \gamma_{11})$  are quantile regression coefficients for  $Y_{11}$ , and  $(\gamma_{02}, \gamma_{12})$  are ones for  $Y_{12}$ . MM stands for our proposed method, and RQ stands for the 'rq' function in R package 'quantreg'. BZ stands for Bottai's approach.

Normal	0.1			0.3			0.5			0.7			0.9		
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ
$\gamma_{01}$	0.06(0.01)	0.08(0.02)	0.08(0.02)	0.09(0.06)	0.09(0.03)	0.09(0.03)	0.23(0.04)	1.13(0.15)	1.13(0.15)	0.05(0.01)	0.07(0.02)	0.07(0.02)	0.05(0.01)	0.06(0.01)	0.06(0.01)
$\gamma_{11}$	0.04(0.01)	0.07(0.01)	0.07(0.01)	0.04(0.02)	0.07(0.02)	0.07(0.02)	0.95(0.04)	2.87(0.20)	2.87(0.20)	0.02(0.01)	0.06(0.01)	0.06(0.01)	0.04(0.01)	0.05(0.01)	0.05(0.01)
$\gamma_{02}$	0.08(0.01)	0.32(0.05)	0.09(0.02)	0.07(0.02)	0.59(0.05)	0.11(0.02)	0.09(0.02)	0.96(0.06)	0.14(0.03)	0.18(0.02)	1.47(0.08)	0.20(0.03)	0.45(0.05)	2.40(0.11)	0.26(0.04)
$\gamma_{12}$	0.05(0.01)	0.11(0.02)	0.08(0.01)	0.06(0.01)	0.08(0.01)	0.09(0.02)	0.07(0.01)	0.34(0.03)	0.20(0.04)	0.10(0.02)	1.00(0.06)	0.13(0.02)	0.11(0.02)	1.07(0.07)	0.12(0.02)
<hr/>															
$T_3$	0.1			0.3			0.5			0.7			0.9		
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ
$\gamma_{01}$	0.21(0.05)	0.12(0.03)	0.12(0.03)	0.14(0.03)	0.11(0.02)	0.11(0.02)	0.13(0.05)	1.35(0.14)	1.35(0.14)	0.12(0.04)	0.10(0.02)	0.10(0.02)	0.16(0.05)	0.12(0.03)	0.12(0.03)
$\gamma_{11}$	0.11(0.03)	0.10(0.02)	0.10(0.02)	0.09(0.02)	0.08(0.02)	0.08(0.02)	0.37(0.05)	1.96(0.20)	1.96(0.20)	0.07(0.02)	0.07(0.01)	0.07(0.01)	0.10(0.02)	0.12(0.02)	0.12(0.02)
$\gamma_{02}$	0.20(0.19)	0.48(0.10)	0.13(0.11)	0.18(0.13)	0.53(0.05)	0.10(0.03)	0.21(0.07)	1.03(0.05)	0.20(0.04)	0.25(0.07)	1.74(0.09)	0.25(0.05)	0.37(0.07)	2.36(0.18)	0.49(0.11)
$\gamma_{12}$	0.09(0.02)	0.19(0.04)	0.09(0.02)	0.09(0.02)	0.06(0.01)	0.06(0.01)	0.09(0.03)	0.30(0.03)	0.20(0.04)	0.16(0.04)	0.96(0.06)	0.15(0.03)	0.16(0.04)	1.14(0.11)	0.16(0.02)
<hr/>															
Laplace	0.1			0.3			0.5			0.7			0.9		
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ
$\gamma_{01}$	2.34(0.26)	1.80(0.22)	1.80(0.22)	0.19(0.03)	0.22(0.04)	0.22(0.04)	0.17(0.03)	0.69(0.09)	0.69(0.09)	0.23(0.05)	0.21(0.05)	0.21(0.05)	1.77(0.20)	1.23(0.20)	1.23(0.20)
$\gamma_{11}$	0.22(0.04)	0.46(0.06)	0.46(0.06)	0.20(0.04)	0.19(0.04)	0.19(0.04)	0.14(0.02)	0.93(0.11)	0.93(0.11)	0.16(0.03)	0.24(0.04)	0.24(0.04)	0.28(0.05)	0.46(0.09)	0.46(0.09)
$\gamma_{02}$	2.94(0.21)	4.67(0.55)	1.89(0.27)	0.49(0.06)	1.28(0.15)	0.26(0.05)	0.24(0.04)	1.07(0.08)	0.20(0.03)	0.59(0.09)	1.04(0.09)	0.39(0.07)	2.82(0.30)	1.10(0.18)	2.87(0.37)
$\gamma_{12}$	0.29(0.04)	1.08(0.15)	0.45(0.09)	0.25(0.04)	0.23(0.05)	0.16(0.04)	0.21(0.03)	0.37(0.06)	0.18(0.04)	0.30(0.04)	1.14(0.12)	0.26(0.06)	0.34(0.05)	1.54(0.21)	0.58(0.14)

Table 2

Scenario 2: MSE(MCSE) for coefficients estimates of quantiles 0.1, 0.3, 0.5, 0.7, 0.9 under MNAR scenario. In this scenario, we adopted MAR assumption for our approach and thus misspecified the MDM. ( $\gamma_{01}, \gamma_{11}$ ) are quantile regression coefficients for  $Y_{11}$ , and ( $\gamma_{02}, \gamma_{12}$ ) are ones for  $Y_{12}$ . MM stands for our proposed method, and RQ stands for the 'rq' function in R package 'quantreg'. BZ stands for Bottai's approach.

Normal	0.1				0.3				0.5				0.7				0.9			
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ			
$\gamma_{01}$	0.08(0.01)	0.10(0.01)	0.10(0.01)	0.08(0.02)	0.14(0.02)	0.14(0.02)	0.30(0.03)	1.35(0.15)	1.35(0.15)	1.35(0.15)	0.10(0.04)	0.13(0.04)	0.06(0.01)	0.08(0.01)	0.08(0.01)	0.08(0.01)	0.08(0.01)			
$\gamma_{11}$	0.05(0.01)	0.08(0.01)	0.08(0.01)	0.03(0.00)	0.11(0.02)	0.11(0.02)	1.00(0.04)	2.87(0.23)	2.87(0.23)	2.87(0.23)	0.04(0.01)	0.09(0.02)	0.04(0.00)	0.06(0.01)	0.06(0.01)	0.06(0.01)	0.06(0.01)			
$\gamma_{02}$	0.15(0.03)	0.42(0.05)	0.14(0.02)	0.10(0.01)	0.85(0.06)	0.15(0.02)	1.30(0.08)	4.05(0.12)	4.05(0.12)	4.05(0.12)	4.88(0.13)	9.95(0.19)	3.71(0.17)	6.89(0.20)	12.67(0.26)	4.43(0.24)	4.43(0.24)			
$\gamma_{12}$	0.08(0.01)	0.13(0.02)	0.09(0.01)	0.08(0.01)	0.08(0.01)	0.09(0.01)	0.05(0.01)	0.31(0.02)	0.31(0.02)	0.31(0.02)	0.12(0.02)	1.03(0.05)	0.15(0.02)	0.10(0.01)	1.06(0.07)	0.11(0.02)	0.11(0.02)			
$T_3$	0.1				0.3				0.5				0.7				0.9			
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ			
$\gamma_{01}$	0.21(0.05)	0.15(0.03)	0.15(0.03)	0.14(0.03)	0.11(0.02)	0.11(0.02)	0.14(0.03)	0.96(0.13)	0.96(0.13)	0.96(0.13)	0.17(0.03)	0.12(0.02)	0.12(0.02)	0.37(0.25)	0.16(0.03)	0.16(0.03)	0.16(0.03)			
$\gamma_{11}$	0.10(0.03)	0.11(0.03)	0.11(0.03)	0.07(0.01)	0.08(0.03)	0.08(0.03)	0.43(0.05)	1.92(0.21)	1.92(0.21)	1.92(0.21)	0.09(0.01)	0.09(0.02)	0.09(0.02)	0.10(0.02)	0.11(0.02)	0.11(0.02)	0.11(0.02)			
$\gamma_{02}$	0.31(0.08)	0.64(0.15)	0.17(0.04)	0.15(0.04)	0.74(0.06)	0.10(0.03)	1.18(0.09)	4.14(0.11)	4.14(0.11)	4.14(0.11)	3.97(0.18)	10.19(0.19)	3.59(0.20)	4.18(0.24)	11.28(0.43)	3.57(0.30)	3.57(0.30)			
$\gamma_{12}$	0.10(0.04)	0.23(0.05)	0.12(0.02)	0.12(0.02)	0.05(0.01)	0.06(0.02)	0.09(0.02)	0.26(0.02)	0.26(0.02)	0.26(0.02)	0.19(0.05)	1.01(0.05)	0.17(0.03)	0.21(0.06)	1.26(0.12)	0.19(0.04)	0.19(0.04)			
Laplace	0.1				0.3				0.5				0.7				0.9			
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ			
$\gamma_{01}$	2.69(0.24)	2.22(0.23)	2.22(0.23)	0.34(0.05)	0.37(0.06)	0.37(0.06)	0.20(0.03)	0.96(0.12)	0.96(0.12)	0.96(0.12)	0.23(0.03)	0.30(0.05)	0.30(0.05)	2.62(0.23)	2.03(0.23)	2.03(0.23)	2.03(0.23)			
$\gamma_{11}$	0.39(0.05)	0.55(0.08)	0.55(0.08)	0.33(0.04)	0.37(0.07)	0.37(0.07)	0.16(0.02)	1.15(0.14)	1.15(0.14)	1.15(0.14)	0.25(0.03)	0.27(0.06)	0.27(0.06)	0.23(0.03)	0.52(0.07)	0.52(0.07)	0.52(0.07)			
$\gamma_{02}$	4.47(0.37)	7.17(0.77)	3.45(0.45)	1.04(0.12)	1.78(0.16)	0.58(0.08)	1.57(0.15)	4.15(0.20)	4.15(0.20)	4.15(0.20)	2.49(0.20)	7.98(0.31)	2.80(0.21)	0.90(0.10)	5.05(0.45)	1.32(0.18)	1.32(0.18)			
$\gamma_{12}$	0.53(0.08)	1.07(0.20)	0.72(0.13)	0.38(0.05)	0.27(0.04)	0.27(0.04)	0.20(0.03)	0.48(0.06)	0.48(0.06)	0.48(0.06)	0.34(0.04)	1.22(0.10)	0.36(0.04)	0.30(0.03)	1.71(0.24)	0.55(0.08)	0.55(0.08)			



Table 3

Scenario 3:  $MSE(MCSE)$  for coefficients estimates of quantiles 0.1, 0.3, 0.5, 0.7, 0.9 under MNAR scenario. In this scenario, we used the correct sensitivity parameters for our approach.  $(\gamma_{01}, \gamma_{11})$  are quantile regression coefficients for  $Y_{i1}$ , and  $(\gamma_{02}, \gamma_{12})$  are ones for  $Y_{i2}$ . MM stands for our proposed method, and RQ stands for the 'rq' function in R package 'quantreg'. BZ stands for Bottai's approach.

Normal	0.1			0.3			0.5			0.7			0.9		
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ
$\gamma_{01}$	0.13(0.03)	0.11(0.01)	0.11(0.01)	0.13(0.02)	0.16(0.02)	0.16(0.02)	0.37(0.04)	1.15(0.12)	1.15(0.12)	0.08(0.01)	0.13(0.02)	0.13(0.02)	0.06(0.01)	0.07(0.01)	0.07(0.01)
$\gamma_{11}$	0.07(0.01)	0.08(0.01)	0.08(0.01)	0.05(0.01)	0.13(0.02)	0.13(0.02)	0.94(0.05)	2.48(0.20)	2.48(0.20)	0.04(0.01)	0.09(0.02)	0.09(0.02)	0.04(0.01)	0.05(0.01)	0.05(0.01)
$\gamma_{02}$	0.13(0.02)	0.48(0.05)	0.14(0.02)	0.15(0.03)	0.92(0.05)	0.15(0.02)	0.37(0.05)	4.25(0.11)	1.31(0.09)	0.64(0.06)	10.20(0.17)	3.70(0.16)	0.97(0.08)	12.95(0.26)	4.27(0.22)
$\gamma_{12}$	0.07(0.01)	0.09(0.02)	0.09(0.01)	0.08(0.01)	0.07(0.01)	0.10(0.02)	0.14(0.02)	0.28(0.03)	0.23(0.05)	0.10(0.02)	0.97(0.05)	0.14(0.02)	0.11(0.02)	1.04(0.07)	0.12(0.02)
$T_3$	0.1			0.3			0.5			0.7			0.9		
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ
$\gamma_{01}$	0.36(0.08)	0.19(0.03)	0.19(0.03)	0.28(0.04)	0.14(0.02)	0.14(0.02)	0.17(0.03)	1.11(0.12)	1.11(0.12)	0.19(0.03)	0.15(0.02)	0.15(0.02)	0.25(0.05)	0.17(0.03)	0.17(0.03)
$\gamma_{11}$	0.16(0.03)	0.14(0.03)	0.14(0.03)	0.16(0.04)	0.11(0.02)	0.11(0.02)	0.56(0.05)	2.00(0.20)	2.00(0.20)	0.10(0.01)	0.11(0.02)	0.11(0.02)	0.11(0.02)	0.12(0.01)	0.12(0.01)
$\gamma_{02}$	0.36(0.08)	0.83(0.12)	0.29(0.05)	0.26(0.04)	0.73(0.05)	0.15(0.02)	0.33(0.05)	3.88(0.11)	1.24(0.09)	0.77(0.08)	9.80(0.21)	3.72(0.20)	0.67(0.08)	10.92(0.40)	3.76(0.26)
$\gamma_{12}$	0.16(0.03)	0.27(0.04)	0.16(0.03)	0.11(0.02)	0.08(0.01)	0.10(0.02)	0.19(0.03)	0.35(0.03)	0.27(0.04)	0.30(0.06)	1.19(0.07)	0.22(0.03)	0.32(0.06)	1.52(0.12)	0.24(0.03)
Laplace	0.1			0.3			0.5			0.7			0.9		
	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ	MM	RQ	BZ
$\gamma_{01}$	2.16(0.20)	1.82(0.21)	1.82(0.21)	0.29(0.04)	0.43(0.07)	0.43(0.07)	0.22(0.03)	0.94(0.12)	0.94(0.12)	0.28(0.04)	0.33(0.04)	0.33(0.04)	2.49(0.24)	2.06(0.26)	2.06(0.26)
$\gamma_{11}$	0.37(0.05)	0.53(0.07)	0.53(0.07)	0.21(0.03)	0.29(0.04)	0.29(0.04)	0.15(0.02)	1.10(0.14)	1.10(0.14)	0.28(0.04)	0.43(0.07)	0.43(0.07)	0.35(0.06)	0.59(0.08)	0.59(0.08)
$\gamma_{02}$	2.42(0.26)	5.97(0.57)	3.34(0.38)	0.38(0.05)	1.48(0.14)	0.55(0.07)	0.44(0.05)	3.89(0.18)	1.50(0.13)	0.56(0.08)	7.58(0.35)	2.81(0.24)	1.48(0.18)	4.43(0.39)	1.37(0.21)
$\gamma_{12}$	0.38(0.06)	0.96(0.12)	0.76(0.10)	0.31(0.05)	0.30(0.05)	0.27(0.04)	0.25(0.04)	0.48(0.05)	0.34(0.06)	0.30(0.05)	1.35(0.14)	0.39(0.07)	0.56(0.11)	1.89(0.27)	0.56(0.08)

**Table 4**

*Estimated marginal quantile regression coefficients with 95% bootstrap percentile confidence interval for weight of participants at 6 and 18 months. Missing data mechanism assumption is MAR.*

		Intercept	Age	White	BaseWeight
6 months					
10%	-6.05 (-10.88, 2.67)	0.34 (-0.25, 0.85)	-3.86 (-5.75, -2.43)	0.92 (0.85, 0.97)	
30%	-2.56 (-7.67, 3.66)	0.33 (-0.25, 0.84)	-3.90 (-5.43, -2.54)	0.92 (0.86, 0.97)	
50%	-0.25 (-5.29, 5.60)	0.31 (-0.25, 0.85)	-4.04 (-5.57, -2.55)	0.93 (0.87, 0.98)	
70%	1.79 (-3.27, 7.81)	0.35 (-0.22, 0.86)	-4.11 (-5.67, -2.68)	0.93 (0.87, 0.98)	
90%	4.81 (-0.05, 11.32)	0.40 (-0.20, 0.94)	-4.07 (-5.68, -2.68)	0.94 (0.88, 0.99)	
18 months(MAR)					
10%	-17.65 (-31.75, 21.41)	-0.73 (-1.99, 0.39)	-0.12 (-10.60, 2.96)	1.01 (0.63, 1.14)	
30%	-18.26 (-28.27, 9.88)	-0.74 (-2.01, 0.32)	1.07 (-8.93, 3.67)	1.07 (0.79, 1.17)	
50%	-12.72 (-24.20, 10.45)	-0.73 (-2.01, 0.30)	1.04 (-6.14, 3.94)	1.06 (0.83, 1.17)	
70%	-9.12 (-19.69, 14.38)	-0.73 (-2.00, 0.31)	1.18 (-5.18, 3.92)	1.06 (0.84, 1.16)	
90%	-3.90 (-12.65, 19.61)	-0.75 (-1.98, 0.36)	1.24 (-4.19, 3.76)	1.08 (0.85, 1.16)	
18 months(MNAR)					
10%	-20.51 (-30.97, 25.00)	-0.69 (-2.23, 0.47)	0.24 (-10.19, 3.04)	1.04 (0.62, 1.14)	
30%	-18.04 (-27.14, 8.73)	-0.74 (-2.04, 0.49)	1.08 (-9.22, 3.94)	1.07 (0.83, 1.16)	
50%	-12.19 (-22.52, 8.79)	-0.73 (-2.06, 0.38)	1.05 (-6.36, 4.17)	1.06 (0.86, 1.16)	
70%	-7.89 (-17.63, 12.26)	-0.73 (-1.95, 0.32)	1.17 (-4.43, 4.20)	1.06 (0.87, 1.16)	
90%	-3.11 (-8.60, 21.70)	-0.73 (-2.02, 0.38)	1.68 (-3.90, 4.05)	1.10 (0.87, 1.15)	