InsDel2a / 2b / 2c per−sample share by Major Cancer Type and stratum

Each sample appears in exactly one stratum. Hypermutator cutoff = 4000 total indels. Black bar = median.

MSS non−hypermutated
MSI−H
MSS hypermutated

Signature share of sample's assigned indels


C_ID2/InsDel2a
C_ID2/InsDel2b
C_ID2/InsDel2c

1.00
0.75
0.50
0.25
0.00

Biliary
Bladder
Bone.SoftTissue
Breast
CNS
Colon
Esophagus
Head
Kidney
Liver
Lung
Lymphoid
Myeloid
Other
Ovary
Pancreas
Prostate
Skin
Stomach
Thymus
Uterus


C_ID2/InsDel2a mutations attributed per sample

Each dot = a sample. Red dashed line = per−type median over nonzero samples. Counts beneath bars: nonzero / total samples in that stratum.



C_ID2/InsDel2b mutations attributed per sample

Each dot = a sample. Red dashed line = per−type median over nonzero samples. Counts beneath bars: nonzero / total samples in that stratum.



C_ID2/InsDel2c mutations attributed per sample

Each dot = a sample. Red dashed line = per−type median over nonzero samples. Counts beneath bars: nonzero / total samples in that stratum.

