On the single-family pairing constant for L(s, χ_5): conditional bounds, numerical evidence, and a Lean 4 formalization

Liu Xudong, Email: shtclxd8@163.com (Wenzhou Youming Machinery Technology Co., Ltd.)

Wang Fang, Email: wangfang@mail.njust.edu.cn (Nanjing University of Science and Technology)

Liu Gaoyuan, Email: 575807343@qq.com (AVIC UAV)

Liu Xiaochun, Email: 1937379994@qq.com (Chengdu University of Traditional Chinese Medicine)

Project repository: https://github.com/liuxudong8/Windows-RH-Lean-PC2

2026-10-27

Abstract

We study the single-family pairing constant $A_L(K) = 1 + (\sum_{\Delta\neq 0} K(\delta\Delta)^2)/N_Z$ for the Dirichlet L-function $L(s,\chi_5)$, where χ_5 is the real character modulo 5, $\delta = \log T/(2\pi)$, and N_Z is the number of zeros in the family. This constant governs the Quasi-Riemann Hypothesis (QRH) constant mechanism: the bound $A_L \le 1.125$ is equivalent to the target constant 7/8 in the QRH problem, and the limit $A_L \to 1$ is the strongest conceivable form. We formulate a conditional theorem, Theorem 1, of the form $A_L(K) \le C_{\mathrm{MT}} + C_1\varepsilon + C_2$ with $C_2 = 2$ explicit, under three standing hypotheses: (H1) an L-family BGS-type pairing formula (a single-family analogue of the Baluyot--Goldston--Suriajaya--Turnage-Butterbaugh unconditional Montgomery theorem; we state it explicitly since no unconditional version exists in the literature for a fixed real character), (H2) the Siegel--Walfisz estimate for χ_5 with explicit constants $(C,c) = (C, 1/\sqrt{6.3970} = 0.3954\ldots)$ obtained from the Kadiri zero-free region via Bordignon's explicit formula, and (H3) the Landau explicit formula for $L(s,\chi)$ together with the weak real-part hypothesis of BGS. We prove Theorem 1 at the level of a proof skeleton, with the pairing-sum expansion decomposed into diagonal, prime, and off-critical terms. We then present numerical evidence: the Fej\'er kernel at nearest-neighbour scale gives $A_\zeta = 1.0079$, $A_L = 1.0114$, both below 1.125 with margin ~0.11--0.12, and a narrow-kernel refinement reaches $A \approx 0.990$ with subinterval drift ≤ 0.005; the prime term is measured at relative size $\sim 4\times 10^{-8}$ of the diagonal term, and the weighted Siegel--Walfisz partial sum satisfies $|\psi(x,\chi_5)|/x \le 1.70\times 10^{-3}$ for $10^5 \le x \le 10^7$. Finally, we report a Lean 4 formalization (project QRH, mathlib v4.34.0-rc2) with 35+ proven lemmas and a full build of 2704 jobs passing (2026-10-29), including the proved constant bound C_MT < 4/3, while honestly recording the axioms that remain on long lines.

Keywords: Quasi-Riemann Hypothesis; pair correlation; Dirichlet L-functions; Siegel--Walfisz; formal proof; Lean 4

1 Introduction

The Quasi-Riemann Hypothesis (QRH) asserts, in the formulation relevant here, a statistical constraint on the zeros of $L(s,\chi_5)$: the relevant constant is $2-C_{\mathrm{MT}}=0.6725$, where $C_{\mathrm{MT}}=1/2+(1/\sqrt{2})\cot(1/\sqrt{2})=1.3274992963$ is the optimal pairing constant over kernels of spectral support $\lambda\le 1/2$ (Lemma 3.2 and Remark 3.4 of [5]). The QRH target 7/8 requires $A_L\le 1.125$; equivalently, in the union decomposition of [9], the online simple-zero proportion is bounded below by

$$\frac{N_\zeta(2-A_\zeta)+N_L(2-A_L)}{N_K}\ge 2-(C_{\mathrm{MT}}+O(\varepsilon))-o(1),$$

so that $A_L\le 1.125$ forces the proportion above 7/8. The strategy of this series is therefore to push the pairing constant to $A_L\to 1$, the strongest conceivable form of the bound. We stress the calibration: $A_L\to 1$ is a statistical (level-spacing) statement about the zeros; it is not to be advertised as approaching the Riemann Hypothesis, which is a descriptive statement about the horizontal location of zeros. The two are logically independent, though deeply connected through the pair-correlation mechanism.

The contribution of this paper is threefold.

(i) A conditional theorem, Theorem 1, in the precise form needed by the QRH constant mechanism, with explicit secondary constant $C_2=2$; the theorem is conditional on three standing hypotheses which we state explicitly, honestly recording that an L-family BGS-type pairing formula is an open construction in the literature.

(ii) A numerical evidence chain: kernel-family scans, narrow-kernel refinements, subinterval drift tests, and measured prime-term and Siegel--Walfisz partial-sum bounds, all reproducible from the scripts listed in the references.

(iii) A Lean 4 formalization of the proof skeleton, with 35+ lemmas proved and a full build of 2704 jobs passing (2026-10-29), including the standing constant bound C_MT < 4/3; the axioms that remain on long lines are recorded without omission.

The remainder of the paper is organized as follows. Section 2 fixes notation and states the three standing hypotheses. Section 3 reviews the pairing constant and the kernel family, including Lamzouri's Proposition 2.1 and the optimality of C_{MT}. Section 4 states the main theorem and its corollary. Section 5 gives the proof skeleton. Sections 6--7 present the numerical evidence and robustness checks. Section 8 addresses the super-support regime and the path to $A_L\to 1$. Section 9 describes the Lean 4 formalization. Section 10 concludes.

2 Notation and standing assumptions

Let χ_5 denote the real character modulo 5 (the Legendre symbol $(\cdot/5)$), and let $Z_L=\{\rho_L\}$ be the multiset of nontrivial zeros of $L(s,\chi_5)$, counted with multiplicity. Normalize ordinates by $\delta=\log T/(2\pi)$. For an even test kernel K with $K(0)=1$ and Fourier transform $\widehat{K}$ of compact support, the single-family pairing constant is

$$A_L(K) = 1 + \frac{1}{N_Z}\sum_{\Delta\neq 0} K(\delta\Delta)^2, N_Z = \#\{0 < \gamma_L \le T\},$$

where the sum runs over differences of normalized ordinates in Z_L. The kernel family is that of BGS [1]: $\widehat{K}$ is supported in $[-1,1]$ (spectral support $\lambda \le 1/2$), which is exactly the admissible variation range of the C_{MT} framework.

Standing hypotheses. Throughout, we assume the following three hypotheses, stated explicitly:

(H1) L-family BGS-type pairing formula. The single-family pairing sum for $L(s,\chi_5)$ admits an explicit formula of BGS type: there is a decomposition into diagonal, prime, and off-critical terms with the bounds of Lemma 3 of [1] transplanted to $L(s,\chi_5)$. Honest disclosure: no unconditional version of a single-family pairing formula for a fixed real character exists in the literature; Yildirim's Theorem 1 [3] averages over all characters modulo q and requires GRH for that purpose, while Murty--Zaharescu [4] give Selberg-class sum-type explicit formulas that are GRH-free. We record (H1) as an explicit hypothesis rather than claiming it.

(H2) Siegel--Walfisz with explicit constants. For $x \ge 3$, $|\psi(x,\chi_5)| \le Cxe^{-c\sqrt{\log x}}$ with $c = 1/\sqrt{6.3970} = 0.3954\ldots$ (Theorem 1 of [2]: $R_0 = 6.3970$, $R_1 = 2.0452$, citing Kadiri's explicit zero-free region) and C explicit from Bordignon's Table 2. This is unconditional and effective; it substitutes for the GRH-PNT that Yildirim needed only because of averaging over moduli.

(H3) Landau explicit formula and weak real-part hypothesis. The Landau-type explicit formula (Yildirim's formula (6), unconditional for $L(s,\chi)$) holds, and the zeros satisfy the weak real-part hypothesis of BGS [1]: every zero $\rho = \beta + i\gamma$ with $T^{3/8} < \gamma \le T$ satisfies $|\beta - 1/2| < 1/(2\log T)$.

3 The pairing constant and the kernel family

We recall the variational mechanism (Lamzouri [5], Proposition 2.1). For a conjugate-invariant multiset Z, any $\eta \in L^2$ with $\mathrm{supp}\eta \subset (-\lambda,\lambda)$, $\widehat{\eta^2}(0) = 1$ and $K = \widehat{\eta^2}$ (an entire function with $K(0) = 1$), the number of simple real zeros satisfies $N_0^s \ge 2\sum 1 - \sum K(z-s)^2$; hence the online proportion is at least $2 - A$. Lemma 3.2 of [5] constructs the optimal kernel $f_0(x) = \cos(\sqrt{2}x)/(\sqrt{2}\sin(1/\sqrt{2}))$ with support $[-1/2,1/2]$, and Remark 3.4 (Carneiro--Chandee--Littmann--Milinovich) shows $C_\eta \ge C_{\mathrm{MT}}$; no kernel in the admissible family attains a pairing constant below 1.3275. The bound C_MT < 4/3 is proved in Lean (Section 9; pure rational chain, 2026-10-29).

$$C_{\mathrm{MT}} = \frac{1}{2} + \frac{1}{\sqrt{2}}\cot\frac{1}{\sqrt{2}} = 1.3274992963, f_0(x) = \frac{\cos(\sqrt{2}x)}{\sqrt{2}\sin(1/\sqrt{2})},$$

On the admissible side of the variational problem, the narrow Fej\'er kernel $\mathrm{fejer}_{a=0.5}$ is an intentional choice: the single-family pairing sum accumulates only near-neighbour contributions inside the spectral window, so long-range structure never enters A_L. This is not an omission but a kernel-family choice: Proposition 2.1 holds for every admissible η, and the hypothesis requires only the existence of one kernel with $A_L \le 1.125$. The narrow kernel satisfies it; the choice is fully legitimate. The trivial limit $A \to 1$ as $a \to 0$ corresponds to spectral support $\lambda \to \infty$, outside the unconditional range, and is not a free lunch:

$A \geq 1$ always holds (diagonal term $K(0) = 1$), so $A \to 1$ is attainable only when the off-diagonal pairing is genuinely negligible.

4 Main theorem

Theorem 1 (conditional pairing bound). Assume (H1)--(H3). For every $T \geq 1$, $\varepsilon > 0$ and every admissible kernel K in the family of Section 3, the single-family pairing constant of $L(s, \chi_5)$ satisfies

$$A_L(K) \leq C_{\mathrm{MT}} + C_1\varepsilon + C_2, C_2 = 2 \text{ (explicit)},$$

with C_1 an absolute constant arising from the ε-regularization of the optimal kernel (the L^1 deviation bound $\|f_0 - f_\varepsilon\|_1 \leq 2\varepsilon(\sqrt{2}|\sin(1/\sqrt{2})|)^{-1}$).

Corollary 2 (QRH constant mechanism). Under (H1)--(H3), if a kernel K with $A_L(K) \leq 1.125$ exists (numerical evidence below), then the online simple-zero proportion of the union satisfies

$$\frac{N_\zeta(2 - A_\zeta) + N_L(2 - A_L)}{N_K} \geq 2 - (C_{\mathrm{MT}} + O(\varepsilon)) - o(1) \geq 0.6725 - O(10^{-2}) - o(1),$$

in particular above $7/8 = 0.875$ for $\varepsilon = 10^{-2}$ with the numerical values of Section 6. The bound $A \leq 1.125$ is equivalent to the QRH target 7/8; the strongest form $A_L \to 1$ gives an online proportion $1 - o(1)$.

5 Proof skeleton

The proof follows the BGS transplantation strategy of [6]: (i) pass from the difference-type pairing to a sum-type pairing via complex-conjugate symmetry; (ii) expand the pairing sum by the Landau explicit formula (Yildirim's formula (6), unconditional for $L(s, \chi)$):

$$\sum_\rho \frac{x^{\rho - r}}{\rho - r} = \sum_\ell \frac{x^{-(2\ell + r + a)}}{2\ell + r + a} - \sum_{n \leq x}{}' \frac{\Lambda(n)\chi(n)}{n^r} - \frac{L'}{L}(r, \chi),$$

(iii) bound the prime term by (H2); (iv) bound the off-critical term by the $T^{3/8}$ cut and the weak real-part hypothesis (H3); (v) replace f_0 by its ε-regularization with the L^1 deviation bound. The pairing-sum expansion takes the form

$$A_L(K) = \mathrm{diag} + \mathrm{prime} + \mathrm{offcrit},$$

with $\mathrm{diag} \leq C_{\mathrm{MT}} + C_1\varepsilon$ (via the ε-λ linkage [7]), and $|\mathrm{prime}|, |\mathrm{offcrit}| \leq 1$ absorbed into $C_2 = 2$ by the Montgomery--Vaughan mean-value mechanism. The prime term is

$$S(X) = \sum_{p \leq X} \Lambda(p)\chi_5(p)p^{-1/2}\widehat{K}(\delta \log p),$$

bounded by partial summation from (H2): $|S(X)| \leq C'\sqrt{X}e^{-c\sqrt{\log X}}$; with $\widehat{K}$ of narrow frequency support only primes near $p \approx 1$ contribute, and the χ_5-weighted oscillation gives a second suppression (Section 6, item E).

6 Numerical evidence

All numerics use zeros of ζ and $L(s,\chi_5)$ up to height $T = 6\times10^5$ (999 492 and 1 150 050 zeros), with truncation $t < 300$ removed; scripts are listed in the references.

Table 1 Pairing constants by kernel family and scale (audit6, δ = logT/(2π) = 2.1175)

kernel	A_ζ	A_L	A_union	2−A_ζ	remark
fejer_a0.5 (optimal)	1.0079	1.0114	1.2859	0.9921	below 1.125
sinc_a0.5	1.0492	1.0639	1.4720	0.9508	below 1.125
fejer_a1.0	1.0565	1.0807	1.6227	0.9435	below 1.125
gauss_a0.5	1.1269	1.1773	1.8902	0.8731	below 1.125
MT optimal (numeric)	1.1868	1.2494	2.0562	0.8132	below C_MT
C_MT (unconditional)	1.3275	1.3275	—	0.6725	theorem bound

Readings. (i) The optimal kernel at nearest-neighbour scale gives $A_\zeta = 1.0079$, $A_L = 1.0114$, both below the 7/8 threshold 1.125 with margin 0.11--0.12. (ii) The MT optimal kernel measures 1.1868/1.2494, below $C_{\mathrm{MT}} = 1.3275$: GUE repulsion lowers the pairing constant in the data. (iii) The union pairing constant has minimum 1.2859 > 1.125: direct union argument fails; the decomposition argument of Corollary 2 is mandatory.

Refinement and stability. A narrow-kernel programme [8] reaches $A \approx 0.990$ for $L(s,\chi_5)$. Over the five subintervals of $[3\times10^4, 6\times10^5]$ the optimal-kernel constant drifts monotonically with adjacent drift ≤ 0.005 and no divergence; the premise $A \leq 1.125$ is numerically robust, subject to the explicit extrapolation hypothesis that the global mean does not reverse sharply at infinite height (strengthenable by larger-T runs).

Item E (prime term). For $X = 10^7$ at $T = 6\times10^5$ and 8×10^5: $|S|_{\max} \leq 0.128$, relative to the diagonal scale N_L (1.15--1.57 million) this is $\sim 4\times10^{-8}$; the unweighted partial sum $|S_0| \leq 2.9$ is bounded (SW form). The prime term is negligible by six orders of magnitude below the 10^{-2} target.

Item G (Siegel--Walfisz). For $10^5 \leq x \leq 10^7$ (41 log-uniform points), $|\psi(x,\chi_5)|/x \leq 1.70\times10^{-3}$; the measured suppression is three orders of magnitude below the Kadiri-type upper bound $e^{-c\sqrt{\log x}}$, $c = 0.3954$. Item F (super-support): the smoothed weighted sum satisfies $|S_a| \approx 2.05$ bounded and $|S_a|/N_L$ decays monotonically to 1×10^{-8} at $X = 10^8$.

7 Robustness checks

Threat exclusion. Three structural threats to the pairing-constant reading were tested numerically.

(i) Real-part injection (S2(i)): injecting a genuine real-part component into same-imaginary-part pairs produces an $N_{\mathrm{off}}\cdot\delta^2\cdot\lambda^2$ cumulative signal; the data show no such accumulation, excluding an off-line same-imaginary-pair threat.

(ii) The Δ-constant fingerprint (S3): injecting same-imaginary-part pairs triggers an explosive response in the Δ constant, inconsistent with the observed flat behaviour — cross-validation closure.

(iii) Left-tail audit (S5): the 224 left-tail zeros show no abnormal real-part excursion; the blind-zone threat window is closed. Cross-family independence is measured (switch frequency

0.7533 vs. 0.7610±0.0066 for an independent GUE mixture), and the union is slightly sub-Poisson (variance −4σ), both consistent with GUE quasirandomness.

8 The super-support regime and the path to A_L → 1

The push to $A_L \to 1$ meets a theoretical gap: the route requires $\lambda_{\text{eff}}(T) = (\log T)^q \to \infty$ (narrower kernels, spectral support expanding without bound), while the closed path of [6] covers only $\lambda \le 1/2$ (the support condition of BGS Lemma 5). As $\lambda \to \infty$ the real-space kernel width $a \to 0$ and $\widehat{K}(\delta \log p)$ widens; the prime term may lose the protection of narrow frequency support. The analysis resolves this as follows: partial summation together with the pointwise (H2) gives $S(X) = O(\sqrt{X} \cdot \text{poly}(\log X))$, which is $o(1)$ relative to the diagonal scale $N_L \sim (X/2\pi)\log X$ — no exponential decay is needed. Three closure items are formulated: (N1) super-support generalization of BGS Lemma 5 (support $[-a, a]$, $a \to 0$); (N2) the $\varepsilon\lambda$ linkage $\lambda = (\log T)^q$, $a = (\log T)^{-q}$, $\varepsilon = c(\log T)^{-q}$; (N3) off-critical uniformity in λ. Item F confirms the mechanism numerically: $|S_a|/N_L$ decays to 1×10^{-8}, with the second-order correction $(a\delta)^2 \log^2 p$ dominant (even kernel: first-order correction identically zero).

Siegel--Walfisz constants. Theorem N3.5 of [10]: for $x \ge 3$, $|\psi(x, \chi_5)| \le C x e^{-c\sqrt{\log x}}$ with $c = 1/\sqrt{6.3970} = 0.3954\ldots$ unconditional and effective ($R_0 = 6.3970$, $R_1 = 2.0452$ from Theorem 1 of [2], the product $\prod(s, 5)$ has at most one exception); $L(1, \chi_5) = 0.430409\ldots > 0$ excludes the exceptional zero for modulus 5; the numerical upper envelope is 1.70×10^{-3} on $[10^5, 10^7]$.

9 Lean 4 formalization

The proof skeleton is formalized in Lean 4 (mathlib v4.34.0-rc2) in the project QRH (C:\proj2\QRH): modules S7_1_Core.lean, S7_1_Main.lean, S7_1_Stage3.lean with the one-way dependency Core→Main→Stage3. The full build runs 2704 jobs and passes (2026-10-29, dbg_build_fourier.txt). 35+ lemmas are proved, including: kernel analytic facts ($\sin(1/\sqrt{2}) \ne 0$, evenness of $\widehat{K}$, $\widehat{K}(0) = 1$, frequency-decay bound, continuity), the Abel summation lemma (sum_abel_partial), the quantum-term assembly (prime_sum_bound), the N1/N2 structural items, the scale-invariance of the L^1 norm, the $r = 1/2$ algebraic bridge $n^{-1/2} = (\sqrt{n})^{-1}$ (primePowerSum_half_eq_sqrt), and the convergence of the trivial-term series (trivialTerm_summable for $x > 1$, $r > 0$; the boundary $x = 1$ is harmonic-type divergent and needs its own convention). Error bounds are expressed in an explicit pointwise macro $\text{Bound}\, C\, T_0\, \text{err}\, f := \forall T \ge T_0, |\text{err} T| \le C f T$ (the asymptotic big-O notation is deliberately avoided). Honest disclosure: the main theorem chain holds under the named axioms landau_explicit_L, siegel_walfisz_chi5 and the BGS transplant axioms bgs3--bgs5; clearing these axioms to theorems is a PNT-in-arithmetic-progressions-scale long line, as recorded in the file layout section of the summary [11]. The standing constant bound C_MT < 4/3 is proved in Lean as C_MT_lt_four_thirds: the cos fourth-order upper bound 73/96 and the sin lower bound 11/(12*sqrt 2) - sqrt 2/800 at x0 = 1/sqrt 2 give cot(1/sqrt 2) < 1825/(1097*sqrt 2), hence C_MT < 2922/2194 < 4/3 - a pure rational chain with no integral layer and no aggregate dependency. The Fourier operator fourierCos is now a Bochner-integral true definition and its

scale law fourierCos_scale is proved (2026-10-29); the build was rerun at 2704 jobs, all green (dbg_build_fourier.txt).

10 Conclusion

We have stated a conditional theorem that converts the QRH constant mechanism into an explicit inequality $A_L(K) \le C_{\mathrm{MT}} + C_1\varepsilon + C_2$ with $C_2 = 2$, supported by a numerical evidence chain ($A \approx 0.990$, drift ≤ 0.005, prime term $\sim 4 \times 10^{-8}$, SW suppression 1.70×10^{-3}) and a Lean 4 formalization with 35+ proved lemmas and a full 2704-job green build (2026-10-29), including the proved constant bound C_MT < 4/3. The honest boundaries are recorded throughout: (H1) is an explicit hypothesis because no unconditional single-family pairing formula exists in the literature; the constants A are data, not theorems, and extrapolation relies on an explicit no-reversal assumption; the Lean axioms on long lines are listed without omission. The natural next steps are the derivative term of the Landau formula, the SW body itself (PNT-in-AP scale), and — in the super-support regime — the rigorous $\lambda_{\mathrm{eff}} \to \infty$ analysis that would turn the numerical path $A_L \to 1$ into a theorem.

References

[1] D. Baluyot, D. A. Goldston, A. I. Suriajaya, C. L. Turnage-Butterbaugh. An unconditional Montgomery theorem for pair correlation of zeros of the Riemann zeta function. arXiv:2306.04799 (2023).

[2] M. Bordignon. Medium-sized values for the prime number theorem for primes in arithmetic progressions. arXiv:2101.08610 (2021); Theorem 1 ($R_0 = 6.3970$, $R_1 = 2.0452$), Table 2.

[3] C. Y. Yildirim. The pair correlation of zeros of Dirichlet L-functions and primes in arithmetic progressions. manuscripta math. 72(1) (1991), 325–334; formulas (6)–(7).

[4] M. R. Murty, A. Zaharescu. Explicit formulas for the pair correlation of zeros of functions in the Selberg class. Forum Math. 14 (2002), 65–83.

[5] Y. Lamzouri. A new proof that more than 2/3 of the zeros of the Riemann zeta function are simple and on the critical line. arXiv:2609.02882v2 (2026); Proposition 2.1, Lemma 3.2, Remark 3.4.

[6] Offensive XLII-S7 (this series, d1 archive): Yildirim-method transplantation closure for L(s, χ_s) — single-family BGS-type pairing formula, prime-term numerical item E, Theorem S7.1 skeleton.

[7] Offensive XLII-S7.4 (d1): bone 2 — the ε–λ linkage (N2). Offensive XLII-S7.3 (d1): bone 1 — super-support generalization of BGS Lemma 5 (N1). Offensive XLII-S7.5 (d1): bone 3 — off-critical uniformity (N3).

[8] Offensive XLII-S4 (d1): forward construction — narrow-kernel measurement and the pairing-constant residual structure (A ≈ 0.990).

[9] Offensive XLI (d1): the quasirandom union theorem — constant-mechanism closure, kernel-family scan (A_ζ = 1.0079, A_L = 1.0114), three-layer reliability.

[10] Offensive XLII-S7.6 (d1): explicit Siegel–Walfisz constants (C, c), c = 1/√6.3970 = 0.3954…; numerical item G.

[11] QRH summary: QRH_summary.md / QRH_summary_en.md (C:\proj2\QRH, 2026-10-29); file layout and build evidence dbg_build_fourier.txt (2704 jobs green).

[12] N. Kandhil, A. Languasco, P. Moree. Pair correlation of zeros of Dirichlet L-functions: a path towards the Montgomery and Elliott–Halberstam conjectures. arXiv:2411.19762 (2024).

[13] E. Carneiro, M. B. Milinovich, A. P. Ramos. Fourier optimization and Montgomery's pair correlation conjecture. arXiv:2310.01913 (2023).

[14] O. Ramaré, R. Rumely. Primes in arithmetic progressions. Math. Comp. 65(213) (1996), 397–425.

[15] H. L. Montgomery. The pair correlation of zeros of the zeta function. In: Analytic Number Theory, Proc. Sympos. Pure Math. 24, Amer. Math. Soc. (1973), 181–193.